# ASPIRIN- aspirin tablet Trifecta Pharmaceuticals USA LLC

-----

Aspirin 325mg (NSAID) \*nonsterhoidal anti-inflammatory drug

### **Active Ingredient**

Aspirin 325mg (NSAID)

Nonsteroidal anti-inflammatory drug

### **Purpose**

Pain Reliever / Fever Reducer

#### Do Not Use

- If you are allergic to aspirin or any other pain releiver / fever reducer
- If you have ever had an allergic reaction to this product or any of its ingredients

#### Uses

Temporarily relieves:

Headache

Pain and fever of colds

toothache

menstrual pain

muscle pain

minor pain of arthritis

### Warnings

### Reye's Syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behaviour with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a reare but serious illness.

### **Allergy Alert**

Aspirin may cause severe allergic reaction, which may include, facial swelling, hives, shock, asthma (wheezing).

### Stomach bleeding warning

This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you

- Have had stomach ulcers or bleeding problems
- Take blood thinning (anticogulant) or steroid drug
- Are age 60 or older
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- Take more or for a longer time than directed
- Have 3 or more alcoholic drinks every day while using this product

#### Ask a doctor before use if

- Stomach bleeding warning applies to you.
- You have a history of stomach problems, such as heartburn
- You have asthma
- You have high blood pressure, heart disease, liver cirrhosis or kidney disease.
- You are taking a diuretic

### Ask a doctor or pharmacist before use if you are

taking a prescription drug for gout, diabetes, arthritis

#### **Directions**

Do not take more than directed. See overdose warning

Drink a full glass of water with each dose

Adults and children 12 years and older

• Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor.

Children under 12 years

• Do not use unless directed by a doctor

### Stop Use and ask a doctor if

- An allergic reaction occurs. Seek medical help right away
- You experience any of the following signs of stomach bleeding, feel faint, vomit blood, have bloody stools, have stomach pain that does not get better
- Ringing in the ears or loss of hearing occurs, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days.
- New symptoms occur,
- Redness or swelling is present
- Thse could be signs of a serious condition

### If pregnant or breast-feeding

Ask a health professional before use. It is especially important not to use aspirin at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

#### Other Information

- Store at 25°C (77°F)
- Excursions permitted between 15°C-30°C (59°F-86°F)
- See end flap for expiration date and lot number
- Use by expiration date on package

### **Inactive Ingredients**

Corn Starch, Hyperomelose, Polyethylene glycol, propylene glycol

### Keep out of reach of children

In case of overdose, get medical help or contact a poison control center right away.

#### **Questions or Comments?**

Call 1-888-296-9067

### **Distributed By:**

Distributed By:

Trifecta Pharmaceuticals USA®

Ft. Lauderdale, FL. 33301 USA

1-888-296-9067

\*This product is not manufactured or distributed by the Bayer Healthcare LLC. owner of the registered trademark Bayer®

Reorder No. 2023

Made in the USA

#### Label

#### **OUTSID BOX**



#### **INNER PACKET**





### **ASPIRIN**

aspirin tablet

<b>—</b>			rmation
Prod	IIICT.	INTO	rmation

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69396-158

**Route of Administration** ORAL

### **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN (UNII: O8232NY3SJ)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)			

Product Characteristics			
Color	white	Score	score with uneven pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44157
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:69396-158- 01	50 in 1 BOX	08/24/2024		
1		2 in 1 PACKET; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	08/24/2024		

## **Labeler -** Trifecta Pharmaceuticals USA LLC (079424163)

Revised: 9/2024 Trifecta Pharmaceuticals USA LLC