

**ZANTAC 360- famotidine tablet, film coated**  
**Select Corporation**

-----  
**Zantac 360<sup>o</sup>™**

***Drug Facts***

**Active ingredient (in each tablet)**

Famotidine USP 20 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy Alert**

Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- read the directions and warnings before use
- store at 20-25°C (68-77°F)
- protect from moisture

**Inactive ingredients**

carnauba wax, corn starch, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, red iron oxide, sodium starch glycolate, talc, titanium dioxide, yellow iron oxide

**Questions or Comments?**

call **1-800-633-1610** or visit **www.zantacotc.com**

Product Repackaged and Distributed  
with permission of manufacturer by:  
**Select Corporation** Carrollton, TX 75007

**PRINCIPAL DISPLAY PANEL - 1 Tablet Pouch Blister Pack**

NEW FORMULA  
MAXIMUM STRENGTH  
See New Warnings

Zantac  
360°™  
Famotidine Tablets USP 20 mg/Acid Reducer

JUST ONE TABLET  
PREVENTS  
& RELIEVES

HEARTBURN  
DUE TO ACID  
INDIGESTION

1 TABLET  
Actual Size

2111926095.A



**MAXIMUM STRENGTH**  
**Zantac**  
 Famotidine Tablets USP 20 mg/Acid Reducer **360°**



**Single - Pack: 1 Tablet**

Product Repackaged and Distributed  
 with permission of manufacturer by:  
 Select Corporation Carrollton, TX 75007

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Famotidine USP 20 mg.....	Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy Alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain • frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting • stomach pain • kidney disease

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). ➔

**Drug Facts (continued)**

**Directions**

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- read the directions and warnings before use
- store at 20-25°C (68-77°F) • protect from moisture

**Inactive ingredients**

carnauba wax, corn starch, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, red iron oxide, sodium starch glycolate, talc, titanium dioxide, yellow iron oxide

**Questions or Comments?** call 1-800-633-1610 or visit [www.zantacotc.com](http://www.zantacotc.com)

Product manufactured for:  
 Chattem, Inc., a Sanofi Company  
 P. O. Box 2219  
 Chattanooga, TN 37409-0219  
 Made in India

**Now with Famotidine**

Just one tablet prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime

- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

2111926095

## ZANTAC 360

famotidine tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:52904-926(NDC:41167-0360)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
famotidine (UNII: 5QZO15J2Z8) (famotidine - UNII:5QZO15J2Z8)	famotidine	10 mg

### Inactive Ingredients

Ingredient Name	Strength
carnauba wax (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
magnesium stearate (UNII: 70097M6I30)	
microcrystalline cellulose (UNII: OP1R32D61U)	
ferric oxide red (UNII: 1K09F3G675)	
talc (UNII: 7SEV7J4R1U)	
titanium dioxide (UNII: 15FIX9V2JP)	
hydroxypropyl cellulose (1600000 WAMW) (UNII: RFW2ET671P)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
hypromellose 2910 (5 mpa.s) (UNII: R75537T0T4)	

### Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	ROUND (biconvex)	<b>Size</b>	5mm
<b>Flavor</b>		<b>Imprint Code</b>	CC;58
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC 52904-926			

1	NDC:52904-920-04	1 in 1 BLISTER PACK	04/12/2021	
1		1 in 1 POUCH; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA206531		04/12/2021	

**Labeler** - Select Corporation (053805599)

Revised: 4/2022

Select Corporation