### ZICLOVIN ORAL PAIN RELIEF- benzalkonium chloride, benzocaine gel Pharmadel LLC

-----

# Ziclovin Oral Pain Relief Gel (K)

# **Drug Facts**

# Active ingredients & Purposes

Active ingredients	Purposes
Benzalkonium chloride 0.1%	Topical antiseptic
Benzocaine 20%	Fever blister and cold sore treatment

#### Uses

For the temporary relief of pain due to

- fever blisters
- cold sores
- antiseptic to help prevent infection

# Warnings

For external use only. Avoid contact with the eyes.

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:** do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

#### Do not use

- longer than 1 week
- on large areas of the body
- in the eyes

#### Ask a doctor before use

• in case of deep or puncture wounds, animal bites, or serious burns

### When using this product

 you may feel a brief stinging sensation after application that will go away shortly thereafter

# Stop use and ask a dentist or doctor if

- a rash or irritation develops
- condition persists or worsens
- symptoms persist for more than 7 days or clear up and occur again within a few days

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

#### **Directions**

- adults and children 2 years of age and older: Clean the affected area. Apply a small amount to affected not more than 3 times daily. May be covered with a sterile bandage. If bandaged, let first dry.
- children under 2 years of age: do not use, consult a doctor

#### Other information

• store between 68-77°F (20-25°C)

# Inactive ingredients

ammonium glycyrrhizate, carboxymethylcellulose sodium, FD&C yellow #6, glycerin, polyethylene glycol, saccharin sodium, sorbic acid, water

#### **Questions?**

Dist. by/por: PHARMADEL LLC

New Castle, DE 19720

Questions? +1-866-359-3478

NDC 55758-438-01

Ziclovin

Gel para el alivio instantáneo del dolor oral

Tratamiento para ampollas febriles y herpes labial

1oz (28g)



**Instant Oral Pain Relief Gel** 

Fever blister and cold sore treatment

Tratamiento para las ampollas febriles y el herpes labial



Fever blister and cold sore treatment

### ZICLOVIN ORAL PAIN RELIEF

benzalkonium chloride, benzocaine gel

**Product Information** 

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55758-438
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	20 g in 100 g	
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII: 7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.1 g in 100 g	

Inactive Ingredients		
Ingredient Name	Strength	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
SORBIC ACID (UNII: X045WJ989B)		
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)		
GLYCERIN (UNII: PDC6A3C0OX)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
WATER (UNII: 059QF0KO0R)		

F	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55758-438- 01	1 in 1 CARTON	11/12/2024	
1		28 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	11/12/2024	

# Labeler - Pharmadel LLC (030129680)

Revised: 11/2024 Pharmadel LLC