

**LAXATIVE- bisacodyl tablet, delayed release**  
**Walgreen Company**

-----  
**Walgreens 44-676A**

***Active ingredient (in each tablet)***

Bisacodyl USP, 5 mg

***Purpose***

Stimulant laxative

***Uses***

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

***Warnings***

**Do not use**

if you cannot swallow without chewing.

**Ask a doctor before use if you have**

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

**When using this product**

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness, and cramps

**Stop use and ask a doctor if**

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

***Inactive ingredients***

carmines, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

***Questions or comments?***

**1-800-426-9391**

***Principal display panel***

***Walgreens***

Walgreens Pharmacist Recommended†

NDC 0363-6760-01

**Laxative Tablets**

BISACODYL USP, 5 mg / STIMULANT LAXATIVE

- Gentle, dependable overnight relief

**30**

ENTERIC -  
COATED  
TABLETS

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

50844

REV0923A67601

†Our pharmacists recommend the Walgreens brand.

DISTRIBUTED BY: **WALGREEN CO.**

**DEERFIELD, IL 60015**

**100% SATISFACTION GUARANTEED**

**walgreens.com**

©2024 Walgreen Co.



**Walgreens**



# Laxative Tablets

BISACODYL USP, 5 mg / STIMULANT LAXATIVE

- Gentle, dependable overnight relief

**30**  
ENTERIC-COATED  
TABLETS



ACTUAL SIZE

Do not print Area  
Lot & Exp. Date

W3OR60821-F  
REV-0924

50844 REV0923A67601

ITEM 504569 W00000-0000-0  
NDC 0363-6760-01

3 11917107350 7

**Drug Facts (continued)**

cellulose, polydextrose, polyethylene glycol, shellac wax, stearic acid, talc, titanium dioxide, tracetin, triethyl citrate

**Questions or comments?** 1-800-426-9391

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

**Drug Facts (continued)**

**Directions**

- do not take more than directed
- take with a glass of water
- adults and children 12 years and over: take 1 to 3 tablets in a single daily dose
- children 6 to under 12 years: take 1 tablet in a single daily dose
- children under 6 years: ask a doctor

**Other information**

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

**Inactive ingredients**

carne, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline

**Drug Facts (continued)**

**Warnings**

- Do not use if you cannot swallow without chewing.
- Ask a doctor before use if you have a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

**When using this product**

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness, and cramps

**Stop use and ask a doctor if**

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding, ask a health professional

Our pharmacists recommend the Walgreens brand.

DISTRIBUTED BY: WALGREEN CO.  
DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED  
walgreens.com  
©2024 Walgreen Co.

how2recycle.info

PAPER BOX TRAY

B-2201-676A-01-HR  
REV/0923A67601

**Drug Facts (continued)**

**Uses**

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

**Drug Facts**

**Active ingredient (in each tablet)**

Bisacodyl USP, 5 mg

**Purpose**

Stimulant laxative

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

# LAXATIVE

bisacodyl tablet, delayed release

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-6760
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISACODYL</b> (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>FD&amp;C RED NO. 40 ALUMINUM LAKE</b> (UNII: 6T47AS764T)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>METHACRYLIC ACID</b> (UNII: 1CS02G8656)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	

## Product Characteristics

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	B
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC 0363			

1	NDC:0363-6760-01	2 in 1 CARTON	09/16/2015	
1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-6760-52	6 in 1 CARTON	09/16/2015	
2		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	09/16/2015	

**Labeler** - Walgreen Company (008965063)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-6760)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-6760) , pack(0363-6760)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-6760)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-6760)

Revised: 2/2026

Walgreen Company