

TRILOGEL- lidocaine hydrochloride gel
Trifluent Pharma LLC

Trilogel™

Drug Facts

Active Ingredient

Lidocaine HCl, USP 2.8%

Purpose

Topical analgesic

Uses

For the temporary relief of pain associated with minor burns, sunburn, minor cuts, scrapes, insect bites, and minor skin irritations.

Warnings

For external use only

Do not use in large quantities, particularly over raw surfaces or blistered areas.

When using this product avoid contact with eyes

Stop use and ask a physician if condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a physician.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) immediately.

Directions

- Adults and children 2 years of age and older: Apply to affected area not more than 3 to 4 times daily.
- Children under 2 years of age: Do not use; consult a physician.

Other Information

Store at 20-25°C (68-77°F). Protect from freezing.

Inactive Ingredients

Caprylyl Glycol, Carica Papaya (Papaya) Fruit Extract, Disodium EDTA, Ethylhexylglycerin, Glycerin, Hexylene Glycol, Hydroxypropylcellulose, Phenoxyethanol, Purified Water, Sodium Lactate, Xanthan Gum.

Questions or Comments?

(210) 944-6920

PRINCIPAL DISPLAY PANEL - 57 G Pouch Label

Tear at notch to open. Do not use if seal is broken.

NDC: 73352-620-01

TRIFLUENT
PHARMA™

Trilogel™

Lidocaine HCl 2.8%

Topical Analgesic Gel

For the temporary relief of pain associated with minor burns, sunburn, minor cuts, scrapes, insect bites, and minor skin irritations.

NET WT. 2 OZ. (57 G)

Tear at notch to open. Do not use if seal is broken.

NDC: 73352-620-01



Trilogel™

Lidocaine HCl 2.8%

Topical Analgesic Gel

For the temporary relief of pain associated with minor burns, sunburn, minor cuts, scrapes, insect bites, and minor skin irritations.

NET WT. 2 OZ. (57 G)

Drug Facts

Active Ingredient**Purpose**
Lidocaine HCl, USP 2.8%.....Topical analgesic

Drug Facts (continued)

Inactive Ingredients
Caprylyl Glycol, Carica Papaya (Papaya) Fruit Extract, Disodium EDTA, Ethylhexylglycerin, Glycerin, Hexylene

Uses

For the temporary relief of pain associated with minor burns, sunburn, minor cuts, scrapes, insect bites, and minor skin irritations

Warnings**For external use only**

Do not use in large quantities, particularly over raw surfaces or blistered areas.

When using this product avoid contact with eyes

Stop use and ask a physician if condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a physician.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) immediately.

Directions

- Adults and children 2 years of age or older: Apply to affected area not more than 3 to 4 times daily.
- Children under 2 years of age: Do not use; consult a physician.

Other Information

Store at 20-25°C (68-77°F). Protect from freezing.

Glycol, Hydroxypropylcellulose, Phenoxyethanol, Purified Water, Sodium Lactate, Xanthan Gum.

Questions/Comments?

(210) 944-6920

MANUFACTURED FOR:

Trifluent Pharma, LLC
San Antonio, TX 78213

MANUFACTURED IN THE U.S.A.

620-01
REV01 05/2024

**TRIOGEL**

lidocaine hydrochloride gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73352-620
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE	28 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
XANTHAN GUM (UNII: TTV12P4NEE)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
PAPAYA (UNII: KU94FIY6JB)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
CAPRYLYL GLYCOL (UNII: 00YIU5438U)	
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	
HEXYLENE GLYCOL (UNII: KEH0A3F75J)	
SODIUM LACTATE (UNII: TU7HW0W0QT)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73352-620-01	1 in 1 POUCH	06/24/2024	
1		57 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M017	06/24/2024	

Labeler - Trifluent Pharma LLC (117167281)

Revised: 6/2024

Trifluent Pharma LLC