HEALCOME TATTOO NUMBING CREAM 5% LIDOCAINE- lidocaine cream YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD

Drug Facts

Drug Facts

Active ingredients Lidocaine 5%

Purpose Topical Anesthetic

Uses

- Temporarily relieves pain, burning, and soreness - Commonly used for tattooing, waxing, piercing, injection, microneedling, and hair removal

Warnings

For external use only

Do not use

- in large quantities - on wounds or on irritated or damaged skin - with a heating pad

When using this product

- avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes - use only as directed - do not bandage tightly - avoid contact with eyes and mucous membranes - do not expose the area to local heat or to direct sunlight

Stop and ask a doctor if discomfort persists.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of accidental ingestion, seek medical advice immediately or contact a Poison Control Center right away.

Directions

Patch Test Warning: Before first use, apply a small amount of cream to a small area of skin and wait 24 hours. If no adverse reaction occurs, proceed with regular use.
Clean and dry skin with mild soap.
Apply a thick, even layer with brush.
Wrap in plastic for 30 minutes.
Wipe off with tissue before starting the procedure.
Numbness duration may vary with temperature and skin type. If it's weak after 30 minutes, extend it to 60 minutes.

Children under 12 years of age: ask a doctor

Other information Store at room temperature, away from heat and moisture. ALWAYS CLOSE CAP TIGHTLY AFTER USE

Inactive ingredients Polyethylene Glycol, Soybean Lecithin, .Alpha.-Tocopherol Acetate, Water, Carbomer 934, Benzyl Alcohol

Active ingredients

Lidocaine 5%

Topical Anesthetic

Uses

- Temporarily relieves pain, burning, and soreness - Commonly used for tattooing, waxing, piercing, injection, microneedling, and hair removal

Do not use

- in large quantities - on wounds or on irritated or damaged skin - with a heating pad

When using this product

- avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes use only as directed - do not bandage tightly - avoid contact with eyes and mucous membranes - do not expose the area to local heat or to direct sunlight

Stop and ask a doctor if discomfort persists.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of accidental ingestion, seek medical advice immediately or contact a Poison Control Center right away.

Warnings

For external use only

Directions

- Patch Test Warning: Before first use, apply a small amount of cream to a small area of skin and wait 24 hours. If no adverse reaction occurs, proceed with regular use. - Clean and dry skin with mild soap. - Apply a thick, even layer with brush. - Wrap in plastic for 30 minutes. - Wipe off with tissue before starting the procedure. - Numbness duration may vary with temperature and skin type. If it's weak after 30 minutes, extend it to 60 minutes.

Children under 12 years of age: ask a doctor

Other information Store at room temperature, away from heat and moisture. ALWAYS CLOSE CAP TIGHTLY AFTER USE

Inactive ingredients Polyethylene Glycol, Soybean Lecithin, .Alpha.-Tocopherol Acetate, Water, Carbomer 934, Benzyl Alcohol

| HEALCOME TATTOO NUMBING CREAM 5% LIDOCAINE | | | | | | | | |
|--|--|---|--|--|--|--|--|--|
| | | | | | | | | |
| Product Information | | | | | | | | |
| HUMAN OTC DRUG | Item Code (Sour | NDC:83364-010 | NDC:84010-020) | | | | | |
| TOPICAL | | | | | | | | |
| | | | | | | | | |
| Active Ingredient/Active Moiety | | | | | | | | |
| Ingredient Name | | Basis of Strength | Strength | | | | | |
| LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987) | | OCAINE | 5 g in 100 g | | | | | |
| | | | | | | | | |
| | HUMAN OTC DRUG TOPICAL Moiety ient Name | HUMAN OTC DRUG Item Code (Sourd TOPICAL Moiety ient Name | HUMAN OTC DRUG Item Code (Source) NDC:83364-010(TOPICAL Moiety ient Name Basis of Strength | | | | | |

| Inactive Ingredients | | | | | | |
|---|--------------------------|---------------------|-------------------------|-----------------------|--|--|
| | | Strength | | | | |
| PF | | | | | | |
| C/ | ARBOMER 934 (U | NII: Z135WT9208) | | | | |
| .ALPHATOCOPHEROL ACETATE (UNII: 9E8X80D2L0) | | | | | | |
| w | WATER (UNII: 059QF0KO0R) | | | | | |
| BENZYL ALCOHOL (UNII: LKG8494WBH) | | | | | | |
| SOYBEAN LECITHIN (UNII: 1DI56QDM62) | | | | | | |
| | | | | | | |
| | | | | | | |
| Packaging | | | | | | |
| # | ltem Code | Package Description | Marketing Start Date | Marketing End Date | | |
| 1 | NDC:83364-010- 01 | 1 in 1 BOX | 09/23/2024 | | | |

| Marketing Information | | | | | | |
|-----------------------|---|-------------------------|-----------------------|--|--|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | | | |
| OTC Monograph Drug | M017 | 09/23/2024 | | | | |

Labeler - YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD (725220463)

60 g in 1 BOTTLE; Type 0: Not a Combination Product

Revised: 9/2024

1

YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD