

**NASAL DECONGESTANT- pseudoephedrine hcl tablet, film coated
H E B**

HEB 44-112

Active ingredient (in each tablet)

Pseudoephedrine HCl 30 mg

Purpose

Nasal decongestant

Uses

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves sinus congestion and pressure

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- diabetes
- heart disease
- high blood pressure
- thyroid disease
- difficulty in urination due to enlargement of the prostate gland

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with fever

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years and over	take 2 tablets every 4 to 6 hours; do not take more than 8 tablets in 24 hours
children ages 6 to 11 years	take 1 tablet every 4 to 6 hours; do not take more than 4 tablets in 24 hours
children under 6 years	do not use

Other information

- **each tablet contains:** calcium 15 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, silicon dioxide, titanium dioxide, triacetin

Questions or comments?

1-800-426-9391

Principal display panel

Compare to Sudafed® Sinus Congestion active ingredient*

NDC 37808-112-22

H - E - B ®

**Maximum Strength
Nasal Decongestant**

Pseudoephedrine HCl 30 mg /
Nasal Decongestant

Sinus Pressure & Congestion

Non-Drowsy

Relief of:

- **Nasal & Sinus Congestion**
- **Sinus Pressure**

actual size

48 TABLETS, 30 mg EACH

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Sudafed® Sinus Congestion.
50844 REV0619B11222

100%
GUARANTEE
promise

**If you aren't completely pleased
with this product, we'll be happy to
replace it or refund your money.
You have our word on it.**

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TX 78204



This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Sudafed® Sinus Congestion. 50844 REV0619B11222

MADE WITH PRIDE AND CARE FOR H-E-B® SAN ANTONIO, TX 78204

HE-B
If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.
100% GUARANTEE
Pseudoephedrine

Drug Facts (continued)

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient (in each tablet)
Pseudoephedrine HCl 30 mg. Nasal decongestant

Purpose
temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Do not use if you have heart disease, diabetes, high blood pressure, thyroid disease, difficulty in urination due to enlargement of the prostate gland.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ symptoms do not improve within 7 days or occur with fever

If pregnant or breast-feeding, ask a health professional before use.

Other information
■ each tablet contains: calcium 15 mg
■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number

Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, silicon dioxide, titanium dioxide, triacetin

Questions or comments? 1-800-426-9391

Directions

adults and children 12 years and over	take 2 tablets every 4 to 6 hours; do not take more than 8 tablets in 24 hours
children ages 6 to 11 years	take 1 tablet every 4 to 6 hours; do not take more than 4 tablets in 24 hours
children under 6 years	do not use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No Print/No Varnish Area
Lot & Exp Date



Nasal Decongestant

Pseudoephedrine HCl 30 mg / Nasal Decongestant

Sinus Pressure & Congestion

Maximum Strength
Non-Drowsy

Compare to Sudafed® Sinus Congestion active ingredient*



Maximum Strength Nasal Decongestant

Pseudoephedrine HCl 30 mg /
Nasal Decongestant

Sinus Pressure & Congestion

Non-Drowsy

Relief of:

- Nasal & Sinus Congestion
- Sinus Pressure



actual size

48 TABLETS, 30 mg EACH

NDC 37808-112-22

HEB 44-112

NASAL DECONGESTANT
pseudoephedrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-112
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	44;112
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-112-22	2 in 1 CARTON	08/25/1981	
1		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/25/1981	

Labeler - H E B (007924756)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(37808-112)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(37808-112) , pack(37808-112)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(37808-112)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(37808-112)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(37808-112)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(37808-112)