

TETROXY HCA- oxytetracycline hcl powder Bimeda, Inc.

Tetroxy® HCA-1400 (oxytetracycline hydrochloride soluble powder)

Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

This packet contains 512 g of oxytetracycline hydrochloride and will make:
2560 gallons (9690 liters) containing 200 mg oxytetracycline hydrochloride per gallon
1280 gallons (4845 liters) containing 400 mg oxytetracycline hydrochloride per gallon
640 gallons (2422 liters) containing 800 mg oxytetracycline hydrochloride per gallon
This packet will treat 51,200 lbs of swine.
This packet will treat 20,480 lbs of turkey at 25 mg/lb body weight.

For Use in Drinking Water Only
Not For Use in Liquid Medicated Feeds
Not For Use in Humans
Keep Out of Reach of Children
HAZARDOUS REMEDY

See back panel for instructions.
Approved by FDA under ANADA # 200-144

Net Weight: 3.09 lb (1400 g)

MIXING DIRECTIONS FOR WATER MEDICATION: CHICKENS - TURKEYS - SWINE

Dosage and indications for use in water proportioners: Add one packet to the following amount of water to make a stock solution for use in a proportioner set to meter at the rate of 1 ounce per gallon.

	Disease	Treatment Level	Gallons Stock/Pack
CHICKENS	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200-400 mg	20-10
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400-800 mg	10-5
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400-800 mg	10-5
TURKEYS	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200-400 mg	20-10
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	10
	GROWING TURKEYS - Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
SWINE	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE: Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption
	FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospirae) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.		

USE DIRECTIONS: Mix fresh solutions daily. Use as sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys.

NOTE: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperature and humidity, each of which affects water consumption.

CAUTION: Use as the sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.

WARNING: Do not feed to birds producing eggs for human consumption.

RECOMMENDED PACKET STORAGE CONDITIONS: Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

CONTACT INFORMATION: To report adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

NDC 61133-5010-2

TETROXY[®] HCA-1400

(oxytetracycline hydrochloride soluble powder)

Antibiotic

CAUTION: Federal (USA) law restricts this drug product to use by or on the order of a licensed veterinarian.

This packet contains 512 grams of oxytetracycline hydrochloride and will make:
2560 gallons (9690 liters) containing 200 mg oxytetracycline hydrochloride per gallon
1280 gallons (4845 liters) containing 400 mg oxytetracycline hydrochloride per gallon
640 gallons (2422 liters) containing 800 mg oxytetracycline hydrochloride per gallon
This packet will treat 51,200 lbs of swine.
This packet will treat 20,480 lbs of turkey at 25 mg/lb body weight.

For Use in Drinking Water Only
Not for Use in Liquid Medicated Feeds
Not for Use in Humans
Keep Out of Reach of Children
HAZARDOUS REMEDY

See back panel for instructions

Approved by FDA under ANADA # 200-144

Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com



Net Weight: 3.09 lb (1400 g)

Bimeda

TETROXY[®] HCA-1400

(oxytetracycline hydrochloride soluble powder)

MIXING DIRECTIONS FOR WATER MEDICATION: CHICKENS - TURKEYS - SWINE

Dosage and indications for use in water preparations: Add one packet to the following amount of water to make a stock solution for use in a proportioner set to meter at the rate of 1 ounce per gallon.

	Disease	Treatment Level	Gallons Stock/Pack
CHICKENS	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200-400 mg	20-10
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400-600 mg	10-5
	Control of fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline.	400-600 mg	10-5
TURKEYS	Control of heartitis caused by <i>Haemophilus meleagridis</i> , susceptible to oxytetracycline.	200-400 mg	20-10
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	10
SWINE	CRYING TURKEYS - Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronavirus enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE: Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella enteritidis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline. FOR BREEDING SWINE: Leptospirosis reducing the incidence of abortions and shedding of leptospira caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline.	10 mg/lb body weight	Varies with age & water consumption

USE DIRECTIONS: Mix fresh solutions daily. Use as sole source of drinking water. Do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. As a generalization, 200 chickens will drink 1 gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer up to 5 days to swine and 7 to 14 days for chickens and turkeys.

NOTE: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperature and humidity, each of which affects water consumption.

CAUTION: Use as the sole source of oxytetracycline. Not to be used for more than 5 consecutive days in swine or 14 consecutive days in chickens and turkeys.

WARNING: Do not feed to birds producing eggs for human consumption.

RECOMMENDED PACKET STORAGE CONDITIONS: Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

CONTACT INFORMATION: To report adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Bimeda, Inc. at 1-888-524-6352. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-9ETS or <http://www.fda.gov/reportarmls>



Tetroxy[®] is a registered trademark of Bimeda, Inc.
11ET014 BIE1006 Rev. 04/24

Bimeda

TETROXY HCA
oxytetracycline hcl powder

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:61133-5010
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYTETRACYCLINE HYDROCHLORIDE (UNII: 4U7K4N52ZM) (OXYTETRACYCLINE ANHYDROUS - UNII:SLF0D9077S)	OXYTETRACYCLINE HYDROCHLORIDE	648 g in 1772 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61133-5010-1	280 g in 1 POUCH		
2	NDC:61133-5010-2	1400 g in 1 POUCH		
3	NDC:61133-5010-3	1772 g in 1 POUCH		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200144	07/02/2004	

Labeler - Bimeda, Inc. (060492923)

Registrant - Bimeda, Inc. (060492923)

Establishment

Name	Address	ID/FEI	Business Operations
Bimeda, Inc.		060492923	manufacture

Revised: 12/2024

Bimeda, Inc.