

**TESALON TESAPERLAS SEVERE COLD MAX RELIEF- dextromethorphan hbr,
guaifenesin capsule, liquid filled
OPMX LLC**

TESALON Tesaperlas Severe Cold Max Relief

Drug Facts

Active ingredients

Active Ingredients (in each softgel)

Active Ingredients (in each softgel)

Dextromethorphan HBr 20 mg
Guaifenesin 400 mg

Purpose

Cough suppressant
Expectorant

Uses

- helps loosen phlegm (mucus)
- helps thin bronchial secretions to make coughs more productive

Warnings

Warnings

Do not use:

- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.
- If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if:

- cough lasts more than 7 days, recurs, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding:

- ask a health professional before use.

Keep out of reach of children

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:

- Do not take more than directed- **See Overdose warning**
- Do not use for more than 7 days.
- Do not take more than 6 softgels in 24 hours

Adults and children 12 years of age & over	Take 1 softgel every 4 hours with a full glass of water while symptoms persist
Children under 12 years of age	Do not use

Other information:

- Store at room temperature between 20-25°C (68-77°F)
- Avoid excessive heat and humidity

Inactive Ingredients:

FD&C yellow No. 10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitan solution, titanium dioxide.

Questions or comments?

Call: **(619) 600-5632** (Mon-Fri 9am-5pm EST)

NDC 69729-788-20

Tesalon®

Tesaperlas

Adults

Expectorant & Cough Suppressant

Expectorante & Supresor de la Tos

Controls cough

Controla la tos

Thins & loosens mucus

Adelgaza y expulsa las flemas

SOFTGELS

Dextromethorphan HBr 20 mg / Guaifenesin 400 mg

20 Softgels Box

Read & keep package for complete product information

TAMPER EVIDENT; DO NOT USE IS PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING



Exclusively distributed by:

 Chula Vista, CA 91910
 Phone: 619-600-5632

Made for:


VERSION EN ESPAÑOL EN EL INTERIOR DE LA CAJA

Call: (619) 600-5632 (Mon-Fri 9am-5pm EST)

Questions or comments?
 P&G yellow #10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Inactive ingredients:
 ■ Avoid excessive heat and humidity
 ■ Store at room temperature between 20-25°C (68-77°F)

Other Information:

Children	Do not use
12 years of age & over	Take 1 softgel every 4 hours with a full glass of water while symptoms persist.

Directions:
 ■ Do not take more than 6 softgels in 24 hours
 ■ Do not use for more than 7 days
 ■ Do not take more than directed. See Overdose warning

Drug Facts (continued)

Drug Facts

Active ingredients (in each softgel)
 Dextromethorphan HBr 20 mg
 Guaifenesin 400 mg

Purpose
 Cough suppressant
 Expectorant

Uses:
 ■ Helps loosen phlegm (mucus)
 ■ Helps thin bronchial secretions to make coughs more productive

Warnings:
 ■ Do not use:
 ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.
 ■ If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:
 ■ Cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ Cough that occurs with too much phlegm (mucus)

Stop and ask a doctor if:
 ■ Cough lasts for more than 7 days, recurs, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding: ask a health professional before use.

Keep out of reach of children

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

TAMPER EVIDENT:
 Do not use if package is
 opened or if blister unit is
 torn, broken or shows
 any sign of tampering



REV 09/2024
 ITEM # RX13144



Adults

Tesaperlas

Expectorant & Cough Suppressant

Dextromethorphan HBr 20 mg Guaifenesin 400 mg

NDC 69729-788-20

Lot No.:
Exp. Date:

NON VARNISH AREA

Read & keep outer package for complete product information



Adults

Tesaperlas

Expectorant & Cough Suppressant

Expectorante & Supresor de la Tos

- Controls cough
Controla la tos
- Thins & loosens mucus
Adelgaza y expulsa las flemas



SOFTGELS

Dextromethorphan HBr 20 mg / Guaifenesin 400 mg

20 Softgels Box



Adults

Tesaperlas

Expectorante & Supresor de la Tos

Dextromethorphan HBr 20 mg Guaifenesin 400 mg

REV. 09.2024
ITEM # RRX1144

8

50043

16912

3

TAMPER EVIDENT:
Do not use if package is
opened or if blister unit is
torn, broken or shows
any sign of tampering

Drug Facts

Active Ingredients (in each softgel)

Dextromethorphan HBr 20 mgCough suppressant
Guaifenesin 400 mgExpectorant

Purpose

Uses:

- helps loosen phlegm (mucus)
- helps thin bronchial secretions to make coughs more productive

Warnings:

Do not use:

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.
- if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if:

- cough lasts for more than 7 days, recurs, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding: ask a health professional before use.

Keep out of reach of children

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions:

- Do not take more than directed - See **Overdose** warning
- Do not use for more than 7 days.
- Do not take more than 6 softgels in 24 hours

Adults and children 12 years of age & over	Take 1 softgel every 4 hours with a full glass of water while symptoms persist
---	--

Children under 12 years of age	Do not use
-----------------------------------	------------

Other Information:

- Store at room temperature between 20-25°C (68-77°F)
- Avoid excessive heat and humidity

Inactive Ingredients:

FD&C yellow #10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Questions or comments?

Call: (619) 600-5632 (Mon-Fri 9am-5pm EST)

VERSION EN ESPAÑOL EN EL INTERIOR DE LA CAJA

Tesalon
Adults
Tesaperlas

Made for:



Exclusively distributed by:



Chula Vista, CA 91910
Phone: 619-600-5632



TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING

Información del Fármaco

Ingredientes Activos (en cada cápsula)

Dextrometorfano HBr 20 mgSupresor de la tos
Guaifenesina 400 mgExpectorante

Propósito

Usos:

- Ayuda a aflojar la flema (moco)
- Ayuda a diluir las secreciones bronquiales para que la tos sea más productiva

Advertencias:

No utilizar:

- si está tomando ahora un inhibidor de la monoaminoxidasa (IMAO) de prescripción médica (ciertos medicamentos para la depresión, afecciones psiquiátricas o emocionales, o la enfermedad de Parkinson), o durante 2 semanas después de dejar el medicamento IMAO.
- si no sabe si su medicamento recetado contiene un IMAO, pregunte a un médico o farmacéutico antes de tomar este producto.

Consulte a un médico antes de usarlo si tiene:

- tos que se produce con demasiada flema (mucosidad)
- tos persistente o crónica como la que se produce con el tabaquismo, el asma, la bronquitis crónica o el enfisema.

Suspenda su uso y consulte a un médico si:

- La tos dura más de 7 días, se repite o va acompañada de fiebre, sarpullido o dolor de cabeza persistente. Estos podrían ser signos de una enfermedad grave.

Si está embarazada o en período de lactancia: consulte a un profesional de la salud antes de utilizarlo.

Mantener fuera del alcance de los niños

Información del Fármaco (continuación)

Advertencia de sobredosis: En caso de sobredosis, busque ayuda médica o póngase en contacto con un Centro de Control de Intoxicaciones de inmediato. La atención médica inmediata es fundamental tanto para los adultos como para los niños, incluso si no se observa ningún signo o síntoma.

Indicaciones:

- No tomar más de lo indicado - Ver **Advertencia de sobredosis**
- No usar durante más de 7 días.
- No tomar más de 6 cápsulas blandas en 24 horas.

Adultos y niños mayores de 12 años	Tomar 1 cápsula blanda cada 4 horas con un vaso lleno de agua mientras persistan los síntomas
---------------------------------------	---

Niños menores de 12 años	No utilizar
--------------------------	-------------

Otras informaciones:

- Almacenar a temperatura ambiente entre 20-25°C (68-77°F)
- Evitar el calor y la humedad excesivos

Ingredientes Inactivos:

FD&C amarillo #10, gelatina, glicerina, polietilenglicol, povidona, propilenglicol, agua purificada, solución de sorbitol y dióxido de titanio

¿Preguntas o comentarios?

Llame: (619) 600-5632 (de lunes a viernes de 9 a 17 horas EST)

A PRUEBA DE MANIPULACIONES: NO UTILIZAR SI EL ENVASE ESTÁ ABIERTO O SI EL BLÍSTER ESTÁ RASGADO, ROTO O PRESENTA ALGÚN SIGNO DE MANIPULACIÓN

TESALON TESAPERLAS SEVERE COLD MAX RELIEF

dextromethorphan hbr, guaifenesin capsule, liquid filled

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:69729-788

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg

Inactive Ingredients

Ingredient Name	Strength
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POVIDONE (UNII: FZ989GH94E)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
WATER (UNII: 059QF0KO0R)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	781
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-788-20	1 in 1 CARTON	09/27/2024	
1		20 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/27/2024	

Labeler - OPMX LLC (029918743)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Medgel Private Limited

677385498

manufacture(69729-788)

Revised: 1/2026

OPMX LLC