

**NO DRIP NASAL MIST- oxymetazoline spray liquid**  
**Walgreens Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Anefrin Nasal Spray Extra Moisturizing**

**Drug Facts**

**Active Ingredient**

Oxymetazoline Hydrochloride 0.05%

**Purpose**

Nasal Decongestant

**Uses**

Temporarily relieves nasal congestion due to:

- common cold
- hay fever
- upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- Shrinks swollen membranes so you can breathe more freely

**Warnings**

**Ask a Doctor before use if you have**

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to enlarged prostate gland.

**When using this product**

- **Do not use more than directed**
- do not use for more than three days, Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur.
- use of this container by more than one person may spread infection.

## stop use and ask a doctor if

symptoms persist.

## if pregnant or breast-feeding

ask a health professional before use.

## Keep out of the reach of children

if swallowed, get medical help or contact a Poison Control Center right away.

## Directions

- Before using for the first time, remove the protective cap from the tip and prime the pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostrile. Fully depress rim with a firm and even stroke, and sniff deeply.
- Wipe nozzle clean after use.
- Adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- Children under 6 years of age consult a doctor.

## Other information

- store between 20° and 25°C (68° to 77°F)
- retain carton for future reference on full labeling

## Inactive Ingredients

Benzalkonium Chloride Solution, Benzyl Alcohol, Edetate Disodium, Flavor, Glycerin, Microcrystalline Cellulose and Carboxymethylcellulose Sodium, Polyethylene Glycol, Povidone, Purified Water, Sodium Phosphate Dibasic, Sodium Phosphate Monobasic

## Principal Panel 0.5 FLOZ Bottle

The image shows the principal panel of a 0.5 FLOZ bottle of Anefrin Nasal Spray. The panel is blue and white with red and black text. It includes the Walgreens logo, product name, active ingredient, and various warnings and directions.

**Walgreens**  
NO DRIP NDC 0363-2307-52  
**Anefrin Nasal Spray**  
OXYMETAZOLINE HCl 0.05% /  
NASAL DECONGESTANT  
• 12 hour  
0.5 FL OZ (15 mL)

**SEE CARTON FOR FULL LABELING**  
**DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING**

**Active Ingredient:** Oxymetazoline hydrochloride 0.05%  
**Uses:** See Carton.  
**Warnings:** Ask a doctor before use if you have:  
• heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland.  
**When using this product:**  
• do not use more than directed • do not use for more than 3 days. Frequent or prolonged use may cause nasal congestion to recur or worsen. • temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur • use of this container by more than one person may spread infection.  
**Stop use and ask a doctor** if symptoms persist. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions:** To use: squeeze cap at base on opposite sides and twist cap counterclockwise to remove. Secure cap after use. • adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period. • children under 6 years of age: ask a doctor.

**Questions or comments?**  
**1-800-222-1087**  
DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
walgreens.com ©2018 Walgreen Co.  
ITEM 821847 W00000-0000-0

REV0319  
ORG0518-F

# Walgreen No Drip Nasal Mist

Extra Moisturizing

Nasal Spray

**Oxymetazoline HCL 0.05%**

**12 hour Nasal Spray**

Net Wt 0.5 FLOZ (15ml)

**Walgreens**  
NO DRIP NDC 0363-2307-52  
**Anefrin Nasal Spray**  
OXYMETAZOLINE HCl 0.05% /  
NASAL DECONGESTANT  
• 12 hour  
0.5 FL OZ (15 mL)

**SEE CARTON FOR FULL LABELING**  
**DO NOT USE IF PRINTED SEAL OVER  
CAP IS TORN OR MISSING**

**Active ingredient:** Oxymetazoline hydrochloride  
0.05% **Uses:** • See Carton.  
**Warnings:** Ask a doctor before use if you have  
• heart disease • high blood pressure • thyroid  
disease • diabetes • trouble urinating due to an  
enlarged prostate gland. **When using this product**  
• do not use more than directed • do not use for more  
than 3 days. Frequent or prolonged use may cause  
nasal congestion to recur or worsen. • temporary  
discomfort such as burning, stinging, sneezing or an  
increase in nasal discharge may occur • use of this  
container by more than one person may spread  
infection. **Stop use and ask a doctor** if symptoms  
persist. **If pregnant or breast-feeding,** ask a health  
professional before use. **Keep out of reach of  
children.** If swallowed, get medical help or contact a  
Poison Control Center right away.  
**Directions:** To use: squeeze cap at base on  
opposite sides and twist cap  
counterclockwise to remove. Secure cap  
after use. • adults and children 6 to under  
12 years of age (with adult supervision): 2 or  
3 sprays in each nostril not more often than  
every 10 to 12 hours. Do not exceed 2 doses  
in any 24-hour period. • children under 6  
years of age: ask a doctor.

**Questions or comments?**  
**1-800-222-1087**  
DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
walgreens.com ©2018 Walgreen Co.  
ITEM 821847 W00000-0000-0

REV0319  
ORG0518-F

**Walgreens**  
NO DRIP NDC 0363-2307-52  
**Anefrin Nasal Spray**  
OXYMETAZOLINE HCl 0.05% /  
NASAL DECONGESTANT  
• 12 hour  
0.5 FL OZ (15 mL)

**SEE CARTON FOR FULL LABELING**  
**DO NOT USE IF PRINTED SEAL OVER  
CAP IS TORN OR MISSING**

**Active ingredient:** Oxymetazoline hydrochloride  
0.05% **Uses:** • See Carton.  
**Warnings:** Ask a doctor before use if you have  
• heart disease • high blood pressure • thyroid  
disease • diabetes • trouble urinating due to an  
enlarged prostate gland. **When using this product**  
• do not use more than directed • do not use for more  
than 3 days. Frequent or prolonged use may cause  
nasal congestion to recur or worsen. • temporary  
discomfort such as burning, stinging, sneezing or an  
increase in nasal discharge may occur • use of this  
container by more than one person may spread  
infection. **Stop use and ask a doctor** if symptoms  
persist. **If pregnant or breast-feeding,** ask a health  
professional before use. **Keep out of reach of  
children.** If swallowed, get medical help or contact a  
Poison Control Center right away.  
**Directions:** To use: squeeze cap at base on  
opposite sides and twist cap  
counterclockwise to remove. Secure cap  
after use. • adults and children 6 to under  
12 years of age (with adult supervision): 2 or  
3 sprays in each nostril not more often than  
every 10 to 12 hours. Do not exceed 2 doses  
in any 24-hour period. • children under 6  
years of age: ask a doctor.

**Questions or comments?**  
**1-800-222-1087**  
DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
walgreens.com ©2018 Walgreen Co.  
ITEM 821847 W00000-0000-0

REV0319  
ORG0518-F

**Walgreens**  
NO DRIP NDC 0363-2307-52  
**Anefrin Nasal Spray**  
OXYMETAZOLINE HCl 0.05% /  
NASAL DECONGESTANT  
• 12 hour  
0.5 FL OZ (15 mL)

**SEE CARTON FOR FULL LABELING**  
**DO NOT USE IF PRINTED SEAL OVER  
CAP IS TORN OR MISSING**

**Active ingredient:** Oxymetazoline hydrochloride  
0.05% **Uses:** • See Carton.  
**Warnings:** Ask a doctor before use if you have  
• heart disease • high blood pressure • thyroid  
disease • diabetes • trouble urinating due to an  
enlarged prostate gland. **When using this product**  
• do not use more than directed • do not use for more  
than 3 days. Frequent or prolonged use may cause  
nasal congestion to recur or worsen. • temporary  
discomfort such as burning, stinging, sneezing or an  
increase in nasal discharge may occur • use of this  
container by more than one person may spread  
infection. **Stop use and ask a doctor** if symptoms  
persist. **If pregnant or breast-feeding,** ask a health  
professional before use. **Keep out of reach of  
children.** If swallowed, get medical help or contact a  
Poison Control Center right away.  
**Directions:** To use: squeeze cap at base on  
opposite sides and twist cap  
counterclockwise to remove. Secure cap  
after use. • adults and children 6 to under  
12 years of age (with adult supervision): 2 or  
3 sprays in each nostril not more often than  
every 10 to 12 hours. Do not exceed 2 doses  
in any 24-hour period. • children under 6  
years of age: ask a doctor.

**Questions or comments?**  
**1-800-222-1087**  
DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
walgreens.com ©2018 Walgreen Co.  
ITEM 821847 W00000-0000-0

REV0319  
ORG0518-F



## Principal display panel- Carton label

### Walgreens No Drip Nasal Mist

Extra Moisturizing

Nasal Spray

**Oxymetazoline HCL 0.05%**

**12 hour Nasal Spray**

Net Wt 0.5 FLOZ (15ml)



# Anefrin<sup>®</sup> Nasal Spray

ON DRIP

Walgreens

DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING

#### Drug Facts

**Active ingredient** Oxymetazoline hydrochloride 0.05% **Purpose** Nasal decongestant

**Uses** temporarily relieves nasal congestion due to:  
• common cold • hay fever • upper respiratory allergies • temporarily relieves sinus congestion and pressure • shrinks swollen nasal membranes so you can breathe more freely

#### Warnings

**Ask a doctor before use if you have** • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland

#### When using this product

• do not use more than directed  
• do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.  
• temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur  
• use of this container by more than one person may spread infection

**Stop use and ask a doctor if symptoms persist**

**if pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

Before using the first time, remove the protective cap from the tip and prime pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use.

• adults and children 6 to under 12 years of age: hold adult supervision; 2 or 3 sprays in each nostril not more often than every 12 to 12 hours. Do not exceed 2 doses in any 24-hour period.  
• children under 6 years of age: ask a doctor

#### Drug Facts (continued)

#### Other information

• store between 20° to 25°C (68° to 77°F)  
• refer carton for future reference to full labeling

#### Inactive ingredients

Borzoalkonium Chloride Solution, Benzyl Alcohol, Edetate Disodium, Flavor, Glycerin, Methylparaben, Methylcellulose and Carbonylmethylcellulose Sodium, Polyethylene Glycol, Purified Water, Sodium Phosphate Dibasic, Sodium Phosphate Monobasic

**IMPORTANT: KEEP OUTER CARTON FOR FUTURE REFERENCE ON FULL LABELING**

**Cap Removal Instructions:** To remove cap, squeeze cap at base on opposite sides on the two firm indentations with thumb while depressing, twist cap counterclockwise to remove. Press down to use. Refer to arrows on cap. To reapply cap, turn cap on pump clockwise until it locks.

Walgreens Pharmacist Recommended  
The Walgreens Pharmacy Survey  
This product is not manufactured or distributed by Walgreens Pharmacy, Inc., the distributor of Anefrin<sup>®</sup> No Drip Extra Moisturizing Pump.

**Questions or comments?**  
1-800-222-1167

DISTRIBUTED BY WALGREEN CO.  
2000 W. 10TH ST., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2010 Walgreen Co.

Walgreens

Compare to Anefrin<sup>®</sup> No Drip Extra Moisturizing Pump Mist active ingredient!

## NO DRIP Anefrin<sup>®</sup> Nasal Spray

NDC 0363-2301-02

OXYMETAZOLINE HCl 0.05% / NASAL DECONGESTANT

EXTRA MOISTURIZING

12 HOUR

- Won't drip from nose or down throat
- Helps relieve congestion

0.5 FL OZ (15 mL)

ACTUAL SIZE

LOT

EXP

ITEM 821947 W00000-0000-0



0H00019-F



## NO DRIP NASAL MIST

oxymetazoline spray liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-2307
<b>Route of Administration</b>	NASAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OXYMETAZOLINE HYDROCHLORIDE</b> (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	OXYMETAZOLINE HYDROCHLORIDE	50 mg in 100 mL

### Inactive Ingredients

Ingredient Name	Strength

<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL 1000</b> (UNII: U076Q6Q621)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>SODIUM PHOSPHATE, MONOBASIC</b> (UNII: 3980JIH2SW)	
<b>SODIUM PHOSPHATE, DIBASIC</b> (UNII: GR686LBA74)	
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-2307-52	1 in 1 CARTON	10/08/2018	
1		15 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/08/2018	

**Labeler** - Walgreens Company (008965063)

Revised: 10/2018

Walgreens Company