XL-3 DAY TIME- acetaminophen, dextromethorphan hbr, phenylephrine hcl liquid Rnv LLC

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Pain reliever/fever reducer

Cough Suppressant

Nasal Decongestant

Uses

temporarily relieves common cold and flu symptoms:

- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Sever liver damage may occur if

- adult takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adults has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause sever skin reactions. Symptoms may include:

- skin reddening
- blisters
- ras

If skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is sever, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly

Do not use

- with any other products containing acetaminophen (prescription or nonprescription).
 If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's diseas), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabete
- high blood pressure
- trouble urinating due to enlarged prostate gland
- cough accompanied by excessive phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product

- do not use more than directed (see overdose warning)
- avoid alcoholic drinks

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms get worse or last more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause serious health problemas including liver damange. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as children even if ou do not notice any signs or symptoms.

- take only as directed- see Overdose warning
- use only dosing cup provided
- keep dosing cup provided with the product
- do not exceed 4 doses per 24 hours
- if taking Day Time during the day and Night Time at night, limit the total to 4 doses per 24 hours
- TBSP=tablespoon
- mL=mililiter

adults & children 12 years and over	30 mL (2 TBSP) every 4 hours
children 6 to under 12 years	15 mL (1 TBSPO) every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

Other information

- each 15 mL contains: sodium 5 mg
- store between 58 86 degrees F (15-30 degrees C)

Inactive ingredients: Citric acid, FD&C Yellow No 6, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or Comments? 1-888-446-4753

Monday - Friday from 8 AM - 5 PM Eastern Standard Time





XL-3 DAY TIME

acetaminophen, dextromethorphan hbr, phenylephrine hcl liquid

Product Information	roduct Information		
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84379-441
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg in 15 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE	5 mg in 15 mL	

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID (UNII: 2968PHW8QP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

F	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84379-441- 06	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2024	
2	NDC:84379-441- 12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2024	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/01/2024	

Labeler - Rnv LLC (118917568)

Revised: 12/2024 Rnv LLC