

**XL-3 DAY TIME- acetaminophen, dextromethorphan hbr, phenylephrine
hcl liquid
Rnv LLC**

XL-3 Day Time

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Pain reliever/fever reducer

Cough Suppressant

Nasal Decongestant

Uses temporarily relieves common cold and flu symptoms:

- nasal congestion • cough due to minor throat and bronchial irritation
- sore throat • headache • minor aches and pains • fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adults has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days or fever unless directed by a doctor
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabete
- high blood pressure
- trouble urinating due to enlarged prostate gland
- cough accompanied by excessive phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product

- **do not use more than directed (see overdose warning)**
- **avoid alcoholic drinks**

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms get worse or last more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause serious health problems including liver damange. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- take only as directed- see Overdose warning
- use only dosing cup provided
- keep dosing cup provided with the product
- do not exceed 4 doses per 24 hours
- if taking Day Time during the day and Night Time at night, limit the total to 4 doses per 24 hours
- TBSP=tablespoon
- mL=milliliter

adults & children 12 years and over	30 mL (2 TBSP) every 4 hours
children 6 to under 12 years	15 mL (1 TBSP) every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

Other information

- each 15 mL contains: **sodium 5 mg**
- store between 58° - 86°F (15°-30°C).

Inactive ingredients: Citric acid, FD&C Yellow No 6, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or Comments? 1-888-446-4753

Monday - Friday from 8 AM - 5 PM Eastern Standard Time

XL-3^{*}

Day Time

Non-Drowsy Cold/Flu relief

COLD MEDICINE

- Acetaminophen - Pain Reliever/Fever Reducer
- Dextromethorphan HBr - Cough Suppressant
- Phenylephrine HCl - Nasal Decongestant

Fast relief of:

- NASAL CONGESTION
- FEVER
- ACHES & PAINS
- SORE THROAT
- COUGH

Orange Flavor

12 FL OZ (354 mL)

XL-3^{*}

Day Time

Non-Drowsy Cold/Flu relief

COLD MEDICINE

- Acetaminophen - Pain Reliever/Fever Reducer
- Dextromethorphan HBr - Cough Suppressant
- Phenylephrine HCl - Nasal Decongestant

Fast relief of:

- NASAL CONGESTION
- FEVER
- ACHES & PAINS
- SORE THROAT
- COUGH

Orange Flavor

12 FL OZ (354 mL)

XL-3^{*} Day Time

COLD MEDICINE

DO NOT USE IF PRINTED SHRINK BAND IS MISSING OR BROKEN
Failure to follow these warnings could result in serious consequences

Drug Facts	
Active ingredients	Purpose
<i>(in each 15 mL, 1 tablespoon)</i>	
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough Suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses: temporarily relieves common cold and flu symptoms:
• nasal congestion • cough due to minor throat and bronchial irritation • sore throat • headache • minor aches and pains • fever

→ DRUG FACTS CONTINUED ON BACK →

6 45981 00028 3

Distributed by:
Midway Importing, Inc.
1807 Brittonmore Road,
Houston, TX 77043 1-800-531-3873
Made in USA

Rev.1 LR-059 REV 03

XL-3^{*} Day Time

COLD MEDICINE

Drug Facts (continued)

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if • adult takes more than 4,000 mg of acetaminophen in 24 hours • child takes more than 5 doses in 24 hours • taken with other drugs containing acetaminophen • adult has 3 or more alcoholic drinks every day while using this product.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • for more than 10 days for pain unless directed by a doctor • for more than 3 days for fever unless directed by a doctor • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • thyroid disease • diabetes • high blood pressure • trouble urinating due to enlarged prostate gland • cough accompanied by excessive phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, or emphysema.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Drug Facts (continued)

When using this product • do not use more than directed (see overdose warning) • avoid alcoholic drinks

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • symptoms get worse or last more than 5 days (children) or 7 days (adults) • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause serious health problems including liver damage. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions • take only as directed - see Overdose warning • use only dosing cup provided • keep dosing cup provided with the product • do not exceed 4 doses per 24 hours • if taking Day Time during the day and Night Time at night, limit total to 4 doses per 24 hours • TBSP=tablespoon • mL=milliliter

adults & children 12 years and over	30 mL (2 TBSP) every 4 hours
children 6 to under 12 years	15 mL (1 TBSP) every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

Other information • each 15 mL contains: sodium 5 mg • store between 58° - 86°F (15° - 30°C).

Inactive ingredients: Citric acid, FD&C Yellow No.6, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum.

Questions or Comments? 1-888-446-4753
Monday - Friday from 8 AM - 5 PM Eastern Standard Time.

XL-3[®] Day Time COLD MEDICINE

DO NOT USE IF PRINTED SHRINK BAND IS MISSING OR BROKEN
Failure to follow these warnings could result in serious consequences

Drug Facts

Active ingredients (in each 15 mL, 1 tablespoon) **Purpose**

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough Suppressant
Phenylephrine HCl 5 mg Nasal decongestant

Uses temporarily relieves common cold and flu symptoms:
• nasal congestion • cough due to minor throat and bronchial irritation
• sore throat • headache • minor aches and pains • fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if • adult takes more than 4,000 mg of acetaminophen in

DRUG FACTS CONTINUED ON BACK

XL-3[®] Day Time

Non-Drowsy Cold/Flu relief

COLD MEDICINE

- Acetaminophen - Pain Reliever/Fever Reducer
- Dextromethorphan HBr - Cough Suppressant
- Phenylephrine HCl - Nasal Decongestant

Fast relief of:

- NASAL CONGESTION
- FEVER
- ACHES & PAINS
- SORE THROAT
- COUGH

6 FL OZ (177 mL)



XL-3[®] Day Time COLD MEDICINE



LR-117

Rev.1

Distributed by:
Midway Importing, Inc.
1807 Brittonmore Road,
Houston, TX
Made in USA

*Trademark

Peel back

XL-3[®] Day Time COLD MEDICINE

Drug Facts (continued)

24 hours • child takes more than 5 doses in 24 hours • taken with other drugs containing acetaminophen • adult has 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued)

Do not use • with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • for more than 10 days for pain unless directed by a doctor • for more than 3 days for fever unless directed by a doctor • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • thyroid disease • diabetes • high blood pressure • trouble urinating due to enlarged prostate gland • cough accompanied by excessive phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, or emphysema.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product • do not use more than directed (see overdose warning) • avoid alcoholic drinks

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • symptoms get worse or last more than 5 days (children) or 7 days (adults) • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause serious health problems including liver damage.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions • take only as directed - see Overdose warning • use only dosing cup provided • keep dosing cup provided with the product • do not exceed 4 doses per 24 hours • if taking Day Time during the day and Night Time at night, limit total to 4 doses per 24 hours • TBSP=tablespoon • mL=milliliter

adults & children 12 years and over 30 mL (2 TBSP) every 4 hours

children 6 to under 12 years 15 mL (1 TBSP) every 4 hours

children 4 to under 6 years ask a doctor

children under 4 years do not use

Other information • each 15 mL contains: sodium 5 mg
• store between 58°- 86°F (15°- 30°C).

Inactive ingredients: Citric acid, FD&C Yellow No.6, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum.

Questions or Comments? 1-888-446-4753

Monday - Friday from 8 AM - 5 PM Eastern Standard Time.

XL-3 DAY TIME

acetaminophen, dextromethorphan hbr, phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84379-441
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL

PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg in 15 mL
---	---------------	------------------

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID (UNII: 2968PHW8QP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MENTHOL (UNII: L7T10EIP3A)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84379-441-06	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2024	
2	NDC:84379-441-12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/01/2024	

Labeler - Rnv LLC (118917568)

Establishment

Name	Address	ID/FEI	Business Operations
Rnv LLC		118917568	manufacture(84379-441)

Revised: 1/2026

Rnv LLC