

NICOTINE POLACRILEX- nicotine polacrilex gum, chewing
Lucy Goods, Inc.

Nicotine Polacrilex Gum Mint 2 mg and 4 mg - Walgreens

Drug Facts

Active ingredient (in each chewing piece)

Nicotine polacrilex 2 mg (nicotine)

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions - 2 mg

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider

Other information - 2 mg

- **each piece contains:** calcium 115 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients - 2 mg

acacia, acesulfame potassium, carnauba wax, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide, xylitol.

Questions or comments?

Call

Principal display panel - 2 mg

*Compare to the active ingredient in Nicorette® Gum

Nicotine Gum

Nicotine polacrilex gum USP, 2 mg (nicotine)

Stop Smoking Aid

SUGAR-FREE

COATED MINT

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE **MORE THAN 30 MINUTES** AFTER WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Gum USP, 4 mg

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

- **not for sale to those under 18 years of age**
- **proof of age required**
- **not for sale in vending machines or from any source where proof of age cannot be verified**

*This product is not manufactured or distributed by Haleon group of companies, distributor of Nicorette® Gum.

Marketed BY Lucy Goods Inc.

2510 E Sunset Road

Suite 5 #A656

Las Vegas, NV 89120

Package label

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.



To remove the gum,
tear off single unit.



Peel off backing
starting at corner
with loose edge.



Push gum through
foil.

LUCY™

*Compare to the active ingredient in Nicorette® Gum

Nicotine Gum

Nicotine polacrilex gum USP, 2 mg (nicotine)
Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE
MORE THAN 30 MINUTES AFTER WAKING UP

If you smoke your first cigarette **WITHIN 30 MINUTES**
of waking up, use Nicotine Polacrilex Gum USP, 4 mg

2
mg

Sugar-Free

100 Pieces, 2 mg EACH

COATED
MINT



Actual Size

LUCY™

Nicotine Gum

Nicotine polacrilex gum USP, 2 mg (nicotine)
Stop Smoking Aid



8 500696 25576 3
Marketed By: Lucy Goods Inc.
2510 E Sunset Road
Suite 5 #A666
Las Vegas, NV 89120
The product is not manufactured or distributed by a member group of companies.
distributor of Nicorette® Gum.
flavors, gum base, polyvinylpyrrolidone, sodium benzoate, sodium carbonate, talc, titanium dioxide, xylitol

Directions
If you are under 18 years of age, ask a doctor before use.
Before using this product, read the enclosed User's Guide for complete directions and other important information.
If you are using the gum on your quit day:
If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum.
If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum.
Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. When used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.
If you have symptoms of an allergic reaction (such as difficulty breathing or rash) or you have experienced or experience such as nausea, vomiting, dizziness, drowsiness, and rapid heartbeat.
If you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, or rapid heartbeat or problems occur.
If mouth, throat, or jaw problems occur.

100 Pieces, 2 mg EACH

2
mg

COATED
MINT



Drug Facts Active ingredient (in each chewing piece) Nicotine polacrilex... 2 mg (nicotine) Purpose Stop smoking aid	Warnings If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risk to your child from this medicine are not fully known. Ask a doctor before use if you have: ■ a sudden-onset chest pain ■ heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. ■ high blood pressure not controlled with medication. Nicotine can increase blood pressure. ■ stomach ulcer or diabetes ■ history of seizures Ask a doctor or pharmacist before use if you are: ■ taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted. ■ taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.
Use ● Nicotine withdrawal symptoms, including nicotine craving, associated with quitting smoking	Other information ■ each piece contains calcium 115 mg, sodium 6 mg ■ store at 20°-25°C (68°-77°F) ■ protect from light ■ protect from light Inactive ingredients acacia, acacia thorne preservative, carnauba wax.

not for sale to those under 18 years of age
proof of age required
not for sale by vending machines or from any source where proof of age cannot be verified

REVIEWER PROVIDED FOR YOUR PROTECTION. DO NOT TAKE IF

LUCY Nicotine Gum Coated Mint

NICOTINE POLACRILEX nicotine polacrilex gum, chewing			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72224-927
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)		NICOTINE	2 mg
Inactive Ingredients			
Ingredient Name			Strength
ACACIA (UNII: 5C5403N26O)			
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)			
MAGNESIUM OXIDE (UNII: 3A3U0G171G)			
SODIUM BICARBONATE (UNII: 8MDF5V39QO)			
SODIUM CARBONATE (UNII: 45P3261C7T)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
XYLITOL (UNII: VCQ006KQ1E)			
Product Characteristics			
Color	white	Score	no score
Shape	SQUARE	Size	14mm
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72224-927-10	100 in 1 CARTON	05/30/2025	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078699	04/30/2025	

Labeler - Lucy Goods, Inc. (080875415)

Revised: 4/2025

Lucy Goods, Inc.