

**BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE- liquid  
bandage liquid  
Chain Drug Marketing Association**

-----  
**Quality Choice Liquid Bandage**

***Active ingredients***

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

***Purpose***

First aid antiseptic

Topical analgesic

***Uses***

- first aid to prevent bacteria contamination in minor cuts and scrapes

***Warnings***

**For external use only.**

**Flammable:**

do not use while smoking or near heat or flame.

**Do not use**

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor
- on infected areas or wounds that are draining
- with other first aid products such as lotions and creams
- over sutures
- on mucous membranes

**Ask a doctor before use if you have**

- deep puncture wounds
- deep cuts
- animal bites
- serious bleeding
- diabetes
- poor circulation
- serious burns

## **When using this product**

- do not store at temperature above 120 °F

Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

## **Stop use and ask a doctor if**

- conditions persists or gets worse
- infection occurs

## **Keep out of reach of children.**

If swallowed, get medical help or call a Poison Control Center right away.

## ***Directions***

- Clean affected area
- apply a small amount on the area 1-3 times daily
- let dry
- a second coating may be applied for extra protection
- to remove, apply more Liquid Bandage and quickly wipe off
- finger nail polish remover may dissolve Liquid Bandage

## ***Other information***

- ***cap bottle tightly***
- store at room temperature away from heat
- may temporarily sting upon application
- do not allow to come in contact with floors, countertops, or other finished surfaces - will stain

## ***Inactive ingredients***

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

## ***Questions***

Call 1-866-964-0939

## **Principal Display Panel**

### **Quality Choice**

### **For Small Cuts & Wounds**

### **Liquid Bandage**

Antiseptic to Help Prevent Infection

Invisible | Flexible | Waterproof | Easy to use

**Use Liquid Bandage For:**

Blisters, chapped & cracked finger tips, hangnails, paper cuts, shaving nicks, & help in preventing the formulation of calluses.

**Also Used By:**

Bowlers, dancers, fishermen, golfers, hikers, musicians, runners, and tennis players

**0.3 FL OZ. (9 mL)**



\*Compare to SKIN SHIELD®

For Small Cuts & Wounds

# Liquid Bandage

Antiseptic To Help Prevent Infection

Invisible | Flexible | Waterproof  
Easy to use

**Use Liquid Bandage For:**

Blisters, chapped & cracked finger tips, hangnails, paper cuts, shaving nicks & help in preventing the formation of calluses.

**Also Used By:**

Bowlers, dancers, fishermen, golfers, hikers, musicians, Runners, and tennis players.

LB00BT



For Small Cuts & Wounds

# Liquid Bandage

Antiseptic To Help Prevent Infection

0.3 FL. OZ. (9mL)

NO VARNISH AREA



©Distributed by C.D.M.A., Inc.  
43157 W. Nine Mile  
Novi, MI 48376-0995  
www.qualitychoice.com  
Questions: 248-449-9300

\* This product is not manufactured or distributed by Insight Pharmaceuticals, LLC, owner of the registered trademark Skin Shield®.

MADE IN USA



**DO NOT USE IF NECKBAND IS BROKEN**

**Drug Facts**

**Purpose**  
First aid antiseptic  
Topical antiseptic

**Active Ingredients**  
Benzethonium chloride 0.75%  
Dyclonine hydrochloride 0.75%

**Uses**  
First aid to prevent bacterial contamination in minor cuts, scrapes and burns

**Warnings**  
For external use only.

**Flammable:** do not use while smoking or near heat or flame.

**Do not use**  
■ in the eyes ■ over large areas of the body ■ longer than 1 week unless directed by a doctor  
■ on infected areas or wounds that are draining ■ with other first aid products such as balm and cream ■ over sutures ■ on mucous membranes

**Ask a doctor before use if you have**  
■ deep puncture wounds ■ deep cuts ■ arterial lacerations ■ serious bleeding ■ diabetes ■ poor circulation ■ serious burns

**When using this product**  
■ do not store at temperature above 120°F ■ avoid use by deliberately concentrating and shaking the contents can be harmful or fatal

**Stop use and ask a doctor if**  
■ condition persists or gets worse ■ reaction occurs

**Keep out of reach of children.** If swallowed, get medical help or call a Poison Control Center right away.

**Directions**  
■ clean affected area ■ apply a small amount on the area 1-3 times daily ■ let dry ■ a second coating may be applied for extra protection ■ to remove, apply more liquid bandage and quickly wipe off ■ fingernail polish remover may dissolve Liquid Bandage

**Other information**  
■ cap bottle tightly ■ store at room temperature away from heat ■ may temporarily sting upon application ■ do not allow to come in contact with hair, countertops, or other finished surfaces — will stain

**Inactive ingredients**  
alcohol, benzalkonium chloride, butyl acetate, castor oil, ethyl acetate, hydroxyacetone, SD alcohol 40

**Questions?** Call 1-248-449-9300

PRINT SIDE

**BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE**  
liquid bandage liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-045
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BENZETHONIUM CHLORIDE</b> (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 9 mL
<b>DYCLONINE HYDROCHLORIDE</b> (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q30O)	DYCLONINE HYDROCHLORIDE	0.75 mg in 9 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>ACETONE</b> (UNII: 1364PS73AF)	
<b>AMYL ACETATE</b> (UNII: 92Q24NH7AS)	
<b>CASTOR OIL</b> (UNII: D5340Y2I9G)	
<b>ETHYL ACETATE</b> (UNII: 76845O8NMZ)	
<b>PYROXYLIN</b> (UNII: KYR8BR2X6O)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:63868-045-03	9 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product	12/19/2017	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M003	12/19/2017	

**Labeler** - Chain Drug Marketing Association (011920774)