

**CAREALL IBUPROFEN- ibuprofen tablet**  
**New World Imports**

-----  
**Drug Facts**

Ibuprofen 200mg (NSAID)\*\*

\*\*nonsteroidal anti-inflammatory drug

Pain reliever / fever reducer

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Temporarily relieves minor aches and pains due to:

Headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis.

Temporarily reduces fever

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you are 60 years or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing a NSAID [aspirin, ibuprofen, naproxen, or others], have 3 or more alcoholic drinks every day while using this product, or take more / for a longer time than directed.

**Heart attack and stroke warning:** NSAID's, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** if you have ever had an allergic reaction to any other pain reliever / fever reducer or right before or after heart surgery.

**Ask a doctor before use if:** stomach bleeding warning applies to you. you have problems or serious side effects from taking pain relievers or fever reducers. You have a history of stomach problems, such as heartburn. You have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke. You are taking a diuretic.

**Ask a doctor or pharmacist before use if you are:** taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin. Under a doctor's care for any serious condition. Taking any other drug.

**When using this product:** take with food or milk if stomach upset occurs.

**Stop use and ask a doctor if:** You experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better You have symptoms of heart problems or stroke: chest pain,

trouble breathing, weakness in one part or side of body, slurred speech, or leg swelling. Pain gets worse or lasts more than 10 days. Fever gets worse or lasts more than 3 days. Redness or swelling is present in the painful area. Any new symptoms appear.

**Pregnant or breast-feeding:** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Do not take more than directed. The smallest effective dose should be used.**

**Adult and children 12 years and older:** Take 1 tablets every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablets, 2 tablets may be used. Do not exceed 6 tablets in 24 hours unless directed by a doctor.

**Children under 12 years:** ask a doctor

colloidal silicon dioxide, corn starch, FD and C yellow 6, hypromellose, polydextrose, polyethylene glycol, povidone, pregelatinized starch, purified water, sodium starch glycolate, stearic acid, titanium dioxide

Store between 20-25 degrees celsius (68-77 degrees farhenheit).

Read all warnings and directions before use

Do not use if seal under bottle cap is broken or missing

*Questions or Comments:* Call 1-888-952-0050 Monday through Friday 9 AM - 5 PM EST

NDC 51824-064-01



# Ibuprofen (NSAID)

Tablets, 200mg

- ▶ Pain Reliever
- ▶ Fever Reducer

Compare to the active ingredient in Advil®\*\*



**100 200mg Tablets**

**DO NOT USE IF SEAL UNDER BOTTLE CAP IS BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)** Purpose  
Ibuprofen USP, 200mg (NSAID)\* .....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
● headache ● toothache ● backache ● menstrual cramps  
● the common cold ● muscular aches ● minor pain of arthritis  
■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older  
■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any

**Drug Facts (continued on inside)**

\*\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings, owner of the registered trademark Advil®. Distributed By: CareAll Products, 160 Athens Way, Nashville, TN 37228  
www.careallproducts.com



0 85317 00502 3

Lot No. Exp. Date:  
**PREP HERE**

**Drug Facts (continued)**  
other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product** take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ● you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ do not take more than directed  
■ the smallest effective dose should be used  
■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor  
■ children under 12 years: ask a doctor

**Other information** ■ store between 20-25 °C (68-77° F) ■ read all warnings and directions before use

**Inactive ingredients** colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, polyketose, polyethylene glycol, povidone, pregelatinized starch, purified water, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or Comments?** Call 1-888-962-0050 Monday through Friday 9AM – 5PM EST

## CAREALL IBUPROFEN

ibuprofen tablet

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:51824-064 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength |
|--|-------------------|----------|
| IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM) | IBUPROFEN         | 200 mg   |

### Inactive Ingredients

| Ingredient Name                    | Strength |
|------------------------------------|----------|
| Starch, Corn (UNII: O8232NY3SJ)    |          |
| HYPROMELLOSES (UNII: 3NXW29V3WO)   |          |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) |          |

|   |  |
|---|--|
| <b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)                 |  |
| <b>POVIDONE</b> (UNII: FZ989GH94E)                              |  |
| <b>WATER</b> (UNII: 059QF0KO0R)                                 |  |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)                          |  |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)                      |  |
| <b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)                          |  |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)      |  |
| <b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2) |  |

### Product Characteristics

|                 |                           |                     |          |
|-----------------|---------------------------|---------------------|----------|
| <b>Color</b>    | brown (Rusty brown color) | <b>Score</b>        | no score |
| <b>Shape</b>    | ROUND (Round)             | <b>Size</b>         | 10mm     |
| <b>Flavor</b>   |                           | <b>Imprint Code</b> | G2       |
| <b>Contains</b> |                           |                     |          |

### Packaging

| # | Item Code        | Package Description                                | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:51824-064-01 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 08/11/2017           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA079174                               | 08/11/2017           |                    |

**Labeler** - New World Imports (075372276)

**Registrant** - New World Imports (075372276)