

LIDOCAINE PAIN RELIEF GEL-PATCH- lidocaine patch
Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.

Liuquanmiao, LIDOCAINE PAIN RELIEF Gel-Patch, Lidocaine 4%, 18 Patches

Active ingredients

Lidocaine 4% Purpose: Topical Analgesic

Purpose

Topical Analgesic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly

Stop use ... if

- skin reactions such as redness, swelling, blistering or other discomfort occur
- symptoms persist for more than 7 days

... ask a doctor if

- skin reactions such as redness, swelling, blistering or other discomfort occur
- symptoms persist for more than 7 days

If pregnant or breast-feeding

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Remove patch from the skin after at most 8-hour application.

Children under age of 12: Consult a doctor

Inactive Ingredients

Dihydroxyaluminum Aminoacetate, Disodium Ethylenediaminetetraacetic Acid, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Water

Usage


Usage



Remove patch from
outer package



Partially peel back
protective film



Place on
affected area



Remove remaining film
while pressing patch to skin

Questions or Comments

sumifunofficial@gmail.com

158-16

www.sumifunofficial.com
 INST: @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunOfficial
 TikTok: @sumifunofficial
 Twitter: @sumifunOfficial

Sumifun®


LIDOCAINE PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- Odor-Free
- Maximum Strength
- Easy to apply and remove
- Single-Use Application

• Knee	• Back	• Neck
• Elbow	• Shoulders	



16 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)

Guangzhou Lidoanmed Import & Export Trade Co., Ltd.
 #12, Building 8, No. 81, Dongfeng Road, Baiyuan,
 Zhuang, Tianhe District, Guangzhou/Guangdong, China

Made in PRC



unjunns

Drug Facts

Active ingredients Purpose
 Lidocaine, etc. Topical Anesthetic

Uses
 Temporarily relieves muscle soreness and minor joint pains in the wrist, knee, back, neck, hip, shoulders, elbows.

Warnings
 For external use only

Do not use
 on wounds or damaged skin
 with a heating pad or device
 with other creams, ointments, sprays, or liniments
 if you are allergic to any ingredients of this product

When using this product
 use only as directed
 avoid contact with the eyes, mucous membranes, or mucus
 do not bandage tightly

Stop use and ask a doctor if
 skin reactions such as redness, swelling, blistering or
 other discomfort occur
 symptoms persist for more than 7 days

If pregnant or breast-feeding,
 ask a health professional before use
KEEP OUT OF REACH OF CHILDREN.
 If swallowed accidentally, get medical help or contact
 with Poison Control Center immediately.



Directions

Adults and Children above 12 years old:
 Clean and dry the affected area.
 Peel off the protective film and apply the exposed part of
 the patch to the affected area.
 Carefully remove remaining film while pressing the patch
 firmly on the skin.
 Remove patch from the skin after at most 8-hour
 application.
 Children under age of 12: Consult a doctor

Inactive ingredients

Dihydroxyaluminum Aminoacetate,
 Disodium Ethylenediaminetetraacetic Acid, Glycolin,
 Kapton, Methylpropane Polycarbonate (3), Polyurethane Glycol,
 Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben,
 Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water

Usage



Questions or Comments: support@sumifunofficial.com
 Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging





www.sumifunofficial.com
 Instagram: @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunofficial
 TikTok: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- Odor-Free
- Maximum Strength
- Easy to apply and remove
- Single-Use Application

• Knee	• Back	• Neck
• Elbow	• Shoulders	

20 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)

Guangzhou Liguangshan Import & Export Trade Co., Ltd.
 17, Zhilong 3, No. 8, Changyang San Road,
 Zhushu, Tianhe District, Guangzhou, Guangdong, China

Made in PRC

Sumifun®

Drug Facts

Active Ingredients: Lidocaine 4% Purpose: Topical Anesthetic

Uses: Temporarily relieves muscle soreness and minor joint pain in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings: For external use only

Do not use on wounds or damaged skin with a healing pad or device with other ointments, creams, sprays, or lotions. If you are allergic to any ingredients of this product.

When using this product: Use only as directed and avoid contact with the eyes, mucous membranes, or mouth. Do not bandage tightly.

Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur; symptoms persist for more than 7 days.

If pregnant or breast-feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.

Inactive Ingredients

Dihydroxyaluminum Ammonioacetate, Stearic Acid, Ethylenediaminetetraacetic Acid, Glycerin, Kaolin, Methylparaben, Polyethylene Glycol, Polyethylene Glycol, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water

Usage



Questions or Comments: support@sumifunofficial.com Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging





www.sumifunofficial.com
 Instagram: @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunofficial
 TikTok: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- Odor-Free
- Maximum Strength
- Easy to apply and remove
- Single-Use Application

• Knee	• Back	• Neck
• Elbow	• Shoulders	

21 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)

Guangzhou Lijianmei Import & Export Trade Co., Ltd.
 17, Zhilong 3, No. 8, Changyang San Road,
 Zhuzhou, Tianhu District, Guangzhou, Guangdong, China

Made in PRC

Sumifun®

Drug Facts

Active Ingredients: Lidocaine 4% Purpose: Topical Anesthetic

Uses: Temporarily relieves muscle soreness and minor joint pain in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only.

Do not use

on wounds or damaged skin with a heating pad or device with other ointments, creams, sprays, or lotions. If you are allergic to any ingredients of this product.

When using this product

use only as directed and avoid contact with the eyes, mucous membranes, or mouth. Do not bandage tightly.

Stop use and ask a doctor if

skin reactions such as redness, swelling, blistering or other discomfort occur; symptoms persist for more than 7 days.

If pregnant or breast-feeding,

ask a health professional before use. **KEEP OUT OF REACH OF CHILDREN.** If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.

Inactive Ingredients

Dihydroxyaluminum Ammonioacetate, Disodium Ethylenediaminetetraacetic Acid, Glycerin, Kaolin, Methylparaben, Polyethylene Glycol, Polyethylene Glycol, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water.

Usage



Questions or Comments: support@sumifunofficial.com Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging



Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch

4%

LIDOCAINE



Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

• Knee	• Back	• Neck
• Elbow	• Shoulders	

28

PATCHES
 3.5" x 3.94"
 (9cm x 10cm)

Guangdong Cooperation Support & Export Trade Co., Ltd.
 617 Building A, No. 8, Song Hong Sun Road,
 Zhuzhen, Nanhai District, Guangzhou, Guangdong, China.

Made in PRC

 Knee
 Back
 Neck
 Elbow
 Shoulder

www.sumifunofficial.com
 Instagram: @sumifunofficial
 Facebook: @sumifunofficial
 YouTube: @sumifun
 Twitter: @sumifunofficial

Sumifun



<p>Drug Facts</p> <p>Active ingredients _____ Purpose Lidocaine 4% _____ Topical Anesthetic</p> <p>Uses Temporarily relieves muscle spasms and minor joint pain in the neck, lower back, neck, hip, shoulders, elbow.</p> <p>Warnings For external use only</p> <p>Do not use on wounds or damaged skin with a heating pad or device with other creams, ointments, sprays, or liniments if you are allergic to any ingredients of this product</p> <p>When using this product use only as directed avoid contact with the eyes, mucous membranes, or rubber do not bandage tightly</p> <p>Stop use and ask a doctor if skin reactions such as redness, swelling, itching or other discomfort occur symptoms persist for more than 7 days</p> <p>If pregnant or breast-feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.</p>	<p>Directions Adults and Children above 12 years old: Clean and dry the affected area. Peel off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor</p> <p>Inactive ingredients Diglycylsuccinate, Antiseptics, Oxidium Ethylenediaminetetraacetic Acid, Glycerin, Kaolin, Methylcellulose, Polyethylene Glycol, Propylene Glycol, Polyethylene Glycol, Polyethylene Glycol, Propylene Glycol, Sodium Polymethacrylate, Titanium Dioxide, Terephthalic Acid, Water</p> <p>Usage</p>    
<p>Child Resistant Packaging</p> 	

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch

Numbing muscle and relieve pain:

4%
LIDOCAINE

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

• Knee	• Back	• Neck
• Elbow	• Shoulders	

30 PATCHES
3.5" x 3.94"
(9cm x 10cm)

Guangdong Cooperation Support & Export Trade Co., Ltd.
617 Building A, No. 8, Dong Hong Sun Road,
Duzhou, Foshan City, Guangdong, China

Made in PRC

www.sumifunofficial.com
INS: @sumifunofficial
YouTube: @sumifun
Facebook: @sumifunofficial
TikTok: @sumifunofficial
Twitter: @sumifunofficial

Sumifun

Drug Facts	Directions Adults and Children above 12 years old: Clean and dry the affected area. Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while peeling the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.
Active Ingredients Lidocaine 4% <small>Typical Appearance:</small>	Warnings For external use only.
Uses Temporarily relieves muscle spasms and minor joint pains in the neck, lower back, neck, hip, shoulders, elbow.	Do not use on wounds or damaged skin with a heating pad or device with other ointments, creams, ointments, or liniments if you are allergic to any ingredients of this product.
When using this product use only as directed avoid contact with the eyes, mucous membranes, or rubber do not bandage tightly.	Inactive Ingredients Diphenylmethane Antimonate, Dioxolane Ethylenechloroacetate, Acid, Glycerin, Kathon, Methacrylates, Polyethylene Glycol, Propylene Glycol, Polyethylene Glycol, Polyethylene Glycol, Propylene Glycol, Sodium Polymethacrylate, Titanium Dioxide, Terephthalic Acid, Water
Stop use and ask a doctor if: skin reactions such as redness, swelling, itching or other discomfort occur symptoms persist for more than 7 days.	Usage
If pregnant or breast-feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.	
	Child Resistant Packaging

Questions or Comments: support@sumifunofficial.com
Business Cooperation: business@sumifunofficial.com

CUT OPEN HERE

Sumifun®

LIDOCAINE


PAIN RELIEF Gel-Patch

Numbing muscle and relieve pain:



- **Odor-Free**
- **Maximum Strength**
- *Easy to apply and remove*
- *Single-Use Application*

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



1 PATCH
5.5" x 3.94"
(14cm x 10cm)

Drug Facts

Active ingredients **Purpose**
Lidocaine 4% Topical Anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly

Stop use and ask a doctor if

- skin reactions such as redness, swelling, blistering or other discomfort occur
- symptoms persist for more than 7 days

If pregnant or breast-feeding,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Remove patch from the skin after at most 8-hour application.

Children under age of 12: Consult a doctor

Inactive Ingredients

Dihydroxyaluminum Aminoacetate, Disodium Ethylenediaminetetraacetic Acid, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Usage



Remove patch from outer package



Partially peel back protective film



Place on affected area



Remove remaining film while pressing patch to skin

Questions or Comments: supports@sumifunofficial.com
Business Cooperation: business@sumifunofficial.com

Guangzhou Liuguang Import & Export Trade Co., Ltd.
417, Building A, No. 8, Dong Heng San Road,
Zhucun, Tianhe District, Guangzhou, Guangdong, China.

Child Resistant Packaging



www.sumifunofficial.com
 INSTA: @sumifunofficial
 YOUTUBE: @sumifun
 FACEBOOK: @sumifunofficial
 TIKTOK: @sumifunofficial
 TWITTER: @sumifunofficial

Sumifun®


LIDOCAINE PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- Odor-Free
- Maximum Strength
- Easy to apply and remove
- Single-Use Application

• Knee	• Back	• Neck
• Elbow	• Shoulders	



15 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)

Guangzhou Lidoanmed Import & Export Trade Co., Ltd.
 #12, Building 8, No. 81, Dongfeng Road,
 Zhuzhou, Tianhe District, Guangzhou/Guangdong, China

Made in PRC



sumifun

Drug Facts

Active ingredients: Lidocaine, etc. Purpose: Topical Anesthetic.

Uses: Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only.

Do not use on wounds or damaged skin with a healing pad or device with other creams, ointments, sprays, or liniments if you are allergic to any ingredients of this product.

When using this product

use only as directed. avoid contact with the eyes, mucous membranes, or mucus. do not bandage tightly.

Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur. symptoms persist for more than 7 days.

If pregnant or breast-feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.


Directions

Adults and Children above 12 years old: Clean and dry the affected area. Peel off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.

Inactive Ingredients

Dihydroxyaluminum Aminoacetate, Disodium Ethylenediaminetetraacetic Acid, Glycolin, Kapton, Methylpropane Polycarbonate (M), Polyurethane Glycol, Polyacrylic Acid, Polyvinylpyrrolidone (PVP), Propylparaben, Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water.

Usage



Questions or Comments: support@sumifunofficial.com
 Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging



Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch




Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	

31 PATCHES
5.5" x 3.94"
(14cm x 10cm)

Sumifun (Lidocaine) Patch & Gel-Patch, Inc. 2020
 217, Building A, No. 4, Changyang San Road,
 Zhuzhou, Hunan District, Guangzhou, Guangdong, China

Made in PRC

Sumifun

<p>Drug Facts</p> <p>Active Ingredients..... Patchlet Lidocaine, 4%..... Topical Anesthetic</p> <p>Uses Temporarily relieves muscle cramps and minor pain points in the wrist, knee, back, neck, hip, shoulder, elbow.</p> <p>Warnings For external use only.</p> <p>Do not use on wounds or damaged skin with a heating pad or device with other creams, sprays, or lotions, if you are allergic to any ingredients of this product.</p> <p>When using this product use only as directed avoid contact with the eyes, mucous membranes, or tubes do not bandage tightly.</p> <p>Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days.</p> <p>If pregnant or breast feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If inhaled accidentally, get medical help or contact with Poison Control Center immediately.</p>	<p>Directions Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.</p> <p>Inactive Ingredients Diphenylmethane Antioxidant, Dioctyl Ethylhexedimethylsilyoxane, AcM Glycerin, Acetic Acid, Polypropylene, Polyethylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Polyisobutylene, Sodium Polyacrylate, Titanium Dioxide, Terephthalic Acid, Water</p> <p>Usage</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  <p>Remove patch from outer package</p> </div> <div style="text-align: center;">  <p>Patch on neck</p> </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 10px;"> <div style="text-align: center;">  <p>Patch on elbow</p> </div> <div style="text-align: center;">  <p>Remove remaining film while pressing patch to skin</p> </div> </div> <p>Questions or Comments: support@sumifun.com Business Cooperation: business@sumifun.com</p> <p style="text-align: center;">Child Resistant Packaging</p> <div style="text-align: right;">  <p>6 18 773 14 249 07 8</p> </div>
--	---

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch




Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

• Knee	• Back	• Neck
• Elbow	• Shoulders	

32 PATCHES
5.5" x 3.94"
(14cm x 10cm)

Guangzhou Sumifun Pharmaceutical Co., Ltd.
 211, Building A, No. 4, Changhong San Road,
 Zhushan, Lantau District, Guangzhou, Guangdong, China.
 Made in PRC

Sumifun

<p>Drug Facts</p> <p>Active Ingredients..... Patched LidoCAINE 4%..... Proprietary Topical Anesthetic</p> <p>Uses Temporarily relieves muscle cramps and minor pain points in the wrist, knee, back, neck, hip, shoulder, elbow.</p> <p>Warnings For external use only.</p> <p>Do not use on wounds or damaged skin with a healing pad or device with other creams, sprays, or lotions, if you are allergic to any ingredients of this product.</p> <p>When using this product use only as directed avoid contact with the eyes, mucous membranes, or teeth do not bandage tightly.</p> <p>Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days.</p> <p>If pregnant or breast feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If inhaled accidentally, get medical help or contact with Poison Control Center immediately.</p>	<p>Directions Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.</p> <p>Inactive Ingredients Diphenylmethane Antioxidant, Dioxolane Ethylene Dimethyleneacrylate, ADM Glycerin, Acrylic Methacrylate, Polyethylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Polyisobutylene, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water</p> <p>Usage</p> <div style="display: flex; justify-content: space-around;"> <div>  <p>Remove patch from outer package</p> </div> <div>  <p>Patch on back neck protective film</p> </div> </div> <div style="display: flex; justify-content: space-around;"> <div>  <p>Tear off affected area</p> </div> <div>  <p>Remove remaining film while pressing patch to skin</p> </div> </div> <p>Questions or Comments: support@sumifun.com Business Cooperation: business@sumifun.com</p> <p>Child Resistant Packaging</p>  <p>6 18 773 14 24 908 5</p>
---	--

www.sumifunofficial.com
 @NSF@sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunofficial
 TikTok: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®


LIDOCAINE PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



Knee



Back



Neck



Elbow



Shoulder

39 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)

Guangzhou Lidaqianhua Import & Export Trade Co., Ltd.
 517 Building A, No. 4, South Ring East Road,
 Zhujiang, Tianhe District, Guangzhou, Guangdong, China.

Back © 2020

Sumifun

<p>Drug Facts</p> <p>Active ingredients Purpose Lidocaine HCl Topical Anesthetic</p> <p>Uses Temporarily relieves muscle soreness and minor joint pains in the wrist, lower back, neck, hips, shoulders, elbow.</p> <p>Warnings For external use only.</p> <p>Do not use - on wounds or damaged skin - with a heating pad or device - with other creams, ointments, sprays, or lotions - if you are allergic to any ingredients of this product</p> <p>When using this product - use only as directed - avoid contact with the eyes, mucous membranes, or neck - do not bandage tightly</p> <p>Stop use and ask a doctor if - skin reactions such as redness, swelling, blistering or other discomfort occur - symptoms persist for more than 7 days</p> <p>Keep out of reach of children. If swallowed, immediately get medical help or contact with Poison Control Center immediately.</p>	<p>Directions Adults and Children above 12 years old - Clean and dry the affected area. - Peel off the protection film and apply the exposed part of the patch to the affected area. - Gently remove remaining film while pressing the patch being on the skin. - Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor</p> <p>Inactive Ingredients Ethylhexylmethacrylate, Dimethylsiloxane, Hydroxyethyl Methacrylate, Glyceryl Acrylate, Methyl Methacrylate, Polyethylene Glycol, Polyethylene Glycol, Polyethylene Glycol, Polyethylene Glycol, Sodium Polycrylate, Titanium Dioxide, Sorbic Acid, Water</p> <p>Usage</p> <div style="display: flex; justify-content: space-around;"> <div> <p>Patch pack (39 patches)</p> </div> <div> <p>Remove remaining film while pressing patch to skin</p> </div> </div> <p>Questions or Comments: support@sumifunofficial.com Business Cooperation: business@sumifunofficial.com</p> <p>OTC Non-Drug Packaging</p>
--	---

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch




Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

• Knee	• Back	• Neck
• Elbow	• Shoulders	


38 PATCHES
5.9" x 3.94"
 (14.9cm x 10cm)

Sumifun (Lidocaine) Patch and Patch Pack, Inc.
 10000 S. Bascom Ave., Suite 100, San Jose, CA 95128
 ©2018 Sumifun (Lidocaine) Patch and Patch Pack, Inc.

Made in PRC

Sumifun

<p>Drug Facts</p> <p>Active ingredients: Lidocaine HCl</p> <p>Purpose: Topical Anesthetic</p> <p>Uses: Temporarily relieves muscle aches and minor joint pains in the wrist, knee, back, neck, hip, shoulder, elbow.</p> <p>Warnings: For external use only.</p> <p>Do not use: on wounds or damaged skin with a heating pad or device with other heating systems, wraps, or treatments if you are allergic to any ingredients of this product.</p> <p>When using this product: use only as directed; avoid contact with the eyes, mucous membranes, or mouth; do not bandage tightly.</p> <p>Stop use and ask a doctor if: skin reactions such as redness, swelling, blistering or other reactions occur; symptoms persist for more than 7 days.</p> <p>If pregnant or breast-feeding: ask a health care provider before use. KEEP OUT OF REACH OF CHILDREN. For more information, get medical help or contact your Poison Control Center immediately.</p>	<p>Directions: Adults and Children about 12 years old: Clean and dry the affected area. Tear off the adhesive front and apply the exposed part of the patch to the affected area. Carefully remove adhesive from back, pressing the patch firmly on the skin. Remove patch from the skin after or just before application. Children under age of 12: Consult a doctor.</p> <p>Inactive Ingredients: Chlorophyllin/Aluminum Antiseptics, Dimethyl Ethylhexanoate/Dimethyl Ethylhexanoate, Glycerin, Kaolin, Methylparaben, Polyethylene Glycol, Propylene Glycol, Polyethylene Glycol, Polyethylene Glycol, Propylene Glycol, Sodium Hydroxide, Talcum Powder, Talcum Powder, Talcum Powder.</p> <p>Usage:</p>  <p>Questions or Comments: support@sumifun.com Business Inquiries: business@sumifun.com</p> <p>Child-Resistant Packaging:</p> 
---	--



Drug Facts

Active Ingredients: LIDOCAINE HCl

Purpose: Temporary relief of minor aches and pains.

Typical Appearance: Clear, adhesive patch.

Use: Temporarily relieves muscle aches and minor joint pains in the arms, back, neck, hip, shoulders, elbow.

Directions: Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8 hour application. Children under age of 12: Consult a doctor.

Warnings: For external use only.

Do not use: if wounds or damaged skin with a heavy coat of ointment with other patches, creams, ointments, lotions, or liniments. If you are allergic to any ingredients of the product.

When using this product: avoid contact with the eyes, mucous membranes, or broken skin. Do not bandage tightly.

Stop use and ask a doctor if: skin reactions such as redness, swelling, blistering or other discomfort occur. Symptoms persist for more than 7 days.

If pregnant or breast-feeding: ask a health professional before use. **KEEP OUT OF REACH OF CHILDREN.** For medical information, get medical help or contact with Poison Control Center immediately.

Directions: Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8 hour application. Children under age of 12: Consult a doctor.

Warnings: For external use only.

Do not use: if wounds or damaged skin with a heavy coat of ointment with other patches, creams, ointments, lotions, or liniments. If you are allergic to any ingredients of the product.

When using this product: avoid contact with the eyes, mucous membranes, or broken skin. Do not bandage tightly.

Stop use and ask a doctor if: skin reactions such as redness, swelling, blistering or other discomfort occur. Symptoms persist for more than 7 days.

If pregnant or breast-feeding: ask a health professional before use. **KEEP OUT OF REACH OF CHILDREN.** For medical information, get medical help or contact with Poison Control Center immediately.

Questions or Comments: support@sumifunofficial.com

Child Resistant Packaging:



LIDOCAINE PAIN RELIEF GEL-PATCH

lidocaine patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83602-158
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.04 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
GLYCERIN (UNII: PDC6A3C0OX)	

POVIDONE K90 (UNII: RDH86HJV5Z)
METHYLPARABEN (UNII: A2I8C7HI9T)
PROPYLPARABEN (UNII: Z8IX2SC1OH)
KAOLIN (UNII: 24H4NWX5CO)
TARTARIC ACID (UNII: W4888I119H)
WATER (UNII: 059QF0KO0R)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JNI2J)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83602-158-15	15 in 1 BOX	10/14/2024	
1	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		
2	NDC:83602-158-16	16 in 1 BOX	10/14/2024	
2	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		
3	NDC:83602-158-20	20 in 1 BOX	10/14/2024	
3	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		
4	NDC:83602-158-21	21 in 1 BOX	10/14/2024	
4	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		
5	NDC:83602-158-28	28 in 1 BOX	10/14/2024	
5	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		
6	NDC:83602-158-30	30 in 1 BOX	10/14/2024	
6	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		
7	NDC:83602-158-31	31 in 1 BOX	10/14/2024	
7	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		
8	NDC:83602-158-32	32 in 1 BOX	10/14/2024	
8	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		

9	NDC:83602-158-38	38 in 1 BOX	10/14/2024	
9	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		
10	NDC:83602-158-39	39 in 1 BOX	10/14/2024	
10	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		
11	NDC:83602-158-40	40 in 1 BOX	10/14/2024	
11	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		
12	NDC:83602-158-41	41 in 1 BOX	10/14/2024	
12	NDC:83602-158-01	12 g in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	08/08/2023	

Labeler - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Registrant - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Revised: 10/2024

Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.