#### DAYTIME COLD AND FLU- acetaminophen dextromethorphan hbr phenylephrine hcl capsule, liquid filled QUALITY CHOICE (Chain Drug Marketing Association)

\_\_\_\_\_

#### **Drug Facts**

#### Active ingredients (in each softgel)

#### Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

#### Purpose

#### Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

#### Uses

- temporarily relieves common cold and flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - fever
  - cough due to minor throat and bronchial irritation

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**: Acetaminophen may cause severe skin reactions. symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## Ask a doctor before use if you have

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

## Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

## When using this product,

## do not exceed recommended dosage.

#### Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

## Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

## Inactive ingredients

FD&C red #40, FD&C yellow#6, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

## **Questions or comments?**

Call 1-800-935-2362 Monday-Friday 9AM-5PM EST

# Principal Display Panel

†Compare to the Active Ingredients in VICKS® DAYQUIL® COLD & FLU LIQUICAPS®

Non-Drowsy

## Daytime

# Multi-Symptom Relief For Cold/Flu

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

Pain Reliever | Fever Reducer

Cough Suppressant | Nasal Decongestant

Non-Drowsy | Alcohol Free | Antihistamine Free

Softgels

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

#### TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWING ANY SIGNS OF TAMPERING.

# KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by: CDMA, Inc. Novi MI 48375

Package Label



trassandque riguo. Dextromethorphan HBr 10 mg. 🔳 usaal congestion Tevel ain reliever/fever reducer gm 225 neriqonimeteoA minor aches and pains 🔳 headache 🔳 sore throat CRES E temborarily relieves common cold and flu symptoms: Purposes (laptios nose ni) stnaibarpni avitoA Drug Facts (continued) Stoe? gund

cough due to minor throat and bronchial initiation

24

Instal decongestant

Phenylephrine HCl 5 mg.

NDC 83324-145-24

Ask a doctor before use if you have mived risease diabetes heart disease thyroid disease mived risease	Other information
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription moneamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	Directions = do not take more than directed (see Overdose warming) = do not take more than 4 doses in 24 hours = do not take more than 4 doses in 24 hours = adults and children 12 years and over: take 2 softgels with water every 4 hours = swallow whole; do not crush, chew, or dissolve = children under 12 years: do not use = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other Daytime or Nighttime products, carefully read each = when using other correct dosing
Do not use the set short date contraction and set and set and set of the set	adults as well as for children even if you do not notice any signs or symptoms.
may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warming: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or womiting, consult a doctor promptly.	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Oulck medical attention is critical for
Warnings Liver warming: This product contains acetaminophen. Severe liver damage may occur if you take: I more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product 3 or more alcoholic drinks every day while using this product 3 or more alcoholic drinks every day while using this product and the sections. Symptoms	Stop use and ask a doctor if m pain, cough, or nasal congestion gets worse or lasts more than 7 days m nervousness, disziness, or sleeplessness occur m fever gets worse or lasts more than 3 days m tedness or swelling is present m cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
Drug Facts (continued)	Drug Facts (continued)

When using this product, do not exceed recommended dosage.	Call 1-800-935-2362 Monday-Friday 9AM-5PPM EST
Ask a succest or prioritic sector as a successful or an example of the sector of the s	Questions or comments?
Persistent or chronic cough such as occurs with smoking, asthma, or emphysema m cough that occurs with too much philegm (mucus) Ask a doctor or pharmacist before use if you are taking the blood	Inactive ingredients lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink
Ask a doctor before use if you have ■ liver disease ■ diabetes ■ heart disease ■ triyroid disease ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland	Other information a store between 15-30°C (59-86°F) a svoid excessive heat
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a deotor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiator, or emotional conditions, or (certain drugs for depression, psychiator, or emotional conditions, or do not know if your prescription drug contains an MAOI, ask a doctor or do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	Directions — do not take more than directed (see Overdose warming) a do not take more than 4 doses in 24 hours a do not take more than 4 doses in 24 hours a dults and children 12 years and over: take 2 softgels with water every 4 hours a swallow whole; to not crush, chew, or dissolve a when under 12 years: do not use children under 12 years: do not use a when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing
vomična, consult a doctor prompty. Do not use	Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
The provide the provided the pr	If pregnant or breast-feeding, ask a health professional before use. Keep out at reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, of get modical help or confacta 7 Poion Confrol Comport
Allergy alert. Acetaminophen may cause severe skin reactions. Symptoms	signs of a serious condition.

Exp. Date Lot No.:



# DAYTIME COLD AND FLU

acetaminophen dextromethorphan hbr phenylephrine hcl capsule, liquid filled

<b>Product Informati</b>	ion						
Product Type		HUMAN OTC DRUG	Item Code (Sour	ce)	NDC:83324-145	(NDC:5	1013-189)
Route of Administrat	ion	ORAL					
	<b>A</b> . <b>1 .</b>						
Active Ingredient/		-		1			
	-	dient Name			asis of Streng	gth	Strength
ACETAMINOPHEN (UNII:				ACET	AMINOPHEN		325 mg
DEXTROMETHORPHAN I (DEXTROMETHORPHAN - U			19КҮН)		ROMETHORPHAN ROBROMIDE		10 mg
PHENYLEPHRINE HYDRO UNII:1WS297W6MV)	OCHLORI	<b>DE</b> (UNII: 04JA59TNSJ	) (PHENYLEPHRINE -		IYLEPHRINE ROCHLORIDE		5 mg
Inactive Ingredien	ts						
		Ingredient Na	me			St	rength
FD&C RED NO. 40 (UNII:	WZ B912	7XOA)					
FD&C YELLOW NO. 6 (U	JNII: H77V	EI93A8)					
GELATIN (UNII: 2G86QN3	27L)						
GLYCERIN (UNII: PDC6A30							
POLYETHYLENE GLYCO	L, UNSPE	CIFIED (UNII: 3WJQ0	SDW1A)				
POVIDONE (UNII: FZ 989G	•						
PROPYLENE GLYCOL (UI		Q167V3)					
WATER (UNII: 059QF0KOC							
SORBITAN (UNII: 6092IC)							
SORBITOL (UNII: 506T60/							
LECITHIN, SOYBEAN (UN							
LIGHT MINERAL OIL (UN		37QVP)					
MANNITOL (UNII: 30WL53	3L36A)						
Product Character	ristics						
		(a.e. #)	<b>C</b> .				
	orange (c		Score			no scor	e
•	CAPSULE	(obiong)	Size	_		20mm	
Flavor			Imprint Co	de	F	PC9	

CO	ntains			
Pa	ckaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:83324-145- 24	24 in 1 CARTON	07/31/2024	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
		Toddet		
		inoduct		
M		nformation		
M			Marketing Start Date	Marketing End Date

Labeler - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 5/2024

QUALITY CHOICE (Chain Drug Marketing Association)