KROGER DAYTIME COLD AND FLU SOFTGEL- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

The Kroger Co.

Kroger DayTime Cold & Flu SOFTGELS

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg Dextromethorphan HBr 10 mg Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer Cough suppressant Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
 sore throat
- cough due to minor throat & bronchial irritation
- headache minor aches & pains fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

• skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease heart disease diabetes
- high blood pressure
 thyroid disease
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product, do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

• store at room temperature between 20°-25°C (68°-77°F) • protect from light, heat, and moisture

Inactive ingredients

FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol 400,

povidone, propylene glycol, sorbitol sorbitan anhydrous, titanium dioxide

Questions or comments?

1-800-632-6900

COMPARE TO active ingredients of VICKS[®] DAYQUIL[®] COLD & FLU RELIEF LIQUICAPS™*

POWERFUL RELIEF

Multi-Symptom Relief

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

DISTRIBUTED BY THE KROGER CO. CINCINNATI, OHIO 45202 QUALITY GUARANTEE www.kroger.com

PRODUCT OF INDIA

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the trademarks Vicks[®], DayQuil[™] and LiquiCaps [™].


REV.00-092024

PARENTS:

Learn about teen medicine abuse www.StopMedicineAbuse.org

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND INFORMATION

Packaging



Acetaminophen

Dextromethorphan HBr Phenylephrine HCI

Pain Reliever Fever Reducer Cough Suppressant Nasal Deconaestant

48 SOFTGELS

Purpose

Pain Reliever Fever Reducer Cough Suppressant Nasal Decongestant

48 SOFTGELS

nor or priarmaciss. You are now taking a prescription mon use inhibitor (MAOI) (certain drugs for e MAOI drug, If you do not know if your escription drug contains an MAOI, ask a actor or pharmacist before taking this pro Directions

2 softgels with water every 4 hrs ask a docto

READ AND KEEP OUTER CARTON FOR C LOT

DRUG FACTS LABEL

Drug Facts

Active ingredients (in each softgel)

.Pain reliever/fever reduce cetaminophen 325 m Dextromethorphan HBr 10 mg......Cough suppressant Phenylephrine HCl 5 mg..... ...Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
 sore throat
- cough due to minor throat & bronchial irritation ■ headache ■ minor aches & pains ■ fever

Warnings

Liver warning: This product contains acetaminophen Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening
■ blisters
■ rash If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ high blood pressure ■ thyroid disease

Drug Facts (continued)

- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product, do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Drug Facts (continued)

Other information

■ store at room temperature between 20°-25°C (68°-77°F) ■ protect from light, heat, and moisture

Inactive ingredients

FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, sorbitol sorbitan anhydrous, titanium dioxide

Questions or comments? 1-800-632-6900

KROGER DAYTIME COLD AND FLU SOFTGEL

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59450-046
I I OGGE I VPC	TIOT IN ALC TO DITOC	item code (source)	1120.33 130 0 10

Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg		
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg		

Inactive Ingredients				
Ingredient Name	Strength			
FD&C RED NO. 40 (UNII: WZB9127XOA)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)				
GLYCERIN (UNII: PDC6A3C0OX)				
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)				
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SORBITOL (UNII: 506T60A25R)				
SORBITAN (UNII: 6092ICV9RU)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				

Product Characteristics				
Color	red	Score	no score	
Shape	OVAL	Size	19mm	
Flavor		Imprint Code	A09	
Contains				

Packaging					
# Item Code	Package Description	Marketing Start Date	Marketing End Date		
NDC:59450- 046-48	4 in 1 CARTON	01/08/2025			
1	12 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information					
Marketing	Application Number or Monograph	Marketing Start	Marketing End		
Category	Citation	Date	Date		

OTC Monograph Drug	M012	01/08/2025	

Labeler - The Kroger Co. (006999528)

Revised: 1/2025 The Kroger Co.