CHILDRENS COLD AND COUGH - brompheniramine maleate, dextromethorphan, phenylephrine hcl liquid CARDINAL HEALTH 110, LLC. DBA LEADER

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Leader Childrens Cold and Cough 577

ACTIVE INGRADIENTS (in each 10 mL)

Brompheniramine maleate, USP 2 mg
Dextromethorphan HBr, USP 10 mg
Phenylephrine HCl, USP 5 mg

PURPOSES

Antihistamine Cough Suppressant Nasal decongestant

USES

- temporarily relieves cough due to minor bronchial irritation as may occur with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
- sneezing
- Itching of the nose or throat
- runny nose
- itchy, watery eyes
- temporarily restores freer breathing through the nose

WARNINGS

.

DO NOT USE

• to sedate a child or to make a child sleepy

 if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, or psychiatric, for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

ASK A DOCTOR BEFORE USE IF YOU HAVE

- heart disease
- glaucoma
- diabetes
- thyroid disease
- high blood pressure
- difficulty in urination due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

- taking any other oral nasal decongestant or stimulant
- taking sedative or tranquilizers

WHEN USING THIS PRODUCT

- do not take more than directed
- may cause drowsiness
- alcohol, sedatives, and tranquillizers may increase drowsiness
- avoid alcoholic beverages
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

STOP USE AND ASK A DOCTOR IF

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

IF PREGNANT OR BREAST-FEEDING,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-

DIRECTIONS

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not exceed recommended dosage
- mL = milliliter

Age	Dose
adults & children	20 mL
12 years & over	every 4 hours
children 6 to under	10 mL
12 years	every 4 hours
children under 6 years	do not use

OTHER INFORMATION

- each 10 mL contains: sodium 10 mg
- store at 20-25°C (68-77°F)
- measuring cup provided

INACTIVE INGREDIENTS

anhydrous citric acid, artificial flavor, FD&C blue #1, FD&C Red #40, flavor, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose

QUESTIONS OR COMMENTS?

call (609) 860-2600 Mon - Fri 8:00 am to 4:30 pm est

PRINCIPAL DISPLAY PANEL

LEADER

NDC 70000-0631-1

Children's Cold & Cough Liquid

Brompheniramine Maleate

Dextromethorphan Hydrobromide

Phenylephrine Hydrochloride

Antihistamine

Cough Suppressant

Nasal Decongestant

Grape

Flavor

Relieves:

Stuffy Nose,

Runny Nose,

Sneezing,

Cough

Alcohol-Free

COMPARE TO CHILDREN'S DIMETAPP®
COLD & COUGH

active ingredients*

4 FL OZ (118 mL)

LEADER

DO NOT USE IF SEAL UNDER CAP IMPRINTED WITH EALED FOR YOUR PROTECTION" IS BROKEN OR MISS

Drug Facts

Active ingredients (in each 10 mL) Purposes Brompheniramine maleate, USP 2 mg....Antihistamine Dextromethorphan HBr, USP 10 mg...Cough suppressant Phenylephrine HCI, USP 5 mg......Nasal decongestant

Uses

- temporarily relieves cough due to minor bronchial irritation as may occur with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis): * sneezing * itching of the nose or throat * runny nose * itchy, watery eyes temporarily restores freer breathing through the nose

- Do not use

 to sedate a child or to make a child sleepy
 if you are now taking a prescription monoamine
 oxidase inhibitor (MAOI) (certain drugs for
 depression, sychiatric, or remotional conditions,
 or Parkinson's disease), or for 2 weeks after
 stopping the MAOI drug. If you do not know if your
 prescription drug contains an MAOI, ask a doctor
 or pharmacist before taking this product.

- Ask a doctor before use if you have
 heart disease glaucoma diabetes
 thyroid disease inigh blood pressure
 difficulty in urination due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
 a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are ■ taking any other oral nasal decongestant or stimulant ■ taking sedatives or tranquilizers

When using this product do not take more than directed

- may cause drowsiness
 alcohol, sedatives, and tranquilizers may increase drowsiness
 avoid alcoholic beverages
 be careful when driving a motor vehicle or
- operating machinery

Drug Facts (continued)

excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
 symptoms do not improve within 7 days or
 are accompanied by fever
 cough lasts more than 7 days, comes back, or
- is accompanied by fever, rash, or persistent headache. These could be signs of a serious

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of

overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
 measure only with dosing cup provided
 do not exceed recommended dosage
 mL = milliliter

Age	Dose
adults & children	20 mL
12 years & over	every 4 hours
children 6 to under	10 mL
12 years	every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 10 mg store at 20-25°C (68-77°F)
- measuring cup provided

Inactive ingredients anhydrous citric acid, artificial flavor, FD&C blue #1, FD&C red #40, glycerin propylene glycol, propyl gallate, purified water sodium benzoate, sodium citrate, sorbitol, sucralose

Questions or comments? call (609) 860-2600 Mon - Fri 8:00 am to 4:30 pm est

*This product is not manufactured or distributed by Pfizer ConsumerHealthcare, the distributor of Children's Dimetapp® Cold and Cough.

KEEP THIS BOX FOR COMPLETE INSTRUCTIONS

LEADER? LEADER?

NDC 70000-0631-1

Children's Cold & Cough

Brompheniramine Maleate Dextromethorphan HBr Phenylephrine HCI

Antihistamine Cough Suppressant Nasal Decongestant

Grape Flavor

Relieves: Stuffy Nose, Runny Nose, Cough

Alcohol-Free



COMPARE TO CHILDREN'S DIMETAPP* COLD & COUGH

4 FL OZ (118 mL)

Children's

Cold & Cough Liquid

©2023 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, ESSENTIAL TO CARE, LEADER, and the LEADER LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

DISTRIBUTED BY
CARDINAL HEALTH
DUBLIN, 0HIO 43017
WWW.myleader.com
1–800–200-6313
Essential to Care** since 1979

CardinalHealth™

A■ LEADER™ Brand Products Have A 100% Money Back Guarantee Return to place of purchase if not satisfied.

No Coating Area



LOT:

UNVARNISHED AREA TO BE LEFT BLANK FOR LOT & EXP. INFORM

EXP:



CHILDRENS COLD AND COUGH

brompheniramine maleate, dextromethorphan, phenylephrine hcl liquid

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0631	
Route of Administration	ORAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
BROMPHENIRAMINE MALEATE (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII: H57G17P2FN)	BROMPHENIRAMINE MALEATE	2 mg in 10 mL		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL		
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE	5 mg in 10 mL		

Inactive Ingredients				
Ingredient Name	Strength			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				

WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	

Product Characteristics				
Color	PURPLE	Score		
Shape		Size		
Flavor	GRAPE	Imprint Code		
Contains				

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:70000- 0631-1	1 in 1 CARTON	03/30/2023			
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product				

Marketing Information				
Marketing Application Number or Monograph Marketing Start Marketing End Category Citation Date Date				
OTC MONOGRAPH FINAL	part341	03/30/2023		

Labeler - CARDINAL HEALTH 110, LLC. DBA LEADER (063997360)

Establishment				
Name	Address	ID/FEI	Business Operations	
Guardian Drug Company		119210276	MANUFACTURE(70000-0631)	

Revised: 3/2023 CARDINAL HEALTH 110, LLC. DBA LEADER