

# **MUCUS RELIEF COLD AND SINUS MAXIMUM STRENGTH- acetaminophen, guaifenesin, phenylephrine hcl liquid**

**Walgreens**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

## **Drug Facts**

### **Active ingredients (in each 20 mL)**

**Acetaminophen 650 mg**

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

## **Purposes**

### **Pain reliever/fever reducer**

Expectorant

Nasal decongestant

## **Uses**

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - sore throat
  - headache
  - nasal congestion
  - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

## **Warnings**

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- diabetes
- liver disease
- heart disease
- thyroid disease
- high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product,**

**do not use more than directed.**

**Stop use and ask a doctor if**

- redness or swelling is present
- nervousness, dizziness or sleeplessness occur
- new symptoms occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- cough comes back, or occurs with rash or persistent headache

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL=milliliter
- adults and children 12 years and older: 20 mL in dosing cup provided every 4 hours

- children under 12 years of age: do not use

**Other information**

- each 20 mL contains: **sodium 12 mg**
- store at 20-25°C (68-77°F). Do not refrigerate

**Inactive ingredients**

anhydrous citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

**Questions or comments?**

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**Principal Display Panel**

Maximum Strength

Mucus Relief

Cold & Sinus

Acetaminophen/Pain Reliever & Fever Reducer

Guaifenesin/Expectorant

Phenylephrine HCl/Nasal Decongestant

- Nasal & chest congestion
- Headache
- Fever
- Thins & loosens mucus
- For ages 12 & over

Compare to Mucinex® Fast-Max® Cold & Sinus Maximum Strength active ingredients≠

FL OZ (mL)

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD. DEERFIELD, IL 60015

walgreens.com

≠This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Fast-Max® Cold & Sinus Maximum Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL IS BROKEN OR MISSING.

**Package Label**

Well at  
Walgreens

WALGREENS PHARMACIST RECOMMENDED<sup>1</sup>

NDC 0363-3350-06

**MAXIMUM STRENGTH**  
**Mucus Relief**  
**Cold & Sinus**  
**Acetaminophen /**  
**Pain Reliever & Fever Reducer**  
**Guaifenesin / Expectorant**  
**Phenylephrine HCl /**  
**Nasal Decongestant**

- Nasal & chest congestion
- Headache • Fever
- Thins & loosens mucus
- For ages 12 & over

W10040-XXXX-F

Compare to Mucinex<sup>®</sup> Fast-Max<sup>®</sup> Cold & Sinus  
Maximum Strength active ingredients<sup>2†</sup>

PLD-B281B  
LB001972

6 FL OZ (177 mL) LIQUID

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
walgreens.com ©2014 Walgreen Co.  
ITEM 280507



<sup>1</sup>Walgreens Pharmacist Survey Study, November 2012.  
<sup>2†</sup>This product is not manufactured or distributed by Beckitt Benckiser, distributor of Mucinex<sup>®</sup> Fast-Max<sup>®</sup> Cold & Sinus Maximum Strength.

PLD-B281B  
LB001994 **TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL IS BROKEN OR MISSING.**

**Drug Facts**

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - sore throat ■ headache ■ nasal congestion
  - sinus congestion and pressure
  - temporarily reduces fever

PEEL CORNER FOR MORE DRUG FACTS

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 6 doses in 24 hours, which is the maximum daily amount
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks daily while using this product

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age

**Drug Facts (continued)**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease ■ heart disease ■ diabetes
- thyroid disease ■ high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or persistent headache. These could be signs of a serious condition.

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not take more than directed (see Overdose warning)

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- adults and children 12 years and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

**Other information**

- each 20 mL contains: sodium 12 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** anhydrous citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

**Drug Facts (continued)**

**Questions or comments?** Call  
1-877-753-3935 Monday-Friday 9AM-5PM EST

REV0914-RF2

**Well at Walgreens Maximum Strength Mucus Relief Liquid**

**MUCUS RELIEF COLD AND SINUS MAXIMUM STRENGTH**

acetaminophen, guaifenesin, phenylephrine hcl liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-3350
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-3350-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/31/2014	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	10/31/2014	

**Labeler** - Walgreens (008965063)