







- Tell your doctor in the past several days, or you have or discuss with any other drug including warfarin (a type of anticoagulant), other over the counter drugs, and herbs or dietary supplements in addition to your prescription drugs. Tell your doctor about any use of the following over-the-counter drugs, and any use of the following over-the-counter products, including but not limited to aspirin, acetaminophen, ibuprofen, naproxen, salicylates, NSAIDs, antiplatelet drugs, and aspirin. Tell your doctor about any use of the following over-the-counter products, including but not limited to aspirin, acetaminophen, ibuprofen, naproxen, salicylates, NSAIDs, antiplatelet drugs, and aspirin. Tell your doctor about any use of the following over-the-counter products, including but not limited to aspirin, acetaminophen, ibuprofen, naproxen, salicylates, NSAIDs, antiplatelet drugs, and aspirin.
- Check for other side effects:
  - Stomach pain or bloating
  - Bleeding (purple bruising)
- If you are continuing treatment:
  - Avoid any activity or exposure that could result in traumatic injury.
  - Avoid any activity or exposure that could result in traumatic injury.
  - Avoid any activity or exposure that could result in traumatic injury.
  - Avoid any activity or exposure that could result in traumatic injury.

**Manufacturer:**

**Call for Information:**  
 Call for information 1-800-838-1161  
**Distributed by:**  
 Zilex Pharmaceuticals USA, Inc.  
 15904 Valley Blvd.  
 Van Nuys, CA 91410  
 818-708-2200  
 Fax: 818-708-2205  
 Website: www.zilex.com

**INDICATIONS:**

**Warfarin Sodium USP, USP**  
 Warfarin sodium USP is indicated for the treatment of thromboembolic disorders, such as pulmonary embolism and deep vein thromboses, and for the prevention of thromboembolism in patients with certain types of heart disease. It is also used to prevent blood clots in patients with atrial fibrillation and to prevent stroke in patients with certain types of heart disease.

**What is the most important information I should know about warfarin sodium?**

- You may have a higher risk of bleeding if you are taking warfarin sodium if you have any of the following conditions or if you are taking any of the following medications:
  - are 65 years of age or older
  - have a history of stomach or intestinal bleeding
  - have high blood pressure (hypertension)
  - have a history of stroke or "mini-strokes" (transient ischemic attack or TIA)
  - have urinary tract disease
  - have a low blood count or cancer
  - have had trauma, such as an accident or surgery
  - have kidney problems
  - have other medicines that increase your risk of bleeding, including:
    - aspirin and other nonsteroidal anti-inflammatory drugs (NSAIDs)
    - other medicines to prevent or treat blood clots
    - tramadol and other pain relievers (opioids)

**Tell your healthcare provider if you take any of these medicines. Ask your healthcare provider if you should stop or change the dose of any of these medicines. Tell your healthcare provider if you are taking any of the following medicines:**

- anti-infective medicines, including penicillins, cephalosporins, and others
- anti-infective medicines, including penicillins, cephalosporins, and others
- anti-infective medicines, including penicillins, cephalosporins, and others
- anti-infective medicines, including penicillins, cephalosporins, and others

**Get your regular blood tests to check for your response to warfarin sodium.**

The healthcare provider will check your blood clotting time before you start taking warfarin sodium. Your healthcare provider will check your blood clotting time before you start taking warfarin sodium. Your healthcare provider will check your blood clotting time before you start taking warfarin sodium.

- Tell your healthcare provider if you are taking any of the following medicines, including:
  - aspirin and other nonsteroidal anti-inflammatory drugs (NSAIDs)
  - other medicines to prevent or treat blood clots
  - tramadol and other pain relievers (opioids)

**Tell your healthcare provider if you have any of the following conditions:**

- bleeding problems, such as nosebleeds, bleeding gums, or bleeding from cuts
- bleeding problems, such as nosebleeds, bleeding gums, or bleeding from cuts
- bleeding problems, such as nosebleeds, bleeding gums, or bleeding from cuts
- bleeding problems, such as nosebleeds, bleeding gums, or bleeding from cuts

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems

**Do not take warfarin sodium if you have any of the following conditions:**

- active bleeding problems
- active bleeding problems
- active bleeding problems
- active bleeding problems