

**ALLERGY MULTI-SYMPTOM- acetaminophen, chlorpheniramine maleate, phenylephrine hcl tablet, film coated**  
**FRED'S, INC.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Freds Pharmacy 44-455C-AMS**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Chlorpheniramine maleate 2 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever  
Antihistamine  
Nasal decongestant

***Uses***

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
  - headache
  - nasal congestion
  - runny nose and sneezing
  - minor aches and pains
  - sinus congestion and pressure
- temporarily relieves these additional symptoms of hay fever:
  - itchy, watery eyes
  - itching of the nose or throat
- helps clear nasal passages
- helps decongest sinus openings and passages

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

***Do not use***

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- glaucoma
- high blood pressure
- a breathing problem such as emphysema or chronic bronchitis
- heart disease
- thyroid disease
- diabetes
- liver disease

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery
- may cause drowsiness

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed**
- adults and children 12 years and over
  - take 2 caplets every 4 hours
  - swallow whole - do not crush, chew, or dissolve
  - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

***Inactive ingredients***

corn starch, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silica gel, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

***Questions or comments?***

**1-800-426-9391**

***Principal Display Panel***

**fred's  
Pharmacy**

NDC 55315-456-08

**ACTUAL  
SIZE**

**ALLERGY  
MULTI-SYMPTOM**

**Acetaminophen**  
*Chlorpheniramine maleate*  
*Phenylephrine HCl*

*Pain Reliever, Antihistamine, Nasal Decongestant*

- *Headache* • *Sinus Pressure* • *Nasal Congestion*
- *Runny Nose & Sneezing* • *Itchy, Watery Eyes*

**24 COOL CAPLETS  
with COOL BLAST FLAVOR**

*Compare to the active ingredients in:*

**Tylenol® Allergy**  
Multi-Symptom\*

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR  
SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by McNeil  
Consumer Healthcare, owner of the registered trademark Tylenol®  
Allergy Multi-Symptom. 50844 REV0616F45508

DISTRIBUTED BY: fred's, Inc.  
4300 NEW GETWELL RD, MEMPHIS, TN 38118  
www.fredsinc.com

*100% satisfaction guaranteed*  
Questions or comments: 1-855-331-FRED (3733)





NDC 55315-456-08

# ALLERGY MULTI-SYMP TOM

**Acetaminophen**  
**Chlorpheniramine maleate**  
**Phenylephrine HCl**

**Pain Reliever, Antihistamine, Nasal Decongestant**

- Headache • Sinus Pressure • Nasal Congestion
- Runny Nose & Sneezing • Itchy, Watery Eyes

**24 COOL CAPLETS**  
with COOL BLAST FLAVOR

Compare to the active ingredients in:

**Tylenol® Allergy  
Multi-Symptom®**



ACTUAL  
SIZE

**Drug Facts**  
KEEP OUTER PACKAGE FOR COMPLETE  
PRODUCT INFORMATION

**Active ingredients (in each caplet)**  
Acetaminophen 325 mg ..... Pain reliever  
Chlorpheniramine maleate 2 mg ..... Antihistamine  
Phenylephrine HCl 5 mg ..... Nasal decongestant

**Purpose**  
temporarily relieves these symptoms of hay fever or other upper respiratory allergies:  
■ runny nose and sneezing  
■ itchy, watery eyes  
■ nasal congestion  
■ headache

**Uses**  
temporarily relieves these symptoms of sinus congestion and pressure  
■ minor aches and pains  
■ sore throat

**Drug Facts (continued)**  
**Questions or comments? 1-800-426-9391**

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Allergy Multi-Symptom. 50844 REV0616F45508



DISTRIBUTED BY: Fred's, Inc.  
4300 NEW GETWELL RD, MEMPHIS, TN 38118  
www.fredsinc.com

**100% satisfaction guaranteed**  
Questions or comments: 1-800-331-4262 (D732)

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

No print/No varnish  
Lot & Exp date

**Drug Facts (continued)**

**Other information**  
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BUSTER IS TORN OR BROKEN  
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Drug Facts (continued)**  
see end flap for expiration date and lot number

**Inactive ingredients**  
corn starch, croscollon, hydroxypropyl methylcellulose, polyethylene glycol, polyvinyl alcohol, povidone, silica gel, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

**Do not use**  
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have** ■ difficulty in urination due to enlargement of the prostate gland


**Directions**  
■ do not take more than directed  
■ adults and children 12 years and over  
■ take 2 caplets every 4 hours  
■ swallow whole - do not crush, chew, or dissolve  
■ do not take more than 10 caplets in 24 hours  
■ children under 12 years: ask a doctor

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**If pregnant or breast-feeding,** ask a health professional before use.

■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ if pregnant or breast-feeding, ask a health professional before use.

B-0510-455C-08AMSR  
REV0616F45508



Freds Pharmacy 44-455C-AMS

## ALLERGY MULTI-SYMP TOM

acetaminophen, chlorpheniramine maleate, phenylephrine hcl tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55315-456
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	

Product Characteristics			
Color	WHITE	Score	no score
Shape	OVAL	Size	17mm
Flavor	MINT	Imprint Code	44;455
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55315-456-08	2 in 1 CARTON	06/28/2005	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	06/28/2005	

**Labeler** - FRED'S, INC. (005866116)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(55315-456)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(55315-456)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(55315-456)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(55315-456)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(55315-456)