

**TRICODE AR- chlorpheniramine maleate, codeine phosphate, pseudoephedrine hydrochloride liquid**

**Respa Pharmaceuticals, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Tricode AR**

**Drug Facts**

**Active ingredients**

**(in each 5 mL teaspoonful)**

Chlorpheniramine Maleate 2 mg

Codeine Phosphate\* 8 mg

**\*WARNING: May be habit forming**

Pseudoephedrine Hydrochloride 30 mg

**Purpose**

Antihistamine

Antitussive (Cough Suppressant)

Nasal Decongestant

**Uses**

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- cough due to minor throat and bronchial irritation
- nasal congestion
- reduces swelling of nasal passages

**Warnings**

**Do not exceed recommended dosage.**

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)

- chronic pulmonary disease or shortness of breath, or children who are taking other drugs
- heart disease
- high blood pressure
- thyroid disease
- diabetes

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.**

**When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.
- may cause or aggravate constipation
- nervousness, dizziness, or sleeplessness occur
- new symptoms occur

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of the reach of children.**

**In case of overdose, get medical help or contact a Poison Control Center right away.**

**Directions**

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 4 to 6 hours, not to exceed 8 teaspoonfuls in a 24 hour period
Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 4 teaspoonfuls in a 24 hour period

Children under 6 years of age:	Consult a doctor.
-----------------------------------	-------------------

A special measuring device should be used to give an accurate dose of this product to children under 6 years of age. Giving a higher dose than recommended by a doctor could result in serious side effects for your child.

**Other information**

Store at 15°- 30°C (59°- 86°F)

**Inactive ingredients**

Berry Vanilla Flavor, Citric Acid, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

**Questions? Comments?**

Call (630) 543-3333

**Rx Only**

**CV**

Manufactured for:

**RESPA<sup>®</sup>**  
**PHARMACEUTICALS, INC.**  
Addison, IL 60101

**Product Packaging**

The packaging below represents the labeling currently used:

Principal display panel and side panel for 473 mL label:

**NDC 60575-313-16**

---

**Tricode AR<sup>®</sup>**

**Antihistamine•Cough Suppressant**  
**Nasal Decongestant**

---

**Each 5 mL (one teaspoonful) for oral administration contains:**

Chlorpheniramine Maleate.....2 mg  
Codeine Phosphate\*.....8 mg  
\*(Warning: May be habit-forming)  
Pseudoephedrine HCl.....30 mg

Dye Free / Sugar Free  
Alcohol Free / Gluten Free

**Berry Vanilla Flavor**

**Rx Only**

**RESPA®  
PHARMACEUTICALS, INC.  
Addison, IL 60101**

**16 fl oz. (473 mL)**

**SEE ATTACHED PRODUCT INSERT FOR COMPLETE  
DOSING INFORMATION.**

Tamper evident by foil seal under cap. Do not use if foil  
seal is broken or missing.

**THIS BOTTLE IS NOT TO BE DISPENSED TO THE CONSUMER.**

Dispense in a tight, light-resistant container as defined in  
the USP/NF with a child-resistant closure.

Store at controlled room temperature, 15°- 30°C (59°- 86°F).

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH  
OF CHILDREN.**

Manufactured for:  
Respa Pharmaceuticals  
Addison, IL 60101

**Rx Only**

Rev. 08/11

SEE ATTACHED PRODUCT INSERT FOR COMPLETE DOSING INFORMATION.

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

**THIS BOTTLE IS NOT TO BE DISPENSED TO THE CONSUMER.**

Dispense in a tight, light-resistant container as defined in the USP/NF with a child-resistant closure.

Store at controlled room temperature, 15°-30°C (59°-86°F).

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

Manufactured for:  
Respa Pharmaceuticals  
Addison, IL 60101

**Rx Only**

Rev. 08/11



NDC 60575-313-16

# Tricode® AR

Antihistamine • Cough Suppressant  
Nasal Decongestant

Each 5 mL (one teaspoonful) for oral administration contains:  
Chlorpheniramine Maleate ..... 2 mg  
Codeine Phosphate\* ..... 8 mg  
\*(Warning: May be habit-forming)  
Pseudoephedrine HCl ..... 30 mg

Dye Free / Sugar Free  
Alcohol Free / Gluten Free

Berry Vanilla Flavor

Rx Only

**RESPA®**  
PHARMACEUTICALS, INC.

Addison, IL 60101

16 fl oz. (473 mL)

## Drug Facts   Lift Here

### Active ingredients   Purpose (in each 5 mL teaspoonful)

Chlorpheniramine Maleate 2 mg .....	Antihistamine
Codeine Phosphate* 8 mg .....	Antitussive
*(Warning: May be habit forming)	
Pseudoephedrine Hydrochloride 30 mg ..	Nasal Decongestant

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:  
■ runny nose ■ sneezing ■ Itching of the nose or throat ■ Itchy, watery eyes  
■ cough due to minor throat and bronchial irritation ■ nasal congestion  
■ reduces swelling of nasal passages

### Warnings

Do not exceed recommended dosage.

#### Do not use this product

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

#### Ask a doctor before use if you have

■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma  
■ trouble urinating due to an enlarged prostate gland  
■ a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ a cough that occurs with too much phlegm (mucus)</li> <li>■ chronic pulmonary disease or shortness of breath, or children who are taking other drugs</li> <li>■ heart disease ■ high blood pressure</li> <li>■ thyroid disease ■ diabetes</li> </ul> <p>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.</p> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>■ excitability may occur, especially in children</li> <li>■ may cause marked drowsiness</li> <li>■ avoid alcoholic drinks</li> <li>■ alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>■ be careful when driving a motor vehicle or operating machinery</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.</li> <li>■ may cause or aggravate constipation</li> <li>■ nervousness, dizziness, or sleeplessness occur</li> <li>■ new symptoms occur</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b> Do not exceed recommended dosage.</p>	<p><b>Drug Facts (continued)</b></p> <table border="1"> <tr> <td>Adults and children 12 years of age and over:</td> <td>2 teaspoonsful (10 mL) every 4 to 6 hours, not to exceed 8 teaspoonsful in a 24 hour period</td> </tr> <tr> <td>Children 6 to under 12 years of age:</td> <td>1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 4 teaspoonsful in a 24 hour period</td> </tr> <tr> <td>Children under 6 years of age:</td> <td>Consult a doctor.</td> </tr> </table> <p>A special measuring device should be used to give an accurate dose of this product to children under 6 years of age. Giving a higher dose than recommended by a doctor could result in serious side effects for your child.</p> <p><b>Other information</b> Store at 15°-30°C (59°-86°F)</p> <p><b>Inactive ingredients</b> Berry Vanilla Flavor, Citric Acid, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.</p> <p><b>Questions? Comments?</b> Call (630) 543-3333</p> <p>Rx Only</p>  <p>Manufactured for: <b>RESPA</b> Pharmaceuticals, Inc. Addison, IL 60101</p>	Adults and children 12 years of age and over:	2 teaspoonsful (10 mL) every 4 to 6 hours, not to exceed 8 teaspoonsful in a 24 hour period	Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 4 teaspoonsful in a 24 hour period	Children under 6 years of age:	Consult a doctor.
Adults and children 12 years of age and over:	2 teaspoonsful (10 mL) every 4 to 6 hours, not to exceed 8 teaspoonsful in a 24 hour period						
Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 4 teaspoonsful in a 24 hour period						
Children under 6 years of age:	Consult a doctor.						

## TRICODE AR

chlorpheniramine maleate, codeine phosphate, pseudoephedrine hydrochloride liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:60575-313
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	CV

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Chlorpheniramine Maleate</b> (UNII: V1Q0O9OJ9Z) (Chlorpheniramine - UNII:3U6IO1965U)	Chlorpheniramine Maleate	2 mg in 5 mL
<b>Codeine Phosphate</b> (UNII: GSL05Y1MN6) (Codeine - UNII:Q830PW7520)	Codeine Phosphate	8 mg in 5 mL
<b>Pseudoephedrine Hydrochloride</b> (UNII: 6V9V2RYJ8N) (Pseudoephedrine - UNII:7CUC9DD9F)	Pseudoephedrine Hydrochloride	30 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
Citric Acid (UNII: 2968PHW8QP)	
Glycerin (UNII: PDC6A3C0OX)	
Propylene Glycol (UNII: 6DC9Q167V3)	
Water (UNII: 059QF0KO0R)	
Sodium Citrate (UNII: 1Q73Q2JULR)	
Saccharin Sodium (UNII: SB8ZUX40TY)	
Sorbitol (UNII: 506T60A25R)	

### Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY, VANILLA	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60575-313-16	473 mL in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	06/20/2011	

**Labeler** - Respa Pharmaceuticals, Inc. (883901019)

Revised: 9/2011

Respa Pharmaceuticals, Inc.