

**ASPIRIN- aspirin tablet, film coated**  
**HF Acquisition Co LLC, DBA HealthFirst**

-----  
Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

Pain reliever/fever reduce

**USES**

temporarily relieves:

headache

pain and fever of colds

toothache

menstrual pain

muscle pain

minor pain of arthritis

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

shock

facial swelling

hives

asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

are age 60 or older

have had stomach ulcers or bleeding problems

take a blood thinning (anticoagulant) or steroid drug

take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

have 3 or more alcoholic drinks every day while using this product

take more or for a longer time than directed

Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer

if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

stomach bleeding warning applies to you

you have a history of stomach problems, such as heartburn

you have asthma

you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

you are taking a diuretic

Ask a doctor or pharmacist before use if you are  
taking a prescription drug for

gout  
diabetes  
arthritis

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away.  
you experience any of the following signs of stomach bleeding:

feel faint  
vomit blood  
have bloody or black stools  
have stomach pain that does not get better  
ringing in the ears or a loss of hearing occurs  
pain gets worse or lasts more than 10 days  
fever gets worse or lasts more than 3 days  
new symptoms occur  
redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

## **DIRECTIONS**

do not take more than directed  
drink a full glass of water with each dose  
adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor  
children under 12 years: do not use unless directed by a doctor

store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
see end flap for expiration date and lot number

corn starch, hypromellose, polyethylene glycol, propylene glycol

1-800-645-2158

Ask a doctor before use if

stomach bleeding warning applies to you  
you have a history of stomach problems, such as heartburn  
you have asthma

you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
you are taking a diuretic

Ask a doctor or pharmacist before use if you are  
taking a prescription drug for

gout  
diabetes  
arthritis

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away.  
you experience any of the following signs of stomach bleeding:

feel faint  
vomit blood  
have bloody or black stools  
have stomach pain that does not get better  
ringing in the ears or a loss of hearing occurs  
pain gets worse or lasts more than 10 days  
fever gets worse or lasts more than 3 days  
new symptoms occur  
redness or swelling is present

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away.  
you experience any of the following signs of stomach bleeding:

feel faint  
vomit blood  
have bloody or black stools  
have stomach pain that does not get better  
ringing in the ears or a loss of hearing occurs  
pain gets worse or lasts more than 10 days  
fever gets worse or lasts more than 3 days  
new symptoms occur  
redness or swelling is present

do not take more than directed

drink a full glass of water with each dose

adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor  
children under 12 years: do not use unless directed by a doctor

temporarily relieves: headache - pain and fever of colds - toothache - menstrual pain - muscle pain - minor pain of arthritis



ASPIRIN

aspirin tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51662-1494(NDC:0536-1054)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg
Inactive Ingredients			
Ingredient Name			Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
STARCH, CORN (UNII: O8232NY3SJ)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			

Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;157;ASPIRIN	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51662-1494-1	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013		02/10/1996	

**Labeler** - HF Acquisition Co LLC, DBA HealthFirst (045657305)

**Registrant** - HF Acquisition Co LLC, DBA HealthFirst (045657305)

Establishment			
Name	Address	ID/FEI	Business Operations
HF Acquisition Co LLC, DBA HealthFirst		045657305	relabel(51662-1494)

Revised: 10/2024

HF Acquisition Co LLC, DBA HealthFirst