

WALGREENS MAX STRENGTH BURN GEL- benzalkonium chloride, lidocaine hydrochloride gel
Walgreens Co.

Walgreens Max Strength Burn Gel

Drug Facts

Active Ingredients

Benzalkonium Cl 0.13%

Lidocaine HCl 4%

Purpose

First aid antiseptic

Topical pain relief

Uses:

First aid to help prevent bacterial contamination or skin infection, and for temporary relief of pain and itching associated with minor:

- Burns
- Cuts
- Sunburn
- Scrapes
- Minor Skin Irritation

Warnings

For external use only

Ask a doctor before use if you have:

- Deep or puncture wounds
- Animal bite
- Serious burns

Do not use:

- In or near the eyes
- In large quantities or over large areas of the body
- On raw surfaces or blistered areas

Stop use and ask a doctor if:

- Condition worsens
- Symptoms last more than 7 days or clear up and occur again within a few days

If pregnant or breast-feeding,

ask a health care professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed, get medical help or contact a Poison Control Center immediately.

Directions:**Adults and children 2 years of age & older:**

- Clean the affected area. Apply a small amount of gel on the affected area 1 to 3 times daily.
- May be covered with a sterile bandage (let dry first).
- cover with adhesive bandage or sterile gauze if needed
- apply 1 to 3 times daily.

Children under 2 years of age:

- ask a doctor

Other information:

- Store at room temperature 15° -30°C (59° -86°F)

Inactive ingredients:

Aloe Barbadensis Leaf Juice, Caprylyl Glycol, Chlorophenesin, Dimethyl Isosorbide, Hydroxyethyl Cellulose, Phenoxyethanol, Propanediol, Tocopheryl Acetate (Vitamin E), Water.

Package Labeling:

Walgreens



Burn Gel

BENZALKONIUM Cl 0.13% /
FIRST AID ANTISEPTIC
LIDOCAINE HCl 4% / TOPICAL PAIN RELIEF

Walgreens Burn Gel

Apply Walgreens Burn Gel to the minor burn area. Repeat application as often as necessary (no more than 3 times daily) to temporarily relieve pain and itching.

Walgreens



Compare to the active ingredients in Alocane® Max*

Burn Gel

BENZALKONIUM Cl 0.13% /
FIRST AID ANTISEPTIC
LIDOCAINE HCl 4% /
TOPICAL PAIN RELIEF

Maximum Strength*

+ Aloe

- Fast pain & itch relief
- For use on minor cuts, scrapes & burns
- Kills 99% of germs
- No-sting formula

*Available without a prescription.



*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

††This product is not manufactured or distributed by Quest Products, LLC, owner of the registered trademark Alocane®.

DISTRIBUTED BY: WALGREEN CO.
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com 1-800-925-4733
©2024 Walgreen Co. MADE IN U.S.A.



2.5 FL OZ (75 mL)

Drug Facts

Active Ingredients	Purpose
Benzalkonium Cl 0.13%	First aid antiseptic
Lidocaine HCl 4%	Topical pain relief

Uses:

First aid to help prevent bacterial contamination or skin infection, and for temporary relief of pain and itching associated with minor:

- Burns
- Cuts
- Sunburn
- Scrapes
- Minor Skin Irritation

Warnings:

For external use only

Ask a doctor before use if you have:

- Deep or puncture wounds
- Animal bite
- Serious burns

Do not use:

- In or near the eyes
- In large quantities or over large areas of the body
- On raw surfaces or blistered areas

Stop use and ask a doctor if:

- Condition worsens
- Symptoms last more than 7 days or clear up and occur again within a few days

If pregnant or breast-feeding, ask a health care professional before use.

KEEP OUT OF REACH OF CHILDREN.

If swallowed, get medical help or contact a Poison Control Center immediately.

Directions:

Adults and children 2 years of age & older:

- Clean the affected area. Apply a small amount of gel on the affected area 1 to 3 times daily.
- May be covered with a sterile bandage (let dry first).
- cover with adhesive bandage or sterile gauze if needed
- apply 1 to 3 times daily.

Children under 2 years of age:

- ask a doctor

Other information:

- Store at room temperature 15° -30°C (59° -86°F)

Inactive ingredients:

Aloe Barbadosis Leaf Juice, Caprylyl Glycol, Chlorophenesin, Dimethyl Isosorbide, Hydroxyethyl Cellulose, Phenoxyethanol, Propanediol, Tocopheryl Acetate (Vitamin E), Water.

NDC 0363-9654-10

W3ORG1224-F

ITEM 897407 W00000-0000-0



LOT
EXP

WALGREENS MAX STRENGTH BURN GEL

benzalkonium chloride, lidocaine hydrochloride gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9654
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	13 mg in 1 mL
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	40 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ALOE VERA LEAF JUICE (UNII: RUE8E6T4NB)	
CAPRYLYL GLYCOL (UNII: 00YIU5438U)	
CHLORPHENESIN (UNII: I670DAL4SZ)	
DIMETHYL ISOSORBIDE (UNII: SA6A6V432S)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
PROPANEDIOL (UNII: 5965N8W85T)	
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9654-10	1 in 1 CARTON	03/01/2025	
1		75 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	03/01/2025	

Labeler - Walgreens Co. (008965063)