

UP AND UP CHILDRENS ACETAMINOPHEN- acetaminophen suspension
Target Corporation

Target Corporation Children's Acetaminophen Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL

72-95	11 years	15 mL
-------	----------	-------

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- **each 5 mL contains:** sodium 3 mg
- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-888-547-7400

Principal Display Panel

Compare to active ingredient in Children's Tylenol® Oral Suspension

For Ages 2 To 11 Years

Children's Acetaminophen 160 mg / 5 mL

Oral Suspension

Fever Reducer / Pain Reliever

- Alcohol, ibuprofen and aspirin free
- Paraben free

Grape Flavor

Ages 2 to 11 Years

Dosage Cup Included

up & up™

4 FL OZ (118 mL)



For Ages 2 to 11 Years
Children's Acetaminophen 160 mg / 5 mL
 Oral Suspension
 Fever Reducer / Pain Reliever



Compare to active ingredient in Children's Tylenol® Oral Suspension**

For Ages 2 to 11 Years
Children's Acetaminophen 160 mg / 5 mL
 Oral Suspension
 Fever Reducer / Pain Reliever

- Alcohol, ibuprofen and aspirin free
- Paraben free



4 FL OZ (118 mL)

Grape Flavor

Ages 2 to 11 Years

Dosage Cup Included

Directions:

Step 1
Shake well before using.



Step 2
Remove child protective cap.



Step 3
Squeeze your child's dose into the dosing cup.



IMPORTANT:
Please see dosing chart or ask a doctor for your child's correct dose.

Drug Facts

Active ingredient (in each 5 mL) Purpose
 Acetaminophen 160 mg.....Pain reliever/fever reducer

- Uses temporarily:**
- reduces fever
 - relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blisters
 - rash
 If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

- Do not use**
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease
 Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

- Stop use and ask a doctor if**
- pain gets worse or lasts more than 5 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)

- Directions**
- this product does not contain directions or complete warnings for adult use
 - do not give more than directed (see overdose warning)
 - shake well before using
 - mL = milliliter
 - find right dose on chart below. If possible, use weight to dose; otherwise, use age.
 - remove the child protective cap and squeeze your child's dose into the dosing cup
 - repeat dose every 4 hours while symptoms last
 - do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)**
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

**or as directed by a doctor
Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

- Other information**
- each 5 mL contains: sodium 3 mg
 - store at 20-25°C (68-77°F)
 - do not use if printed neckband is broken or missing

Inactive ingredients
 anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments? 1-888-547-7400

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Tylenol® Oral Suspension.

Distributed by Target Corporation
 Minneapolis, MN 55403
 TM & ©2024 Target Brands, Inc.

NDC 11673-130-26



13026 UW C12

UP AND UP CHILDRENS ACETAMINOPHEN

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-130
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TGW28D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics

Color	PURPLE (viscous)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-130-26	1 in 1 CARTON	07/08/2009	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11673-130-28	1 in 1 CARTON	08/30/2012	06/27/2014
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:11673-130-34	2 in 1 CARTON	07/08/2009	07/08/2009
3		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug	M013	07/08/2009	
--------------------	------	------------	--

Labeler - Target Corporation (006961700)

Revised: 7/2025

Target Corporation