

CHILDREN ACETAMINOPHEN- acetaminophen liquid
AARNA USA INC.

Children's Acetaminophen
Oral Liquid
Alcohol-free and Aspirin-free
Pain Reliever and fever reducer
Acetaminophen 160 mg per 5 mL
Alcohol Free, Aspirin Free
For Ages 2 to 11 years

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain Reliever/Fever Reducer

Uses temporarily

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- If a skin reaction occurs, stop use and seek medical help right away.

Soar throat warning: if sore throat is severe, persists or more than 2 days, is accompanied or followed by fever headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product: Do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use

Overdose warning. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain direction or complete warnings for adult use**
- **Shake well before using**
- ml = milliliter
- find right dose on chart below
- if possible, use weight to dose; otherwise use age
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

Other Information store at room temperature 20 °-25 °C (68 °-77 °F). Protect from Freezing. Protect from light.

Inactive ingredients: Anhydrous Citric Acid, Bubble gum flavor, FD&C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

Questions or comments? call toll-free 1-877-225-6999.

Leland, NC-28451

This product is not manufactured by or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol Elixir.

**Bubble Gum
Flavor**

NDC: 82568-0008-4

Children's Acetaminophen Oral Liquid

160mg per 5mL

PAIN RELIEVER
and fever Reducer

Alcohol Free Aspirin Free

For Ages 2 to 11 Years

Dose Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg	Pain reliever / fever reducer
----------------------	-------------------------------

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to:
■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:
 - more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen.**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning).

Stop use and ask a doctor if: pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (**1-800-222-1222**) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions ■ This product does not contain directions or complete warnings for adult use ■ shake well before using ■ mL = milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ use only the endorsed dosing cap designed for use with product.

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)	Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor	48 to 59	6 to 8	2 tsp or 10 mL
24 to 35	2 to 3	1 tsp or 5 mL	60 to 71	9 to 10	2 ½ tsp or 12.5 mL
36 to 47	4 to 5	1 ½ tsp or 7.5 mL	72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Other information store at room temperature 20°-25°C (68°-77°F). Protect from Freezing & Light.

Inactive Ingredients Anhydrous Citric Acid, Bubble gum flavor, F O C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Chloride, Sorbitol Solution, Sucralose.

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark "Tylenol".
***Temporary Exempted: Do Not Use if Printed Safety Seal Around Bottle or Under Cap Is Broken or Missing.

LOT NO.
EXPDATE.

Aarna USA
Manufactured for:
AARNA USA Inc.
Leland, NC 28451

KTK/25/666/2017

KTG2568 00084 6

Children's Acetaminophen Oral Liquid

160mg per 5mL



NDC: 82568-0008-4

Drug Facts

Active ingredient (in each 5 mL) **Purpose**
Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:
■ more than 5 doses in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen.

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days.
■ new symptoms occur ■ redness or swelling is present
These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

■ this product does not contain directions or complete warnings for adult use, shake well before using
■ mL = milliliter
■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.
■ use only the enclosed dosing cup designed for use with product. Do not use any other dosing device.
■ if needed, repeat dose every 4 hours while symptoms last

NDC: 82568-0008-4



NDC: 82568-0008-4



Children's Acetaminophen Oral Liquid

160mg per 5mL

PAIN RELIEVER
and fever Reducer

Alcohol Free
Aspirin Free

For Ages
2 to 11 Years

Children's Acetaminophen Oral Liquid

160mg per 5mL

PAIN RELIEVER
and fever Reducer

Alcohol Free
Aspirin Free

For Ages
2 to 11 Years



4 FL.OZ (118 mL)



4 FL.OZ (118 mL)

Drug Facts (continued)

■ do not give more than 5 times in 24 hours
■ do not give more than 5 days unless directed by doctor
■ do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Other Information

store at room temperature 20-25°C (68-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients

Anhydrous Citric Acid, Bubble gum flavor, F D & C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

Questions or comments?

call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elivir®

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Aarna USA
Manufactured for:
AARNA USA Inc.
Leland, NC 28451



K10A-C2501-00

NDC: 82568-0008-8



Children's Acetaminophen Oral Liquid

160mg per 5mL

PAIN RELIEVER
and fever Reducer

Alcohol Free
Aspirin Free

For Ages
2 to 11 Years

Bubble Gum
Flavor

8 FL.OZ (236 mL)

Drug Facts

Active ingredient (in each 5 mL) Purpose

Acetaminophen 160 mgPain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold
■ flu ■ headache ■ sore throat ■ toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

■ this product does not contain directions or complete warnings for adult use, shake well before using ■ mL = milliliter

■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

■ use only the enclosed dosing cup designed for use with product.

Do not use any other dosing device. ■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours.

Drug Facts (continued)

■ do not give more than 5 days unless directed by doctor.
■ do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Other Information


store at room temperature 20-25°C (68-77°F). Protect from Freezing.
Protect from Light.

Inactive Ingredients Anhydrous Citric Acid, Bubble gum flavor, FD & C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elvir®

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.



Aarna USA
Manufactured for:
AARNA USA Inc.
Leland, NC-28451



K10B-L2501-00

Children's Acetaminophen Oral Liquid

160mg per 5mL



NDC: 82568-0008-8

Drug Facts

Active ingredient (in each 5 mL) **Purpose**
Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:
■ more than 5 doses in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen.

Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days.
■ new symptoms occur ■ redness or swelling is present
These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use, shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- use only the enclosed dosing cup designed for use with product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last

NDC: 82568-0008-8



Children's Acetaminophen Oral Liquid

160mg per 5mL

PAIN RELIEVER
and fever Reducer

Alcohol Free
Aspirin Free

For Ages
2 to 11 Years



8 FL.OZ (236 mL)

NDC: 82568-0008-8



Children's Acetaminophen Oral Liquid

160mg per 5mL

PAIN RELIEVER
and fever Reducer

Alcohol Free
Aspirin Free

For Ages
2 to 11 Years



8 FL.OZ (236 mL)

Drug Facts (continued)

- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by doctor
- do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Other Information

store at room temperature 20-25°C (68-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients Anhydrous Citric Acid, Bubble gum flavor, F D & C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

Questions or comments?

call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Extra

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Aarna USA
Manufactured by:
AARNA USA Inc.
Leland, NC 28451



Drug Facts

Active ingredient (in each 5 mL) **Purpose**

Acetaminophen 160 mgPain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to:
■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if
■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur redness or swelling is present These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions
■ this product does not contain directions or complete warnings for adult use, shake well before using ■ mL = milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

NDC: 82568-0008-6

Children's Acetaminophen Oral Liquid

160mg per 5mL

PAIN RELIEVER
and fever Reducer

Alcohol Free
Aspirin Free

For Ages
2 to 11 Years

Bubble Gum Flavor

16 FL.OZ (473 mL)

Drug Facts (continued)

■ if needed, repeat dose every 4 hours while symptoms last
■ do not give more than 5 times in 24 hours
■ do not give more than 5 days unless directed by doctor.
■ do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Other Information
store at room temperature 20-25°C (68-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients Anhydrous Citric Acid, Bubble gum flavor, F D & C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elixir

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Aarna USA
Manufactured for:
AARNA USA Inc.
Leland, NC-28451

KTK/25/666/2017

N 3 82568 00086 0

CHILDREN ACETAMINOPHEN			
acetaminophen liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82568-0008
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	160 mg in 5 mL
Inactive Ingredients			
Ingredient Name			Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
FD&C RED NO. 40 (UNII: WZ B9127XOA)			
GLYCERIN (UNII: PDC6A3C0OX)			
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SODIUM CITRATE (UNII: 1Q73Q2JULR)			
SORBITOL SOLUTION (UNII: 8KW3E207O2)			

SUCRALOSE (UNII: 96K6UQ3ZD4)

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82568-0008-4	1 in 1 PACKAGE	02/18/2025	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:82568-0008-8	1 in 1 PACKAGE	02/18/2025	
2		236 mL in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:82568-0008-6	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/18/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/18/2025	

Labeler - AARNA USA INC. (118515992)

Revised: 5/2025

AARNA USA INC.