

**ANBESOL COLD SORE THERAPY- allantion, benzocaine, camphor, petrolatum ointment**  
**Wyeth Consumer Healthcare LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Anbesol® Cold Sore Therapy**

***Drug Facts***

<b><i>Active ingredients</i></b>	<b><i>Purposes</i></b>
Allantoin 1%	Skin protectant
Benzocaine 20%	Fever blister/cold sore treatment
Camphor 3%	Fever blister/cold sore treatment
White petrolatum 64.9%	Skin protectant

***Uses***

- temporarily relieves pain associated with fever blisters and cold sores
- relieves dryness and softens fever blisters and cold sores

***Warnings***

**METHEMOGLOBINEMIA WARNING**

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**For external use only**

**Allergy alert**

Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

**Do not use**

- on deep or puncture wounds
- on animal bites
- on serious burns
- for teething
- in children under age 2

**When using this product**

- avoid contact with the eyes

- do not exceed recommended dosage

**Stop use and ask a doctor if**

- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

- to open tube, cut tip of the tube on score mark with scissors
- adults and children 2 years of age and older: apply to the affected area not more than 3 to 4 times daily
- children under 12 years of age: adult supervision should be given in the use of this product
- children under 2 years of age: do not use

***Other information***

store at 20-25°C (68-77°F)

***Inactive ingredients***

aloe barbadensis leaf extract, benzyl alcohol, butylparaben, glyceryl monostearate, isocetyl stearate, menthol, methylparaben, mineral oil, propylparaben, sodium lauryl sulfate, vitamin E, white wax

***Questions or comments?***

Call weekdays 9AM to 5PM EST at **1-888-797-5638**

**PRINCIPAL DISPLAY PANEL - 9 g Tube Label**

Anbesol®

Cold Sore THERAPY

MAXIMUM STRENGTH

Fever Blister/

Cold Sore

Treatment

ointment

NET WT

0.33 OZ

(9 g)



**PRINCIPAL DISPLAY PANEL - 9 g Tube Blister Pack Label**

ointment

WITH

Vitamin E

& Aloe

See new warnings information

Anbesol®

Fever Blister/Cold Sore Treatment

Cold Sore THERAPY

MAXIMUM

STRENGTH

DOCTOR

RECOMMENDED

Instant

Pain Relief

- Cold Sores and  
Fever Blisters
- Moisturizes While It  
Treats and Protects

NET WT 0.33 OZ (9 g)

Safety Sealed Tube:

Do Not Use if tube tip is cut prior to opening.



See new warnings information

ointment  
WITH  
Vitamin E  
& Aloe

# Anbesol<sup>®</sup>

Fever Blister/Cold Sore Treatment

## Cold Sore THERAPY

**MAXIMUM**  
STRENGTH



### Instant Pain Relief



- ✓ Cold Sores and  
Fever Blisters
- ✓ Moisturizes While It  
Treats and Protects

NET WT 0.33 OZ (9 g)

Safety Sealed Tube:  
Do Not Use if tube tip is cut prior to opening.

READ AND KEEP  
CARD FOR  
COMPLETE  
WARNINGS AND  
INFORMATION



AREA FOR LOT,  
EXPIRATION DATE AND  
COUNTRY OF ORIGIN



## Drug Facts

### Active ingredients

Allantoin 1% .....Skin protectant  
Benzocaine 20% .....Fever blister/cold sore treatment  
Camphor 3% .....Fever blister/cold sore treatment  
White petrolatum 64.9% .....Skin protectant

### Purposes

### Uses

- temporarily relieves pain associated with fever blisters and cold sores
- relieves dryness and softens fever blisters and cold sores

### Warnings

**METHEMOGLOBINEMIA WARNING:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

### For external use only

**Allergy alert:** Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

### Do not use

- on deep or puncture wounds
- on animal bites
- on serious burns
- for teething
- in children under age 2

## Drug Facts (continued)

### When using this product

- avoid contact with the eyes
- do not exceed recommended dosage

### Stop use and ask a doctor if

- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### Directions

- to open tube, cut tip of the tube on score mark with scissors
- adults and children 2 years of age and older: apply to the affected area not more than 3 to 4 times daily
- children under 12 years of age: adult supervision should be given in the use of this product
- children under 2 years of age: do not use

### Other information

store at 20-25°C (68-77°F)

**Inactive ingredients** aloe barbadensis leaf extract, benzyl alcohol, butylparaben, glyceryl monostearate, isocetyl stearate, menthol, methylparaben, mineral oil, propylparaben, sodium lauryl sulfate, vitamin E, white wax

**Questions or comments?** Call weekdays 9AM to 5PM EST at 1-888-797-5638

Marketed by: Pfizer, Madison, NJ 07940 USA

© 2018 Pfizer Inc.

For most recent product information, visit [www.anbesol.com](http://www.anbesol.com)



PAA112308

3 0573 0246 25 5

## ANBESOL COLD SORE THERAPY

allantoin, benzocaine, camphor, petrolatum ointment

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0573-0246
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALLANTOIN (UNII: 344S277G0Z) (ALLANTOIN - UNII:344S277G0Z)	ALLANTOIN	10 mg in 1 g

BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g
CAMPHOR (SYNTHETIC) (UNII: 5TJD82A1ET) (CAMPHOR (SYNTHETIC) - UNII:5TJD82A1ET)	CAMPHOR (SYNTHETIC)	30 mg in 1 g
PETROLATUM (UNII: 4T6H12BN9U) (PETROLATUM - UNII:4T6H12BN9U)	PETROLATUM	649 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QP1IU3FV8)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
ISOCETYL STEARATE (UNII: 3RJ7186O9W)	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)	
METHYLPARABEN (UNII: A218C7H9T)	
MINERAL OIL (UNII: T5L8T28FGP)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
WHITE WAX (UNII: 7G1J5DA97F)	

### Product Characteristics

Color	WHITE (White ointment)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0573-0246-25	1 in 1 BLISTER PACK	05/09/2006	06/21/2010
1		9 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0573-0246-26	1 in 1 BLISTER PACK	09/15/2008	
2		9 g in 1 TUBE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	05/09/2006	

**Labeler** - Wyeth Consumer Healthcare LLC (828831730)