

**MAXIMUM STRENGTH ANTACID- aluminum hydroxide, magnesium hydroxide,
dimethicone suspension**

Geri-Care Pharmaceuticals, Corp

gc 664 mylanta vanilla

Active ingredients (in each 10 mL dose)

Aluminum hydroxide 800 mg (equivalent to dried gel, USP)

Magnesium hydroxide 800 mg

Simethicone 80 mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

taking a prescription drug.

Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals as needed, at bedtime,

or as directed by a doctor

- do not take more than 60 mL (6 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor
- measure with dosing cup provided
- mL = milliliter

Other information

- **each 10 mL dose contains:** magnesium 340 mg, sodium 10 mg
- do not freeze
- store at room temperature tightly closed

Inactive ingredients

benzyl alcohol, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose,
purified water, saccharin sodium, sorbitol solution

Questions or comments?

1-800-540-3765

package Label



Compare to the active ingredients in
Mylanta® Maximum Strength*

NDC 57896-664-12

MAXIMUM STRENGTH

Antacid Liquid & Anti-Gas

**ALUMINA, MAGNESIA
& SIMETHICONE ORAL
SUSPENSION, USP**

Fast-Acting

Soothing relief of:

- Acid indigestion
- Sour stomach
- Pressure & bloating
- Heartburn

**Plus a smooth,
creamy taste**



Vanilla Caramel Flavor

12 FL OZ (355 mL)

210-06142-1
Rev 0924

**TAMPER-EVIDENT: DO NOT USE IF
IMPRINTED NECKBAND IS MISSING
OR BROKEN.**

Drug Facts

Active ingredients (in each 10 mL dose)	Purposes
Aluminum hydroxide (equivalent to dried gel, USP) 800 mg.....	Antacid
Magnesium hydroxide 800 mg.....	Antacid
Simethicone 80 mg	Antigas

Uses relieves • heartburn • sour stomach
• acid indigestion • the symptoms referred to as gas

Warnings

Ask a doctor before use if you have
• kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently
taking a prescription drug. Antacids may interact with certain
prescription drugs.

If pregnant or breast-feeding, ask a health professional
before use.
Keep out of reach of children.

Directions

• shake well before use

• adults and children 12 years of age and older: take 10 mL to
20 mL (1 to 2 doses) between meals as needed, at bedtime,
or as directed by a doctor

• do not take more than 60 mL (6 doses) in a 24 hour period
or use the maximum dosage for more than 2 weeks

• children under 12 years of age; ask a doctor

• measure with dosing cup provided

• mL = milliliter

Other information

• each 10 mL dose contains: magnesium 340 mg, sodium 10 mg
• do not freeze • store at room temperature tightly closed

Inactive ingredients benzyl alcohol, carboxymethylcellulose
sodium, flavor, hypromellose, microcrystalline cellulose,
purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

*This product is not manufactured
or distributed by the owner of the
registered trademark Mylanta®
Maximum Strength.

910-06142-0 REV 0524



Distributed by
Geri-Care Pharmaceuticals
1295 Towbin Ave.
Lakewood, NJ 08701

MAXIMUM STRENGTH ANTACID

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-664
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	VANILLA (Vanilla Caramel)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-664-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	05/01/2024	

Labeler - Geri-Care Pharmaceuticals, Corp (611196254)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(57896-664) , pack(57896-664)

Revised: 12/2024

Geri-Care Pharmaceuticals, Corp