

SHISEIDO SPORTS BB LIGHT- octinoxate and titanium dioxide lotion
SHISEIDO SPORTS BB MEDIUM- octinoxate and titanium dioxide lotion
SHISEIDO SPORTS BB DARK- octinoxate and titanium dioxide lotion
SHISEIDO AMERICAS CORPORATION

SHISEIDO SPORTS BB

Drug Facts

<i>Active ingredients</i>	<i>Purpose</i>
OCTINOXATE 4.9%	Sunscreen
TITANIUM DIOXIDE 15.0%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable

Do not use near heat, flame, or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses

- children under 6 months: Ask a doctor

Inactive Ingredients

DIMETHICONE•SILICA•BIS-PEG-18 METHYL ETHER DIMETHYL
SILANE•ISODODECANE•TRIMETHYLSILOXYSILICATE•SD ALCOHOL 40-
B•TRIMETHYLSILOXYSILICATE/DIMETHICONOL CROSSPOLYMER•ISOPROPYL
MYRISTATE•ALUMINUM HYDROXIDE•PEG-9 POLYDIMETHYLSILOXYETHYL
DIMETHICONE•STEARIC ACID•ISOSTEARIC ACID•DEXTRIN
PALMITATE•DISTEARDIMONIUM HECTORITE•BIS-BUTYLDIMETHICONE POLYGLYCERYL-
3•LAUROYL LYSINE•PEG/PPG-14/7 DIMETHYL ETHER•PCA
DIMETHICONE•PHYTOSTERYL/OCTYLDODECYL LAUROYL GLUTAMATE•PHYTOSTERYL
MACADAMIATE•LECITHIN•GLYCYRRHIZA GLABRA (LICORICE) ROOT
EXTRACT•SUCROSE TETRASTEARATE
TRIACETATE•POLYMETHYLSILSESQUIOXANE•TALC•METHYL METHACRYLATE
CROSSPOLYMER•ALUMINA•POLYSILICONE-2•TRIETHOXYCAPRYLYLSILANE•HYDROGEN
DIMETHICONE•TOCOPHEROL•BHT•ETHYLPARABEN•FRAGRANCE•TITANIUM
DIOXIDE•IRON OXIDES•MICA•

Other information

- protect this product in this container from excessive heat and direct sun.
- avoid contact with clothing and other textiles-may stain.

Questions or comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Container Carton - Light

SHISEIDO
GINZA TOKYO

Sports BB

50+
WetForce
Quick Dry

BROAD SPECTRUM
SPF 50+

WATER RESISTANT
(80 MINUTES)

SUNSCREEN

30mL 1 FL. OZ.



Sports BB



BROAD SPECTRUM SPF 50+

SHISEIDO
GINZA TOKYO

Sports BB

50+

WetForce

Quick Dry

BROAD SPECTRUM
SPF 50+

WATER RESISTANT
(80 MINUTES)

SUNSCREEN

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE • SILICA •
BIS-PEG-18 METHYL ETHER DIMETHYL SILANE •
ISODODECANE • TRIMETHYLSILOXYSILICATE •
SD ALCOHOL 40-8 •
TRIMETHYLSILOXYSILICATE/DIMETHICONDOL
CROSSPOLYMER •
ISOPROPYL MYRISTATE • ALUMINUM HYDROXIDE •
PEG-8 POLYDIMETHYLSILOXYETHYL DIMETHICONE •
STEARIC ACID • ISOSTEARIC ACID •
DEXTRIN PALMITATE •
DISTEARDIMONIUM HECTORITE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
LAUROYL LYSINE • PEG/PPG-14/7 DIMETHYL ETHER •
PCA DIMETHICONE •
PHYTOSTERYL OCTYLDODECYL LAUROYL GLUTAMATE •
PHYTOSTERYL MACADAMIATE • LECITHIN •
GLYCYRRHIZA GLABRA (LICORICE) ROOT EXTRACT •
SUCROSE TETRASTEARATE TRIACETATE •
POLYMETHYLSILSESQUOXANE • TALC •
METHYL METHACRYLATE CROSSPOLYMER • ALUMINA •
POLYSILICONE-2 • TRIETHOXYCAPRYLYLSILANE •
HYDROGEN DIMETHICONE • TOCOPHEROL • BHT •
ETHYLPARABEN • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.
- avoid contact with clothing and other textiles-may stain.

54.5x33x82
1 2 3 4 5 6 7 8 9 0



SHISEIDO
SPORTS BB
BROAD SPECTRUM SPF 50+

A BB type sunscreen ideal for outdoor scenes such as sports, featuring WetForce technology so the UV protection veil actually becomes stronger when it comes into contact with sweat and water.* Furthermore, Quick Dry technology dries your sweat quickly** to keep you looking fresh, no matter how active you have been.

- Contains contouring powder to utilize the effects of light and shadow for creating a sculpted, three-dimensional look.
- Reducing the quantity of application will lower the level of sunscreen protection significantly.
- Shake well before use.

NON-COMEDOGENIC. DERMATOLOGIST AND OPHTHALMOLOGIST-TESTED.

*To activate WetForce technology, expose the applied area to water or perspiration for 30 minutes.

**in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 4.9%	Sunscreen
TITANIUM DIOXIDE 15.0%	Sunscreen

Uses

■ helps prevent sunburn ■ if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10022
MADE IN U.S.A.
www.shiseido.com

Drug Facts (continued)

Flammable : Do not use near heat, flame, or while smoking.

Directions

For sunscreen use:



- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
 - children under 6 months: Ask a doctor ▶

<http://s1872.com/14657AA>
See our website for more information.

Drug Facts (continued)

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Container Carton - Medium

SHISEIDO
GINZA TOKYO

Sports BB

50+
WetForce
Quick Dry

BROAD SPECTRUM
SPF 50+

WATER RESISTANT
(80 MINUTES)

SUNSCREEN

30mL 1 FL. OZ.



Sports BB



BROAD SPECTRUM SPF 50+

SHISEIDO
GINZA TOKYO

Sports BB

50+

WetForce
Quick Dry

BROAD SPECTRUM
SPF 50+

WATER RESISTANT
(80 MINUTES)

SUNSCREEN

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE • SILICA •
BIS-PEG-18 METHYL ETHER DIMETHYL SILANE •
ISODODECANE • TRIMETHYLSILOXYSILICATE •
SD ALCOHOL 40-8 •
TRIMETHYLSILOXYSILICATE/DIMETHICONDOL
CROSSPOLYMER •
ISOPROPYL MYRISTATE • ALUMINUM HYDROXIDE •
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE •
STEARIC ACID • ISOSTEARIC ACID •
DEXTRIN PALMITATE •
DISTEARDIMONIUM HECTORITE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
LAUROYL LYSINE • PEG/PPG-14/7 DIMETHYL ETHER •
PCA DIMETHICONE •
PHYTOSTERYL OCTYLDODECYL LAUROYL GLUTAMATE •
PHYTOSTERYL MACADAMIATE • LECITHIN •
GLYCYRRHIZA GLABRA (LICORICE) ROOT EXTRACT •
SUCROSE TETRASTEARATE TRIACETATE •
POLYMETHYLSILSESQUOXANE • TALC •
METHYL METHACRYLATE CROSSPOLYMER • ALUMINA •
POLYSILICONE-2 • TRIETHOXYCAPRYLYLSILANE •
HYDROGEN DIMETHICONE • TOCOPHEROL • BHT •
ETHYLPARABEN • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.
- avoid contact with clothing and other textiles-may stain.

54.5x33x82
1 2 3 4 5 6 7 8 9 0



SHISEIDO
SPORTS BB
BROAD SPECTRUM SPF 50+

A BB type sunscreen ideal for outdoor scenes such as sports, featuring WetForce technology so the UV protection veil actually becomes stronger when it comes into contact with sweat and water.* Furthermore, Quick Dry technology dries your sweat quickly** to keep you looking fresh, no matter how active you have been.

- Contains contouring powder to utilize the effects of light and shadow for creating a sculpted, three-dimensional look.
- Reducing the quantity of application will lower the level of sunscreen protection significantly.
- Shake well before use.

NON-COMEDOGENIC. DERMATOLOGIST AND OPHTHALMOLOGIST-TESTED.

- *To activate WetForce technology, expose the applied area to water or perspiration for 30 minutes.
- **in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 4.9%	Sunscreen
TITANIUM DIOXIDE 15.0%	Sunscreen

Uses

■ helps prevent sunburn ■ if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10022
MADE IN U.S.A.
www.shiseido.com

Drug Facts (continued)

Flammable : Do not use near heat, flame, or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply: · after 80 minutes of swimming or sweating · immediately after towel drying · at least every 2 hours
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including: · limit time in the sun, especially from 10 a.m. – 2 p.m. · wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶



<http://s1872.com/14657AA>

See our website for more information.

Drug Facts (continued)

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Container Carton - Dark

SHISEIDO
GINZA TOKYO

Sports BB

50+
WetForce
Quick Dry

BROAD SPECTRUM
SPF 50+

WATER RESISTANT
(80 MINUTES)

SUNSCREEN

30mL 1 FL. OZ.



Sports BB



BROAD SPECTRUM SPF 50+

SHISEIDO
GINZA TOKYO

Sports BB

50+

WetForce
Quick Dry

BROAD SPECTRUM
SPF 50+

WATER RESISTANT
(80 MINUTES)

SUNSCREEN

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE • SILICA •
BIS-PEG-18 METHYL ETHER DIMETHYL SILANE •
ISODODECANE • TRIMETHYLSILOXYSILICATE •
SD ALCOHOL 40-B •
TRIMETHYLSILOXYSILICATE/DIMETHICONDOL
CROSSPOLYMER •
ISOPROPYL MYRISTATE • ALUMINUM HYDROXIDE •
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE •
STEARIC ACID • ISOSTEARIC ACID •
DEXTRIN PALMITATE •
DISTEARDIMONIUM HECTORITE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
LAUROYL LYSINE • PEG/PPG-14/7 DIMETHYL ETHER •
PCA DIMETHICONE •
PHYTOSTERYL OCTYLDODECYL LAUROYL GLUTAMATE •
PHYTOSTERYL MACADAMIATE • LECITHIN •
GLYCYRRHIZA GLABRA (LICORICE) ROOT EXTRACT •
SUCROSE TETRASTEARATE TRIACETATE •
POLYMETHYLSILSESQUOXANE • TALC •
METHYL METHACRYLATE CROSSPOLYMER • ALUMINA •
POLYSILICONE-2 • TRIETHOXYCAPRYLYLSILANE •
HYDROGEN DIMETHICONE • TOCOPHEROL • BHT •
ETHYL PARABEN • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.
- avoid contact with clothing and other textiles-may stain.

54.5x33x82
 1 2 3 4 5 6 7 8 9 0



SHISEIDO
 SPORTS BB
 BROAD SPECTRUM SPF 50+

A BB type sunscreen ideal for outdoor scenes such as sports, featuring WetForce technology so the UV protection veil actually becomes stronger when it comes into contact with sweat and water.* Furthermore, Quick Dry technology dries your sweat quickly** to keep you looking fresh, no matter how active you have been.

- Contains contouring powder to utilize the effects of light and shadow for creating a sculpted, three-dimensional look.
- Reducing the quantity of application will lower the level of sunscreen protection significantly.
- Shake well before use.

NON-COMEDOGENIC. DERMATOLOGIST AND OPHTHALMOLOGIST-TESTED.

- *To activate WetForce technology, expose the applied area to water or perspiration for 30 minutes.
- **in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 4.9%	Sunscreen
TITANIUM DIOXIDE 15.0%	Sunscreen

Uses

■ helps prevent sunburn ■ if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away. ▶

SHISEIDO AMERICAS CORPORATION DIST.
 NEW YORK, NY 10022
 MADE IN U.S.A.
www.shiseido.com

Drug Facts (continued)

Flammable : Do not use near heat, flame, or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure ■ reapply: · after 80 minutes of swimming or sweating · immediately after towel drying · at least every 2 hours ■ Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including: · limit time in the sun, especially from 10 a.m. - 2 p.m. · wear long-sleeve shirts, pants, hats, and sunglasses ■ children under 6 months: Ask a doctor ▶


<http://s1872.com/14657AA>

See our website for more information.

Drug Facts (continued)

Questions or Comments?

Call toll free 1-800-906-7503



SHISEIDO SPORTS BB LIGHT

octinoxate and titanium dioxide lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-450
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1815 mg in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	5558 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
BIS-PEG-18 METHYL ETHER DIMETHYL SILANE (UNII: OEB4R3VM9C)	
ISODODECANE (UNII: A8289P68Y2)	
TRIMETHYLSILOXYSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)	

STEARIC ACID (UNII: 4ELV7Z65AP)
ISOSTEARIC ACID (UNII: X33R8U0062)
DEXTRIN PALMITATE (CORN; 20000 MW) (UNII: 89B2BSF9I3)
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)
LAUROYL LYSINE (UNII: 113171Q70B)
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)
PHYTOSTERYL/OCTYLDODECYL LAUROYL GLUTAMATE (UNII: 65954KGO9Q)
PHYTOSTERYL MACADAMIATE (UNII: 233VSF903M)
GLYCYRRHIZA GLABRA (UNII: 2788Z9758H)
SUCROSE TETRASTEARATE TRIACETATE (UNII: 1K7LBQ045N)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
TALC (UNII: 7SEV7J4R1U)
ALUMINUM OXIDE (UNII: LMI26O6933)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YOI)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ETHYLPARABEN (UNII: 14255EXE39)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-450-10	1 in 1 CARTON	02/01/2019	
1		30 mL in 1 CONTAINER; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2019	

SHISEIDO SPORTS BB MEDIUM

octinoxate and titanium dioxide lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-451
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1815 mg in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	5558 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
BIS-PEG-18 METHYL ETHER DIMETHYL SILANE (UNII: OEB4R3WW9C)	
ISODODECANE (UNII: A8289P68Y2)	
TRIMETHYLSILOXYSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
ISOSTEARIC ACID (UNII: X33R8U0062)	
DEXTRIN PALMITATE (CORN; 20000 MW) (UNII: 89B2BSF9I3)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
LAUROYL LYSINE (UNII: 113171Q70B)	
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)	
PHYTOSTERYL/OCTYLDODECYL LAUROYL GLUTAMATE (UNII: 65954KGO9Q)	
PHYTOSTERYL MACADAMIATE (UNII: 233VSF903M)	
GLYCYRRHIZA GLABRA (UNII: 2788Z9758H)	
SUCROSE TETRASTEARATE TRIACETATE (UNII: 1K7LBQ045N)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
TALC (UNII: 7SEV7J4R1U)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YOI)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ETHYLPARABEN (UNII: 14255EXE39)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-451-10	1 in 1 CARTON	02/01/2019	
1		30 mL in 1 CONTAINER; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2019	

SHISEIDO SPORTS BB DARK

octinoxate and titanium dioxide lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-452
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1815 mg in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	5558 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
BIS-PEG-18 METHYL ETHER DIMETHYL SILANE (UNII: OEB4R3VM9C)	
ISODODECANE (UNII: A8289P68Y2)	
TRIMETHYLSILOXYSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IU00)	
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
ISOSTEARIC ACID (UNII: X33R8U0062)	
DEXTRIN PALMITATE (CORN; 20000 MW) (UNII: 89B2BSF9I3)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
LAUROYL LYSINE (UNII: 113171Q70B)	
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)	
PHYTOSTERYL/OCTYLDODECYL LAUROYL GLUTAMATE (UNII: 65954KGO9Q)	
PHYTOSTERYL MACADAMIATE (UNII: 233VSF903M)	
GLYCYRRHIZA GLABRA (UNII: 2788Z9758H)	
SUCROSE TETRASTEARATE TRIACETATE (UNII: 1K7LBQ045N)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
TALC (UNII: 7SEV7J4R1U)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YOI)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	

BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ETHYLPARABEN (UNII: 14255EXE39)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-452-10	1 in 1 CARTON	02/01/2019	04/21/2020
1		30 mL in 1 CONTAINER; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2019	04/21/2020

Labeler - SHISEIDO AMERICAS CORPORATION (193691821)

Establishment			
Name	Address	ID/FEI	Business Operations
SHISEIDO AMERICA INC.		782677132	MANUFACTURE(58411-450, 58411-451, 58411-452) , ANALYSIS(58411-450, 58411-451, 58411-452)

Revised: 12/2023

SHISEIDO AMERICAS CORPORATION