

AMAZON CHILDRENS MULTI SYMPTOM COLD- dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution
AMAZON.COM SERVICES LLC

Amazon Children's Multi-Symptom Cold

Drug Facts

<i>Active ingredients (in each 5 mL)</i>	<i>Purposes</i>
Dextromethorphan HBr 5 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Directions

- do not give more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor

Age	Dose
children 6 years to under 12 years of age	10 mL every 4 hours
children 4 years to under 6 years of age	5 mL every 4 hours
children under 4 years of age	do not use

Other information

- **each 5 mL contains:** sodium 2 mg
- store at room temperature
- do not refrigerate
- dosing cup provided

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue No.1, FD&C Red #40, flavors, potassium citrate, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

Distributed by:

PRINCIPAL DISPLAY PANEL

***Compare to the Active Ingredients in Children's Mucinex® Multi-Symptom Cold**

NDC 72288-649-04

Children's

Multi-Symptom Cold

Dextromethorphan HBr 5 mg - Cough Suppressant

Guaifenesin 100 mg - Expectorant

Phenylephrine HCl 2.5 mg - Nasal Decongestant

- Relieves Stuffy Nose
- Controls Cough
- Relieves Chest Congestion
- Breaks up Mucus

Dosing Cup Included

For Ages 4 & Over

Berry Flavor

Naturally & Artificially Flavored

4 FL OZ (118 mL)

TAMPER EVIDENT: DO NOT USE IF NECKBAND ON BOTTLE CAP IS BROKEN OR MISSING.

IMPORTANT: Keep this carton for future reference on full labeling.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Multi-Symptom Cold.

Children's Multi-Symptom Cold

Drug Facts

Active ingredients (in each 5 mL)

Dextromethorphan HBr 5 mg.....Cough suppressant
 Guaifenesin 100 mg.....Expectorant
 Phenylephrine HCl 2.5 mg.....Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Drug Facts (continued)

Directions

- do not give more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor

Age	Dose
children 6 years to under 12 years of age	10 mL every 4 hours
children 4 years to under 6 years of age	5 mL every 4 hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 2 mg
- store at room temperature
- do not refrigerate
- dosing cup provided

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C Blue No. 1, FD&C Red No. 40, flavors, potassium citrate, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum.

Questions or comments? 1-866-467-2748

TAMPER-EVIDENT: DO NOT USE IF PRINTED INNER SEAL UNDER CAP IS TORN OR MISSING.

IMPORTANT: Keep this carton for future reference on full labeling.

Distributed by:

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Multi-Symptom Cold.

*Compare to the Active Ingredients in Children's Mucinex® Multi-Symptom Cold

NDC 72288-649-04

Children's Multi-Symptom Cold

Dextromethorphan HBr 5 mg • Cough Suppressant
 Guaifenesin 100 mg • Expectorant
 Phenylephrine HCl 2.5 mg • Nasal Decongestant

- Relieves Stuffy Nose
- Controls Cough
- Relieves Chest Congestion
- Breaks Up Mucus

Dosing Cup Included

For Ages 4 & Over

Berry Flavor

Naturally and Artificially Flavored
 4 FL OZ (118 mL)

*Compare to the Active Ingredients in Children's Mucinex® Multi-Symptom Cold

Children's Multi-Symptom Cold

Dextromethorphan HBr 5 mg • Cough Suppressant
 Guaifenesin 100 mg • Expectorant
 Phenylephrine HCl 2.5 mg • Nasal Decongestant

- Relieves Stuffy Nose
- Controls Cough
- Relieves Chest Congestion
- Breaks Up Mucus

Dosing Cup Included

For Ages 4 & Over

Berry Flavor

Naturally and Artificially Flavored
 4 FL OZ (118 mL)



Lot:
 NON-VARNISH AREA
 DO NOT PRINT
 Exp:

AMAZON CHILDRENS MULTI SYMPTOM COLD

dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72288-649
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	pink	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-649-04	1 in 1 CARTON	02/16/2026	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/16/2026	

Labeler - AMAZON.COM SERVICES LLC (128990418)

