

**FAMOTIDINE- famotidine tablet**  
**OUR FAMILY (Nash Finch Company)**

-----  
**Drug Facts**

**Active ingredient (in each tablet)**

Famotidine USP 10 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers.

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck, or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to **prevent** symptoms, swallow 1 tablet with a glass of water **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

### **Other information**

- read the directions and warnings before use
- store at 25°-30°C (77°-86°F)
- protect from moisture

### **Inactive ingredients**

hydroxypropyl cellulose, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

### **Questions or comments?**

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

\*Compare to the active ingredient in Pepcid® AC®

#### **Acid Control**

famotidine tablets USP, 10 mg / acid reducer

Just One Tablet

Relieves & Prevents Heartburn Due to Acid Indigestion

ORIGINAL STRENGTH

Tablets

\*This product is not manufactured or distributed by ©McNeil Consumer Pharmaceuticals Co., owner of the registered trademark Pepcid® AC®.

**DISTRIBUTED BY NASH FINCH COMPANY**

7600 FRANCE AVE S, MPLS, MN 55435

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

#### **Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight

- Quit smoking

Product of India

## Product Label

\*Compare to the active ingredient in Pepcid® AC\*



# Acid Control

famotidine tablets USP, 10 mg / acid reducer

Just One Tablet

Relieves & Prevents Heartburn  
Due to Acid Indigestion

ORIGINAL STRENGTH



Actual Size

90 Tablets

\*This product is not manufactured or distributed by ©Moliel Consumer Pharmaceuticals Co., owner of the registered trademark Pepcid® AC\*.

DISTRIBUTED BY NASH FINCH COMPANY  
©2004, 1806 MFC BRANDS  
7500 FRANCE AVE S, MPLS, MN 55435  
www.ourfamilyfoods.com MF18662



If you are not satisfied with any Our Family brand product, simply return it to the store where purchased. We will refund your money PLUS replace the item with a 90¢ item of the brand of your choice - FREE.

**Tip for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



0 70253 96371 1

**Drug Facts**

Active ingredient (in each tablet)	Famotidine USP 10 mg	Purpose	Acid reducer
------------------------------------	----------------------	---------	--------------

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Always alert:** Do not use if you are allergic to famotidine or other acid reducers.

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, swelling, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck, or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Step use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to prevent symptoms, swallow 1 tablet with a glass of water 15 to 60 minutes before eating food or drinking beverages that cause heartburn
  - do not use more than 2 tablets in 24 hours
  - children under 12 years: ask a doctor

**Other information**

- read the directions and warnings before use
- store at 25°-30°C (77°-85°F)
- protect from moisture

**Inactive ingredients:** hydroxypropyl cellulose, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

**Questions or comments?** Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**Our Family Acid Control Tablet**

## FAMOTIDINE

famotidine tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70253-128
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
HYDROXYPROPYL CELLULOSE (TYPE H) (UNII: RFW2ET671P)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	W972
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70253-128-90	1 in 1 BOX		
1		90 in 1 BOTTLE, PLASTIC		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077146	10/15/2013	

**Labeler** - OUR FAMILY (Nash Finch Company) (006962294)

Revised: 10/2013

OUR FAMILY (Nash Finch Company)