

STOOL SOFTENER PLUS STIMULANT LAXATIVE- docusate sodium and sennosides tablet, film coated
Meijer, Inc.

GS 303B (304)

Active ingredient (in each tablet)

Docusate Sodium 50 mg
Sennosides 8.6 mg

Purpose

Stool softener
Stimulant laxative

Uses

- relieves occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

- if you are now taking mineral oil, unless told to do so by a doctor
- laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that lasts over 2 weeks

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- Take only by mouth. Doses may be taken as a single daily dose, preferably in the evening, or in divided doses
-

adults and children 12 years and over	take 2-4 tablets daily
children under 12 years	ask a doctor

Other information

- each tablet contains: **calcium 20 mg, sodium 3 mg**
- store at controlled room temperature
- **Tamper Evident:** Do not use if imprinted seal under cap is missing or broken.
- product of India

Inactive ingredients

cellulose, croscarmellose sodium, dicalcium phosphate, FD&C blue #2 lake, FD&C red #40 lake, hypromellose, magnesium stearate, PEG, silica, talc, titanium dioxide.

Questions or comments?

1-800-540-3765

Package Label

meijer. NDC 79481-0304-1

Stool Softener + Stimulant Laxative

COMPARE TO THE ACTIVE INGREDIENTS OF COLACE® 2-IN-1*

GENTLE, EFFECTIVE, DEPENDABLE CONSTIPATION RELIEF IN 6-12 HOURS

Docosate Sodium 50 mg
Sennosides 8.6 mg

100 TABLETS

Drug Facts

Active ingredients (in each tablet)

Docosate Sodium 50 mg.....Stool softener
Sennosides 8.6 mg.....Stimulant laxative

Uses • relieves occasional constipation (irregularity)
• this product generally produces a bowel movement in 6 to 12 hours

Warnings

Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that lasts over 2 weeks

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • Take only by mouth. Doses may be taken as a single daily dose, preferably in the evening, or in divided doses, adults and children 12 years and over take 2-4 tablets daily children under 12 years ask a doctor

Other information • each tablet contains: calcium 20 mg, sodium 3 mg • store at controlled room temperature

Tamper Evident: Do not use if imprinted seal under cap is missing or broken. • product of India

Inactive ingredients: cellulose, croscarmellose sodium, dicalcium phosphate, FD&C blue #2 lake, FD&C red #40 lake, hypromellose, magnesium stearate, PEG, silica, talc, titanium dioxide.

Questions or comments? 1-800-540-3765

* This product is not manufactured or distributed by Atlantis Consumer Healthcare, Inc., owner of the registered trademark Colace®.

DIST. BY MEIJER DISTRIBUTION, INC. REV 303B-1124
GRAND RAPIDS, MI 49544
www.meijer.com PID 395542

7 13733 50759 2

STOOL SOFTENER PLUS STIMULANT LAXATIVE

docosate sodium and sennosides tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0304
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	8.6 mg

DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)

DOCUSATE SODIUM

50 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CALCIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	PSD21
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0304-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	03/01/2025	

Labeler - Meijer, Inc. (006959555)

Registrant - Geri-Care Pharmaceutical Corp (611196254)

Revised: 3/2025

Meijer, Inc.