

**DAYTIME NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr, doxylamine succinate and phenylephrine hcl liquid filled capsules
Walmart Inc.**

634T_Daytime Nighttime Severe Cold and Flu Softgels

Drug Facts - Daytime Cold & Flu

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan hydrobromide 10 mg

Phenylephrine hydrochloride 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold or flu:

minor aches and pains

headache

cough

sore throat

nasal congestion

sinus congestion and pressure

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

-more than 4,000 mg of acetaminophen in 24 hours

-with other drugs containing acetaminophen

-3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions.

Symptoms may include:

-skin reddening

-blisters

-rash

-hives

-facial swelling

-asthma (wheezing)

-shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough with excessive phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product, do not exceed recommended dosage

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after the last Night dose before starting Day product.
- adults and children 12 years and over: take 2 softgels with water every 4 hours. Do

not exceed 6 softgels in 12 hours or as directed by a doctor.

- children under 12 years: do not use

Other information

store at room temperature

avoid temperature above 25°C (77°F)

Inactive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions or comments?

Call **1-888-287-1915**

Drug Facts - Nighttime Cold & Flu

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan hydrobromide 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine hydrochloride 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold or flu:

minor aches and pains

headache

cough

nasal congestion

runny nose

sneezing

sore throat

sinus congestion and pressure

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions.

Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- may cause marked drowsiness
- avoid alcoholic drinks

- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after the last Day dose before starting Night product.
- adults and children 12 years and over: take 2 softgels with water every 4 hours. Do not exceed 4 softgels in 12 hours or as directed by a doctor.
- children under 12 years: do not use


Other information

- store at room temperature
- avoid temperature above 25°C (77°F)

Inactive ingredients FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions or Comments?

Call **1-877-290-4008**



Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

LET HERE FOR MORE INFORMATION

Drug Facts

Active ingredients (in each softgel) Purposes

Acetaminophen 325 mg. Pain reliever/fever reducer
 Dextromethorphan hydrobromide 10 mg. Cough suppressant
 Phenylephrine hydrochloride 5 mg. Nasal decongestant

Uses

temporarily relieves these symptoms due to a cold or flu:
 minor aches and pains ■ headache
 sore throat ■ sinus congestion and pressure
 cough ■ nasal congestion

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:
 skin redness ■ hives ■ rash ■ skin swelling ■ blisters ■ asthma (wheezing) ■ shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Drug Facts

Active ingredients (in each softgel) Purposes

Acetaminophen 325 mg. Pain reliever/fever reducer
 Dextromethorphan hydrobromide 10 mg. Cough suppressant
 Doxylamine succinate 6.25 mg. Antihistamine
 Phenylephrine hydrochloride 5 mg. Nasal decongestant

Uses

temporarily relieves these symptoms due to a cold or flu:
 minor aches and pains ■ headache
 nasal congestion ■ sneezing
 runny nose ■ sore throat

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:
 skin redness ■ hives ■ rash ■ skin swelling ■ blisters ■ asthma (wheezing) ■ shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

DO NOT TAKE THESE PRODUCTS AT THE SAME TIME

SAFETY SEALED: DO NOT USE IF THE INDIVIDUAL BLISTER UNIT IS OPEN OR TORN

REMAIN CARTON FOR COMPLETE WARNINGS AND INFORMATION

equate™

DAYTIME MULTI-SYMPTOM

Cold & Flu Formula

equate™

NIGHTTIME MULTI-SYMPTOM

Cold & Flu Formula

equate™

DAYTIME MULTI-SYMPTOM

Cold & Flu Formula

Acetaminophen -
 Pain Reliever/Fever Reducer
Dextromethorphan HBr -
 Cough Suppressant
Phenylephrine HCl -
 Nasal Decongestant



Actual Size

Relieves:

- Nasal congestion
- Headache & body ache
- Cough
- Sore throat
- Sinus pressure

12
SOFTGELS

20 TOTAL SOFTGELS

equate™

NIGHTTIME MULTI-SYMPTOM

Cold & Flu Formula

Acetaminophen -
 Pain Reliever/Fever Reducer
Dextromethorphan HBr -
 Cough Suppressant
Doxylamine Succinate -
 Antihistamine
Phenylephrine HCl -
 Nasal Decongestant



Actual Size

Relieves:

- Nasal congestion
- Headache & body ache
- Cough
- Runny nose
- Sore throat

8
SOFTGELS

Compare to
Alka-Seltzer PLUS®
 Day and Night
 Cold and Flu
 active ingredients*



CONTRIBUTED BY: Wal-Mart Inc., Bentonville, AR 72716
PRODUCT OF: MEXICO
 *This product is not manufactured or distributed by Bayer
 HealthCare LLC, owner of the registered trademark
 Alka-Seltzer TABLETS Day and Night Cold and Flu. ©2011 1136

Substitution guaranteed—Or we'll replace it or give you your money back. For questions or comments please call 1-888-888-8888.

LOT:
 EXP.:

PrintWamish Omit Area

PARENTS:
 Let us know what you think


Inactive ingredients: FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions or comments? Call 1-888-287-1915

Inactive ingredients: FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions or comments? Call 1-888-287-1915





DAYTIME NIGHTTIME COLD AND FLU
 acetaminophen, dextromethorphan hbr, doxylamine succinate and phenylephrine hcl liquid filled capsules kit

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79903-315

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-315-20	1 in 1 CARTON	03/01/2025	

Quantity of Parts		
Part #	Package Quantity	Total Product Quantity
Part 1	12 BLISTER PACK	12 in 2
Part 2	8 BLISTER PACK	8 in 2

Part 1 of 2
DAYTIME COLD AND FLU
 acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Product Information

Item Code (Source) NDC:79903-320

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
POVIDONE (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL (Oblong shaped capsules)	Size	16mm
Flavor		Imprint Code	70
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

Part 2 of 2**NIGHTTIME COLD AND FLU**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride liquid filled capsules capsule, liquid filled

Product Information

Item Code (Source) NDC:79903-321

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
WATER (UNII: 059QF0KO0R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POVIDONE (UNII: FZ989GH94E)	
SHELLAC (UNII: 46N107B71O)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL (oblong shaped capsules)	Size	16mm
Flavor		Imprint Code	72
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2025	

Labeler - Walmart Inc. (051957769)

Registrant - TIME CAP LABORATORIES INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LTD		925822975	manufacture(79903-315)

Revised: 4/2025

Walmart Inc.