HEARTBURN RELIEF- famotidine tablet, film coated Meijer Distribution Inc

Meijer Distribution, Inc. Heartburn Relief Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutesbefore eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose (monohydrate), magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Meijer®

Compare to Maximum Strength Pepcid[®] AC active ingredient

MAXIMUM STRENGTH

heartburn relief

Famotidine Tablets, 20 mg


Acid Reducer

SEE NEW WARNINGS

Just One Tablet! Prevents & Relieves Heartburn Due to Acid Indigestion

25 Tablets

actual size



| GODE AREA 194A7 6E C7 | OPEN OTHER END | Drug Facts Active ingredient Purpose (in each tablet) Famotidine 20 mgAcid reducer Uses Active ingredient I relieves heartburn associated with acid indigestion and sour stomach prevents heartburn associated with acid indigestion and sour stomach brought on by eating or d inking certain fcod and beverages Warnings Allergy alert: Do not use i you are allergic to famotidine or otheracid reducers Do not use if you have trouble or pain swallowing food, vorniting with bod, or bbody or black stools. These may be signs of a serbus condition. See your doctor. With theracid reducers Ask adoctor before use if you have had heatburn over 3 months. This may be a sign of a more serious condition. haetburn with lightheadedness, sweating, or dizziness chaet painor shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or should ers; or lightheadedness if requent chest pain frequent wheezing, particularly with heartburn | Drug Facts (continued) Aska doctor or pharmacist before use if you are taking a prescription drug. Acid red ucers may interact with certain prescription drugs. Stop useand ask a doc br if •your heartburn continues or worsens •you needto take this product for more than 14 days If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Centerright away. (1-800-222-1222) Direct ions •adults and children 12 years and over: •to relieve symptoms, swallow 1 tablet with aglass of water. Do not chew. •to relieve symptoms, swallow 1 tablet with aglass of water at any time from 10to 60 minutes before eating foodor drinking beverages that cause heartburn •dnidren under 12 years: aska doctor Other information •cad the directions and warnings before use •keep the carton. It contains important information. •store at 20° -25°C (68° -77°F) •protect from moisture | The for Managing Heartburn • Do not leaf Hator bend over after eaching • Do not wear tight-fitting dothing around the sbmach • Do not each before bedtime • Raise the head of your bed • Aroid hearthum-causing foods such as rich, spior, fathy or friedfoods, chocolate, cafferine, alcohol, and cartain fruits and vegetables • Eat shwly and avoid big meals • If overweight, lose weight • Outh smoking CONVENTENT RECLOSING TAB |
|---------------------------------|----------------|---|--|--|
| | ϕ | unexplained weight bss a nausea or vom ting stomachpain ■ kidney disease Drug Facts (continued) Questions or comments?1-800-719-9260 *Thisproduct is not manufac bired or distributed by Johnson & Johnson, owner of the regis bred tademark Pepcid [®] . | monohy drate, magnesium steatate, microcry stalline cellulos e, polyethy lene glycol, polyvin yl alcohol, talc, titanium dioxide PID 467404 7 19283 65644 5 | Tips: • bo: bo: such veget |

| HEARTBURN RELIEF famotidine tablet, film coated | | | | | |
|---|--|--------------------|-------|---------------|--|
| Product Information | | | | | |
| Product Type | HUMAN OTC DRUG | Item Code (Source) | | NDC:41250-712 | |
| Route of Administration | ORAL | | | | |
| | | | | | |
| Active Ingredient/Active Moiety | | | | | |
| Ingredient Name Basis of Streng | | | ength | Strength | |
| FAMOTIDINE (UNII: 5QZ015J2Z8) | 5J2Z8) | FAMOTIDINE | | 20 mg | |
| | | | | | |
| Inactive Ingredients | | | | | |
| Ingredient Name | | | 9 | Strength | |
| CARNAUBA WAX (UNII: R12CBM0E | IZ) | | | | |
| SILICON DIOXIDE (UNII: ETJ7Z6XB | U4) | | | | |
| CROSCARMELLOSE SODIUM (UN | CROSCARMELLOSE SODIUM (UNII: M280L1HH48) | | | | |

| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X) | |
|---|--|
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| | |

Product Characteristics

| Color | WHITE | Score | no score |
|----------|-------|--------------|----------|
| Shape | ROUND | Size | 8mm |
| Flavor | | Imprint Code | L194 |
| Contains | | | |

Packaging

| # | ltem Code | Package Description | Marketing Start Date | Marketing End Date |
|---|----------------------|--|-------------------------|-----------------------|
| 1 | NDC:41250-712- 02 | 25 in 1 CARTON | 04/17/2019 | |
| 1 | | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |
| 2 | NDC:41250-712- 71 | 1 in 1 CARTON | 04/17/2019 | |
| 2 | | 50 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 3 | NDC:41250-712- 78 | 2 in 1 CARTON | 08/11/2022 | |
| 3 | | 50 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| | | | | |
| Μ | larketing | Information | | |
| | | Marketing Application Number or Monograph Category Citation | | Marketing End |
| | Category | | Marketing Start Date | Date |

Labeler - Meijer Distribution Inc (006959555)

Revised: 8/2022

Meijer Distribution Inc