

**XT 22C CONC. PART B- sodium chlorite solution
AgroChem USA, LLC.**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

XT-22C CONCENTRATE | PART B

XT-22C CONCENTRATE | PART B

CONCENTRATED ClO₂ TEAT DIP PART B

DANGER

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE.

AVOID FREEZING.

KEEP OUT OF DIRECT SUNLIGHT.

CAUSES SERIOUS EYE IRRITATION.

FOR FARM AND INDUSTRIAL USE ONLY.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/
face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

NDC: 17307-3011-3

NET CONTENTS: 55 GALLON (208.2L)

LOT:

EXP. DATE:

ACTIVE INGREDIENT: Sodium Chlorite 3.5%

XT- 22C Concentrate is a two-part chlorine dioxide teat dip that can be effectively used to reduce the spread of mastitis causing organisms. Recommended for post milking applications.

DIRECTIONS:

IMPORTANT- THIS IS A TWO-PART CONCENTRATE - MUST BE MIXED PRIOR TO USE. MIX RATIO is (20 Water: 1 Part A: 1 Part B). In a clean container, add (20) parts water, then add (1) part XT- 22C Concentrate Part A, next add (1) part XT- 22C Concentrate Part B. FOLLOW THE DIRECTIONS. DO NOT MIX PART A AND B TOGETHER. CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels (> 1000 ppm) within 15 minutes of mixing.

General Use

Once mixed according to directions, do not mix product with any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

This product can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forest ripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels.

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

Manufactured By:

AgroChem USA, LLC 26 Freedom Way • Saratoga Springs, NY 12866 • (518)226-4850

XT-22C CONCENTRATE | PART B

**CONCENTRATED
ClO₂ TEAT DIP
PART B**

DANGER



**NOT FOR HUMAN USE.
AVOID FREEZING.
CAUSES EYE IRRITATION.
KEEP OUT OF DIRECT SUNLIGHT.
KEEP OUT OF REACH OF CHILDREN.
FOR FARM AND INDUSTRIAL USE ONLY.**

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID/ RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant.

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

ACTIVE INGREDIENT: Sodium Chlorite 3.5%

NDC: 17307-3011-3

NET CONTENTS: 55 GALLON (208.2L)

LOT:

EXP. DATE:



UN 1908 Chlorite solution, 8, III

XT-22C Concentrate is a two-part chlorine dioxide teat dip that can be effectively used to reduce the spread of mastitis causing organisms. Recommended for post milking applications.

DIRECTIONS:

IMPORTANT- THIS IS A TWO-PART CONCENTRATE - MUST BE MIXED PRIOR TO USE. MIX RATIO is (20 Water : 1 Part A : 1 Part B). In a clean container, add (20) parts water, then add (1) part XT-22C Concentrate Part A, next add (1) part XT-22C Concentrate Part B. FOLLOW THE DIRECTIONS. DO NOT MIX PART A AND B TOGETHER. CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels (> 1000ppm) within 15 minutes of mixing.

General Use

If using a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

This product can also be foamed or sprayed using recommended equipment

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior to milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels.

Just before each cow is milked, dip, foam, or spray, the full length of each teat with the product. Allow product to stand for 30 seconds. Wipe teats dry after application, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip, foam, or spray, the full length of each teat.

Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the product should be applied to the teats, once a day for 3- 4 days after the last milking. Applications should resume at least twice daily one week before calving.

Manufactured By: AgroChem USA, LLC
26 Freedom Way
Saratoga Springs, NY 12866
Phone: 518-226-4850



XT 22C CONC. PART B

sodium chlorite solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:17307-3011
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORITE (UNII: G538EBV4VF) (CHLORITE ION - UNII:Z63H374SB6)	SODIUM CHLORITE	0.114 kg in 1 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-3011-1	19 kg in 1 CONTAINER		
2	NDC:17307-3011-2	57 kg in 1 DRUM		
3	NDC:17307-3011-3	208 kg in 1 DRUM		
4	NDC:17307-3011-4	1003 kg in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/25/2024	

Labeler - AgroChem USA, LLC. (118336116)

Registrant - AgroChem USA, LLC. (118336116)

Establishment

Name	Address	ID/FEI	Business Operations
AgroChem USA, LLC.		118336116	api manufacture, manufacture, analysis, label, pack

Revised: 1/2024

AgroChem USA, LLC.