

# **ASPIRIN REGULAR STRENGTH- aspirin 325mg tablet**

## **Ulai Health LLC**

-----

### **Drug Facts**

#### **Active Ingredient (in each tablet)**

Aspirin 325mg (NSAID) †

†nonsteroidal anti-inflammatory drug

#### **Purpose**

Pain reliever/fever reducer

#### **Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscle pain
- toothache
- menstrual pain
- colds
- minor pains of arthritis
- temporarily reduces fever

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

## **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

## **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

## **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

## **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur

## **If pregnant or breast-feeding,**

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## **Directions**

- drink a full glass of water with each dose

---

|                                       |  |
|---------------------------------------|--|
| adults and children 12 years and over | take 1 to 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours |
| children under 12 years               | ask a doctor   |

---

## Other information

- store at 20-25°C (68-77°F)
- avoid excess heat and moisture

## Inactive ingredient

corn starch

## Questions or comments?

(866) 562-2756 (Mon- Fri 8 AM to 4 PM EST)

**PHARBEST**

**NDC 73057-365-11**

**Manufactured in the USA**

**\*Compare to the active ingredient in Genuine Bayer® Aspirin**

**Aspirin**

**Regular Strength**

**Pain Reliever/Fever Reducer (NSAID)**

**1000 TABLETS**

**Aspirin, 325 mg each**

**PHARBEST** **Manufactured in the USA**  
NDC 73057-365-11

*"Compare to the active ingredient in Genuine Bayer® Aspirin"*

**Aspirin**  
*Regular Strength*  
**Pain Reliever/Fever Reducer (NSAID)**

**1000 TABLETS**  
Aspirin, 325 mg each

**THIS PACKAGE FOR HOUSEHOLD WITHOUT YOUNG CHILDREN**  
**TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**  
**Active ingredient (in each tablet)**  
Aspirin 325 mg (NSAID)..... Pain reliever/fever reducer  
nonsteroidal anti-inflammatory drug

**Uses**  
Temporarily relieves minor aches and pains due to:  
■ headache ■ muscle pain ■ toothache ■ menstrual pain ■ colds ■ minor pain of arthritis ■ temporarily reduces fever

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor immediately. Symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription drugs (aspirin, ibuprofen, naproxen, etc.) ■ are 3 or more times taking this drug every day using this drug ■ also take or use for a long time other drugs that may increase the risk of bleeding such as aspirin, ibuprofen, naproxen, etc.

**Do not take if:**  
■ you have an allergic reaction to this product or any of its ingredients ■ if you have ever had an allergic reaction to this product or any of its ingredients ■ if you are taking aspirin or another NSAID

**Ask doctor before use if:**  
■ you are taking aspirin or another NSAID ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for:**  
■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if:**  
■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not go away ■ have dark or bloody urine ■ have black or tarry stools ■ have nose or rectal bleeding ■ dizziness ■ weakness ■ swelling in the legs ■ ringing in the ears or a loss of hearing occurs ■ new symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**  
■ drink a full glass of water with each dose

|                                       |  |
|---------------------------------------|--|
| adults and children 12 years and over | take 1 to 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours |
| children under 12 years               | ask a doctor   |

**Other information**  
■ store at 20-25°C (68-77°F) ■ avoid excess heat and moisture

**Inactive ingredient** corn starch

**Questions or comments?** (866) 562-2756 (Mon-Fri 8 AM to 4 PM EST)

This product has been manufactured and distributed by Bayer HealthCare LLC, owner or registered trademark of Genuine Bayer® Aspirin.

**Distributed by:**  
**ULAI HEALTH LLC,**  
Alexandria, VA 22312

Org. 06/19 9  
7 3 0 5 7 | 3 6 5 1 1

## ASPIRIN REGULAR STRENGTH

aspirin 325mg tablet

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:73057-365 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name  |  | Basis of Strength  | Strength             |                    |
|--|--|--|----------------------|--------------------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) |  | ASPIRIN  | 325 mg               |                    |
| <b>Inactive Ingredients</b>                            |  |  |                      |                    |
| Ingredient Name  |  | Strength   |                      |                    |
| STARCH, CORN (UNII: O8232NY3SJ)                        |  |  |                      |                    |
| <b>Product Characteristics</b>                         |  |  |                      |                    |
| Color  | white                                    | Score  | no score             |                    |
| Shape  | ROUND (ROUND TABLET)                     | Size   | 10mm                 |                    |
| Flavor   |  | Imprint Code   | PH033                |                    |
| Contains   |  |  |                      |                    |
| <b>Packaging</b>                                       |  |  |                      |                    |
| #  | Item Code                                | Package Description  | Marketing Start Date | Marketing End Date |
| 1  | NDC:73057-365-11                         | 1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 06/28/2019           |                    |
| 2  | NDC:73057-365-08                         | 100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product  | 06/28/2019           |                    |
| <b>Marketing Information</b>                           |  |  |                      |                    |
| Marketing Category                                     | Application Number or Monograph Citation | Marketing Start Date   | Marketing End Date   |                    |
| OTC Monograph Drug                                     | M013                                     | 06/28/2019   |                      |                    |

**Labeler** - Ulai Health LLC (081181535)

Revised: 1/2025

Ulai Health LLC