

MUCOSAN MAX WITH ACETAMINOPHEN- acetaminophen, dextromethorphan hydrobromide, guaifenesin and phenylephrine hydrochloride liquid
Menper Distributors, Inc.

Mucosan Max with Acetaminophen

Active Ingredients (in each 20 mL) Purpose

Acetaminophen, 650 mg.....Pain reliever/Fever reducer

Dextromethorphan HBr, 20 mg.....Cough suppressant

Guaifenesin, 400 mg.....Expectorant

Phenylephrine HCl, 10 mg.....Nasal Decongestant

Pain reliever/Fever reducer

Cough suppressant

Expectorant

Nasal Decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
- nasal congestion
- cough
- minor aches and pains
- sore throat
- headache
- temporarily reduces fever
- helps loosen phlem (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: this product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4 doses (20 mL each) in 24 hours, wich is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters and rash. If skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- With any other drug containing acetamonophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

- if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or Pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)
- trouble urination due to enlarged prostate gland.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not use more than direct.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough comes back or occurs with fever, rash, or persistent headache
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur. These could be signs of a serious condition.

If pregnant or breast - feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **do not take more than direct**
- do not take more than 4 doses in any 24 hours
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- dose as follows

age	dose
-----	------

adults and children 12 years of age and older	20 mL every 4 hours
children under 12 years of age	do not use

Other information

- **each (20 mL) contains:** sodium 10mg
- store between 15-30°C (59-86°F)
- do not refrigerate.

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum.

Questions or comments?

1-800-560-5223 Monday to Friday 9am-4pm, Eastern Time





Drug Facts (continued)

Directions

- Do not take more than directed
- Do not take more than 4 doses in any 24 hours
- This adult strength product is not intended for use in children under 12 years of age
- Reseal only with the cap provided
- Keep out of reach of children
- Keep this and all other medicines out of the reach of children
- Use as directed
- Do not use for 14 days

Age	Dose
Adults and children 12 years of age and older	20 mL every 4 hours
Children under 12 years of age	Do not use

Other information

- Each 20 mL contains acetaminophen 10 mg
- Store between 15-30°C (59-86°F) Do not refrigerate.

Inactive ingredients

benzoylcholine chloride, croscarmellose sodium, FD&C blue #1, FD&C red #40, flavors, glycerin, hydroxypropyl methylcellulose, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum.

Questions or comments? 1-800-800-8223 Monday to Friday 9am-4pm, Eastern Time

Información del Medicamento

Ingredientes activos (en cada 20 mL)

Acetaminofén, 100 mg. Anilísico/Paralolol
Dextrometorfano HBr, 20 mg. Fenilefrina HCl, 10 mg.

Usos

- Alivia temporalmente los síntomas de la gripe
- comparación nasal
- tos
- coqueo y molestias menores
- dolor de garganta
- dolor de cabeza
- reduce temporalmente la fiebre
- ayuda a disminuir la fiebre (mucho) y a reducir las secreciones bronquiales para disminuir las molestias respiratorias y generar las más productivas.

Advertencias

Advertencia acerca del líquido: este producto contiene acetaminofén. Puede producirle maleficio hepático grave si toma más de 4 dosis (de 20 mL cada una) en 24 horas. Si está en la cantidad diaria máxima para este producto con otros fármacos que contienen acetaminofén, 3 o más bebidas alcohólicas distintas mientras usa este producto.

Advertencia acerca del dolor de garganta: si el dolor de garganta es intenso, persiste por más de 2 días, está acompañado a menudo de fiebre, dolor de cabeza, nariz roja, inflamación o dificultad, consulte a un médico de inmediato.

No utilice si con ningún otro fármaco que contenga acetaminofén (incluido el de venta libre). Si no se está seguro si el fármaco contiene acetaminofén, pregunte a un médico o farmacéutico. Si está tomando inhibidores de la monoaminoxidasa (MAOI) recetados como fármacos para la depresión, esclerosis múltiple o esclerosis, o la enfermedad de Parkinson o durante 2 semanas después de interrumpir el fármaco MAOI, si no está si el fármaco recetado contiene un MAOI, pregunte a un médico o farmacéutico antes de tomar este producto.

Información del Medicamento (Continuación)

Consulte a su médico antes de usar si padece:

- enfermedad hepática
- enfermedad cardíaca
- hipertensión
- enfermedad tiroidea
- diabetes
- problemas renales
- problemas de la vejiga
- problemas de la próstata
- problemas de la glándula tiroides
- problemas de la glándula paratiroides
- problemas de la glándula pituitaria
- problemas de la glándula hipófisis
- problemas de la glándula hipófisis anterior
- problemas de la glándula hipófisis posterior
- problemas de la glándula hipófisis superior
- problemas de la glándula hipófisis inferior
- problemas de la glándula hipófisis media
- problemas de la glándula hipófisis lateral
- problemas de la glándula hipófisis anterior superior
- problemas de la glándula hipófisis anterior inferior
- problemas de la glándula hipófisis posterior superior
- problemas de la glándula hipófisis posterior inferior
- problemas de la glándula hipófisis media superior
- problemas de la glándula hipófisis media inferior
- problemas de la glándula hipófisis lateral superior
- problemas de la glándula hipófisis lateral inferior

Consulte a su médico o farmacéutico antes de usarlo si está tomando el fármaco antidiabético insulina.

Cuando use este producto, no use más de lo indicado.

Suspenda su uso y consulte a su médico si:

- tiene náuseas, vómitos o diarrea
- los síntomas no mejoran en los siguientes 7 días o están acompañados de fiebre
- los ojos se ponen rojos o se ponen amarillos, o dolor de la cabeza persiste
- la fiebre empeora o dura más de 3 días
- hay erupción o hinchazón
- se presentan nuevos síntomas.
- Estos pueden ser signos de un efecto grave.

Si está embarazada o lactando, pregunte a un profesional de la salud antes de usar.

Mantenga fuera del alcance de los niños.

Advertencia de sobredosis: tomar más de la dosis recomendada (sobredosis) puede provocar daño hepático. En caso de sobredosis, busque ayuda médica y comuníquese con un centro de toxicología de inmediato. La atención médica rápida es fundamental aunque no se observen signos ni síntomas.

Indicaciones

- No tomar más de lo indicado
- Los tomar más de 4 dosis en un periodo de 24 horas
- este producto de concentración para reducir su estrés
- Estadillo para usarse en niños mayores de 12 años
- medir más con la copa dosificadora incluida
- Mantenga la tapa dosificadora con este producto
- 1 mL = 20 gotas
- Administrador de dosis según se indica a continuación

edad	dosis
niños y niñas de 12 años de edad y mayores	20 mL cada 4 horas
niños menores de 12 años	no usar

Otra información

- Each 20 mL contains 10 mg
- Store between 15-30°C (59-86°F)
- Do not refrigerate.

Ingredientes inactivos

ácido cítrico, amilo, amilo, FD&C azul #1, FD&C rojo #40, saborizante, propilglicol, agua purificada, benzoato de sodio, sorbitol, sucralosa, goma xantana.

Preguntas o comentarios? 1-800-800-8223 Lunes a Viernes 9am-4pm, Hora del Este

Distributed by / Distribuido por: The Menzer Group 11425 NW 107th St, Miami, FL 33178, USA

MADE IN USA HECHO EN EU

Drug Facts

Active Ingredients

Acetaminophen, 650 mg... Pain reliever/fever reducer
Dextromethorphan HBr, 20 mg... Cough suppressant
Guafenesin, 400 mg... Decongestant
Phenylephrine HCl, 10 mg... Nasal decongestant

Purposes

- temporarily relieve these common cold and flu symptoms
- nasal congestion
- cough
- minor aches and pains
- sore throat
- headache
- temporarily reduce fever
- helps loosen phlegm (mucus) and thin bronchial secretions to help the bronchial passages of bothersome mucus and make coughs more productive.

Uses

- temporarily relieve these common cold and flu symptoms
- nasal congestion
- cough
- minor aches and pains
- sore throat
- headache
- temporarily reduce fever
- helps loosen phlegm (mucus) and thin bronchial secretions to help the bronchial passages of bothersome mucus and make coughs more productive.

Warnings

Liver warning: this product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses (20 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Alert: acetaminophen may cause severe skin reactions. Symptoms may include skin redness, blisters and rash. If skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are taking a prescription medicine and/or other (OTC) over-the-counter drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drugs, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease
- kidney disease
- high blood pressure
- thyroid disease
- diabetes
- breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to enlarged prostate gland.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not use more than directed.

Stop use and ask a doctor if:

- you get nervous, dizzy or drowsy
- symptoms do not get better within 7 days or are accompanied by fever
- cough turns black or occurs with fever, rash, or persistent headache
- fever gets worse or lasts more than 3 days
- nausea or vomiting is present
- new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: taking more than the recommended dose (intended) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.



MUCOSAN MAX WITH ACETAMINOPHEN

acetaminophen, dextromethorphan hydrobromide, guaifenesin and phenylephrine hydrochloride liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53145-696
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL	

PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL
---	------------------------------------	-------------------

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
XANTHAN GUM (UNII: TTV12P4NEE)	
WATER (UNII: 059QF0KOOR)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53145-696-06	1 in 1 CARTON	11/01/2024	
1		177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/01/2024	

Labeler - Menper Distributors, Inc. (101947166)

Registrant - Menper Distributors, Inc. (101947166)

Revised: 11/2024

Menper Distributors, Inc.