

# **MEIJER BISMUTH STOMACH RELIEF- bismuth tablet, chewable**

## **Meijer**

-----

### **Bismuth 262 mg**

### **Active ingredient (in each chewable tablet)**

Bismuth subsalicylate 262 mg

### **Purpose**

Upset stomach reliever and antidiarrheal

### **Uses** relieves

- Traveler's diarrhea
- Diarrhea
- Upset stomach due to overindulgence in food and drink, including
- Heartburn
- Indigestion
- Nausea
- Gas
- Belching
- Fullness

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

### **Allergy alert: Contains salicylate. Do not take if you are**

- Allergic to salicylates (including aspirin)
- taking other salicylates products

### **Do not use if you have**

- An ulcer
- A bleeding problem
- Bloody or black stool

### **Ask a doctor before use if you have**

- Fever
- Mucus in the stool

### **Ask a doctor or pharmacist before use if you are** taking any drug for

- Anticoagulation (thinning the blood)
- Diabetes
- Gout
- Arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- Symptoms get worse or last more than 2 days
- Ringing in the ears or loss of hearing occurs
- Diarrhea lasts more than 2 days

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- Fully chew before swallowing
- Adults and children 12 years and over:
  - 2 chewable tablets every 1/2 hour or 4 chewable tablets every hour as needed for diarrhea.
  - 2 chewable tablets every 1/2 hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea).
- Do not exceed 8 doses (16 chewable tablets) in 24 hours.
- Use until diarrhea stops but not more than 2 days
- Children under 12 years: ask a doctor.
- Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.

**Inactive ingredients:** Beeswax, calcium carbonate, carmine, carnauba wax, citric acid, corn starch, corn syrup, dextrin, ethyl acetate, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, glycerin, gum arabic, hydrogenated coconut oil, maltodextrin, medium chain triglycerides, natural and artificial flavors, phosphoric acid, propylene glycol, shellac, sodium benzoate, sorbic acid, sorbitol, soy lecithin, soybean oil, sucrose, titanium dioxide, vegetable oil (coconut and/or canola oil), water.

**Other information**

- **Each chewable tablet contains:** Calcium 50mg
- Salicylate 123 mg
- Contains soy
- Store between 20° to 25°C (68° to 77°F) in a dry place. Avoid excessive heat (over 40°C).

**Questions?** Call **1-800-245-2898** Mon - Fri, 9AM - 4PM EST

TAMPER EVIDENT: DO NOT USE  
IF INNER PRINTED SEAL ON  
BOTTLE IS BROKEN OR MISSING.

PN 60002510



PID 5451199  
DIST. BY MEIJER  
DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com



**meijer.**  
**Stomach  
Relief  
Superchews®**

Bismuth Subsalicylate 262 mg  
Upset Stomach Reliever/Antidiarrheal

**5 SYMPTOM RELIEF OF:** UPSET STOMACH, INDIGESTION, NAUSEA, HEARTBURN, DIARRHEA  
**MINT FLAVOR**

NDC-00000-00000-0

COMPARE TO  
PEPTO BISMUL®  
ACTIVE INGREDIENT\*

Chewy,  
Not Chalky®

**24**  
CHEWABLE  
TABLETS

**Drug Facts**

**Active ingredient (in each chewable tablet)** Purpose  
Bismuth subsalicylate 262 mg.....Upset stomach  
reliever and antidiarrheal

**Uses** relieves ■ Traveler's diarrhea ■ Diarrhea  
■ Upset stomach due to overindulgence in food and drink,  
including: ■ Heartburn ■ Indigestion  
■ Nausea ■ Gas ■ Belching ■ Fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or  
are recovering from chicken pox or flu-like symptoms  
should not use this product. When using this product, if  
changes in behavior with nausea and vomiting occur,  
consult a doctor because these symptoms could be an  
early sign of Reye's syndrome, a rare but serious illness. ▶

PEEL BACK FOR DRUG FACTS ▶

**Drug Facts (continued)**

**Allergy alert:** Contains salicylate. Do not take if you are  
■ Allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use if you have**  
■ An ulcer ■ A bleeding problem ■ Bloody or black stool

**Ask a doctor before use if you have** ■ Fever ■ Mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for**  
■ Anticoagulation (thinning the blood) ■ Diabetes ■ Gout ■ Arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or  
tongue may occur

**Stop use and ask a doctor if** ■ Symptoms get worse or last more than 2 days  
■ Ringing in the ears or loss of hearing occurs ■ Diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach  
of children.** In case of overdose, get medical help or contact a Poison Control Center  
right away. ▶

**Drug Facts (continued)**

**Directions**  
■ Fully chew before swallowing  
■ Adults and children 12 year and over:  
■ 2 chewable tablets every ½ hour or 4 chewable tablets every hour as needed  
for diarrhea. ■ 2 chewable tablets every ½ hour to 1 hour as needed for  
overindulgence (upset stomach, heartburn, indigestion, nausea).  
■ Do not exceed 8 doses (16 chewable tablets) in 24 hours.  
■ Use until diarrhea stops but not more than 2 days  
■ Children under 12 years: ask a doctor.  
■ Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.

**Inactive ingredients** Beeswax, calcium carbonate, carmine, carnauba wax,  
citric acid, corn starch, corn syrup, dextrin, ethyl acetate, FD&C blue no. 1  
aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake,  
FD&C yellow no. 6 aluminum lake, glycerin, gum arabic, hydrogenated coconut  
oil, maltodextrin, medium chain triglycerides, natural and artificial flavors,  
phosphoric acid, propylene glycol, shellac, sodium benzoate, sorbic acid, sorbitol,

**Drug Facts (continued)**

soy lecithin, soybean oil, sucrose, titanium dioxide, vegetable oil (coconut and/or  
canola oil), water.

**Other information**

■ Each chewable tablet contains: Calcium 50 mg ■ Salicylate 123 mg ■ Contains soy  
■ Store between 20° to 25°C (68° to 77°F) in a dry place. Avoid excessive heat (over 40°C).

**Questions?** Call 1-800-245-2898 Mon - Fri, 9AM - 4PM EST

\*This product is not manufactured or distributed by the owner of the registered trademark Pepto Bismol®.

Empty Before  
Recycling



how2recycle.info

**MEIJER BISMUTH STOMACH RELIEF**

bismuth tablet, chewable

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79481-0097
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE)	<b>BISMUTH SUBSALICYLATE</b>	262 mg
--	------------------------------	--------

### Inactive Ingredients

Ingredient Name	Strength
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	

### Product Characteristics

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	14mm
<b>Flavor</b>	MINT	<b>Imprint Code</b>	SR
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0097-1	24 in 1 BOTTLE; Type 0: Not a Combination Product	11/11/2024	



## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	11/11/2024	

**Labeler** - Meijer (006959555)

**Registrant** - Bestco (002149136)

## Establishment

Name	Address	ID/FEI	Business Operations
Bestco		002149136	manufacture(79481-0097)

