

MUCOSAN SEVERE MULTI-SYMP TOM- dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid
Menper Distributors, Inc.

Mucosan Severe Multi-Symptom

Active ingredients (in each 20 mL) Purpose

Dextromethorphan HBr, 20 mg.....Cough suppressant

Guaifenesin, 400 mg.....Expectorant

Phenylephrine HCl, 10 mg.....Nasal decongestant

Purposes

Cough suppressant

Expectorant

Nasal descong estant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation.

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional condition, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

Do not exceed recommended dosage.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a poison Control Center right away.

Directions

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- mL=milliliter

age	dose
adults and children 12 years of age and older	20 mL every 4 hours
children under 1 years of age	do not use

Other information

- **each (20mL) dose contains:** sodium 10 mg
- store between 15-30°C (59-86°F)
- do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum.

Questions or comments?

1-800-560-5223 Monday to Friday 9am-4pm, Eastern Time

Muco san

max-action

relief · alivio

Dextromethorphan HBr/
Guaifenesin/
Phenylephrine HCl //
Dextrometorfano HBr/
Guaifenesina/
Fenilefrina HCl



**COUGH &
SEVERE CONGESTION**

CONGESTIÓN Y TOS SEVERA

RELIEVES / ALIVIA:

Cough/Tos

Nasal Congestion/
Congestión Nasal

Chest Congestion/
Congestión de Pecho

12+
years/
años

100-1

6 FL oz (177 mL)

Muco san


max-action

DO NOT USE IF PRINTED SEAL UNDER CAP IS
TORN OR MISSING / NO SE USE SI EL SELLO
IMPRESO DEBAJO DE LA TAPA ESTA ROTO O FALTA

Muco san

max-action

Dextromethorphan HBr/Guaifenesin/Phenylephrine HCl //
Dextrometorfano HBr/ Guaifenesina/Fenilefrina HCl



COUGH & SEVERE CONGESTION

CONGESTIÓN Y TOS SEVERA

RELIEVES / ALIVIA:
Cough/Tos
Nasal Congestion/
Congestión Nasal
Chest Congestion/
Congestión de Pecho

12+
years/
años

6 FL oz (177 mL)

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter

age	dose
adults and children 12 years of age and older	20 mL every 4 hours
children under 12 years of age	do not use

Other information

- each (20 mL) dose contains: sodium 10 mg
- store between 15-30°C (59-86°F) do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, propyl galate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum.

Questions or comments?

1-800-680-5223 Monday to Friday 9am-4pm,
Eastern Time

Información del Medicamento

Ingredientes activos (en cada 20 mL)	Propósitos
Dextrometorfano HBr, 20 mg	Inhibidor de la tos
Guaifenesina, 400 mg	Expectorante
Fenilefrina HCl, 10 mg	Descongestionante nasal

Usos

- ayuda a desprender las flemas (mucosidad) y a adelgazar las secreciones bronquiales para drenar las vías respiratorias
- alivia temporalmente los siguientes síntomas que se presentan con el resfriado:
 - congestión nasal
 - tos debida a irritación leve de garganta y bronquios.

Advertencias

No utilice si usted está tomando un inhibidor de la monoamino oxidasa (IMAO) de prescripción (ciertos medicamentos para la depresión, condiciones psicológicas o emocionales, o enfermedad de Parkinson), o durante 2 semanas después de suspender el medicamento IMAO. Si usted desconoce si su medicamento de prescripción contiene un IMAO, consulte a su médico o farmacéutico antes de tomar este producto.

Consulte a su médico antes de usar, si padece

- enfermedad cardíaca
- hipertensión
- enfermedad de la tiroides
- diabetes
- dificultad al orinar debido a distensión de la glándula prostática
- tos que se presenta con exceso de flema (mucosidad)
- tos persistente o crónica tal como ocurre con el tabaquismo, asma, bronquitis crónica o enfisema.

Información del Medicamento (continuación)

Consulte a su médico o farmacéutico antes de utilizarlo si usted está tomando cualquier otro descongestionante o estimulante nasal oral.

No exceda la dosis recomendada.

- Suspenda su uso y consulte a su médico si
 - se siente nervioso, mareado o con insomnio
 - los síntomas no mejoran en los siguientes 7 días o se acompañan de fiebre
 - la tos persiste por más de 7 días, regresa, o se presenta acompañada por fiebre, erupción o dolor de cabeza persistente. Estos podrían ser signos de una condición grave.

En caso de embarazo o lactancia, consulte a un profesional médico antes de su uso. Manténgase fuera del alcance de los niños. En caso de sobredosis, obtenga ayuda médica o contacte al Centro de Control de Envenenamiento de manera inmediata.

Indicaciones

- no tomar más de 6 dosis en un periodo de 24 horas
- este producto de concentración para adultos no está previsto para su uso en niños menores de 12 años de edad
- medir solo con la copa dosificadora incluida
- mantenga la copa dosificadora con este producto
- mL = mililitro

edad	dosis
adultos y niños de 12 años de edad y mayores	20 mL cada 4 horas
niños menores de 12 años	no usar

Otra información

- cada 20 mL contiene: sodio 10 mg
- almacenar entre 15 y 30°C (59 a 86°F)
- no refrigerar.

Ingredientes inactivos

ácido cítrico anhidro, edetato disódico, FD&C azul #1, FD&C rojo #40, sabores, glicerina, propilenglicol, galato de propilo, agua purificada, benzoato sódico, sorbitol, sucralosa, goma xantana.

¿Preguntas o comentarios?

1-800-680-5223 Lunes a Viernes 9am-4pm,
Hora del Este

Distributed by / Distribuido por:
The Menzer Group
11421 NW107th ST,
Miami, FL 33178, USA



EXP
LOT No./Lote:

MUCOSAN SEVERE MULTI-SYMPOM

dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53145-697
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SORBITOL (UNII: 506T60A25R)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53145-697-06	1 in 1 CARTON	11/01/2024	
1		177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/01/2024	

Labeler - Menper Distributors, Inc. (101947166)**Registrant** - Menper Distributors, Inc. (101947166)