

ASPIRIN 81MG DYE-FREE- aspirin 81mg tablet, film coated
Allegiant Health

457 - Aspirin 81mg Dye Free

Active ingredient(s)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Use(s)

for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you take a blood thinning (anticoagulant) or steroid drug

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breastfeeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- TAMPER EVIDENT DO NOT USE IF IMPRINTED

SAFETY SEAL UNDER CAP IS BROKEN OR
MISSING

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

colloidal anhydrous silica, corn starch, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions/Comments

Call 1-888-952-0050
Monday through Friday 9AM - 5PM

Principal Display Panel

HealthA2Z®
 NDC 69166-457-88
 "COMPARE TO BAYER" LOW DOSE ACTIVE INGREDIENT
ASPIRIN
 DYE FREE (NSAID)



ENTERIC COATED

Talk to your doctor or other healthcare provider before using this product for your heart.

81mg SAFETY COATED

LOW STRENGTH ASPIRIN REGIMEN



200 TABLETS
DELAYED RELEASE ASPIRIN

Drug Facts
Active ingredient (in each tablet)
 Aspirin 81 mg (NSAID)*Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses
 for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an

Drug Facts (continued on inside)

*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark BAYER® Low Dose Aspirin.



X0048SS6DJ

HealthA2Z® Aspirin ...s Headache | (NSAID)
 New



Distributed by:
 Allegiant Health
 Deer Park, NY 11729

LB2210
 R0524



Lot: _____
 Exp.: _____

UNVARNISH

Drug Facts (continued)

early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives ■ facial swelling ■ stroking
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug ■ are age 60 or older ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have had stomach ulcers or bleeding problems ■ take more or for a longer time than directed ■ have 3 or more alcoholic drinks every day while using this product

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as

- heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following

- signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools ■ feel faint ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing ■ occurs ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more

Warnings (continued)

redness or swelling is a symptom occur signs of a serious condition.

breast-feeding, ask a health care provider if it is especially important to use aspirin at 20 weeks or more pregnancy unless definitely directed by your doctor because it may cause

in an unborn child or during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

do not take more than 4 tablets in a full glass of water with

children 12 years and over. ■ take every 4 hours not to exceed 4 tablets in 24 hours unless directed by your doctor

if 12 years: do not use unless directed by your doctor

Do not tamper with the cap. Do not use if the cap is broken or the cap is missing.

Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F). See USP Controlled Room Temperature. See USP Excursions. See USP Storage. See USP Stability. See USP Container Closure Integrity.

Ingredients

- aspirin
- colloidal silicon dioxide
- croscarmellose sodium
- dicalcium phosphate dihydrate
- hydroxypropyl methylcellulose
- hydroxypropyl methylcellulose K100
- hydroxypropyl methylcellulose K150
- hydroxypropyl methylcellulose K400
- hydroxypropyl methylcellulose K1000
- hydroxypropyl methylcellulose K1500
- hydroxypropyl methylcellulose K4000
- hydroxypropyl methylcellulose K10000
- hydroxypropyl methylcellulose K15000
- hydroxypropyl methylcellulose K40000
- hydroxypropyl methylcellulose K100000
- hydroxypropyl methylcellulose K150000
- hydroxypropyl methylcellulose K400000
- hydroxypropyl methylcellulose K1000000
- hydroxypropyl methylcellulose K1500000
- hydroxypropyl methylcellulose K4000000
- hydroxypropyl methylcellulose K10000000
- hydroxypropyl methylcellulose K15000000
- hydroxypropyl methylcellulose K40000000
- hydroxypropyl methylcellulose K100000000
- hydroxypropyl methylcellulose K150000000
- hydroxypropyl methylcellulose K400000000
- hydroxypropyl methylcellulose K1000000000
- hydroxypropyl methylcellulose K1500000000
- hydroxypropyl methylcellulose K4000000000

Other information

See important information about this medicine on the inside of the cap.

© 2015 HealthA2Z, Inc. All rights reserved. HealthA2Z, the HealthA2Z logo, and UNVARNISH are trademarks of HealthA2Z, Inc. All other trademarks are the property of their respective owners.

Drug Facts
 ■ than 3 days
 ■ present
 ■ These could be
If pregnant or
 professional be
 important not to
 later in pregnan
 to do so by a do
 problems in the
 complications d
 reach of childr
 medical help or
 Center (1-800-2
Directions
 directed ■ dr
 each dose
 ■ adults and ch
 take 4 to 8 tabl
 exceed 48 tabl
 directed by a dc
 ■ children unde
 directed by a dc
Other info
 EVIDENT DO N
 SAFETY SEAL I
 MISSING ■ st
 permitted betw
 ■ use by expira
Inactive ing
 anhydrous silica
 methacrylic acid
 polydextrose, p
 wax, simethico
 sodium lauryl s
 triacetin, methyl
Questions
 Call 1-888-952-
 Monday through

Aspirin

ASPIRIN 81MG DYE-FREE

aspirin 81mg tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69168-457
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69168-457-98	200 in 1 BOTTLE; Type 0: Not a Combination Product	05/09/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/09/2024	

Labeler - Allegiant Health (079501930)

Revised: 5/2024

Allegiant Health