

BURKHART- sodium fluoride gel
Burkhart Dental Supply Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Active Ingredient:

Fluoride Ion 1.23%

Available from 2.09% Sodium Fluoride and Hydrofluoric Acid

Purpose:

Fluoride Treatment Gel

Indications and Usage:

- A stable thixotropic fluoride treatment gel used to help prevent dental decay.
- For Professional Use Only. This product is not intended for home or unsupervised use.

Warnings:

- Keep out of reach of children.
- Do not swallow. If product is accidentally swallowed in quantities greater than would normally occur with a treatment gel, seek medical help or contact a Poison Control Center right away.
- Read directions carefully before using.

Dosage and Administration:

Shake well before use. This is a one minute or four minute fluoride gel for in-office patient use. It is normally used as a preventative caries treatment two times a year.

1. After thorough prophylaxis, fill two single or one dual tray, one third full with gel. Air dry teeth and insert trays into the mouth.
2. Instruct patient to bite down lightly but firmly for one minute (80% effectiveness) or four minutes (100% effectiveness).
3. Remove trays. Instruct patient to expectorate any excess gel and not to eat or drink for at least 30 minutes.

Other Information:

Store at controlled room temperature 59° to 86°F (15°-30°C).

Protect from freezing.

Inactive Ingredients:

Citric Acid, FD&C Red #3, FD&C Blue #1, Flavor, Magnesium Aluminum Silicate, Phosphoric Acid, Polysorbate 20, Sodium Benzoate, Sodium Saccharin, Titanium Dioxide, Water, Xanthan Gum, Xylitol.

APF GEL

**Acidulated Phosphate
Fluoride Treatment Gel
with Xylitol**

1.23% Fluoride Ion

Gluten Free

R_x Only

GRAPE



Reorder No. 72721422



Active Ingredient:

Fluoride Ion 1.23%.....Fluoride Treatment Gel Available from 2.09% Sodium Fluoride and Hydrofluoric Acid.

Purpose:

Indications and Usage:

- A stable thixotropic fluoride treatment gel used to help prevent dental decay.
- For Professional Office Use Only. This product is not intended for home or unsupervised consumer use.

Warnings:

- Keep out of reach of children.
- Do not swallow. If product is accidentally swallowed in quantities greater than would normally occur with a treatment gel, seek medical help or contact a Poison Control Center right away.
- Read directions carefully before using.

Dosage and Administration:

Shake well before use. This is a one minute or four minute fluoride gel for in-office patient use. It is normally used as a preventative caries treatment two times a year.

1. After thorough prophylaxis, fill two single or one dual tray, one third full with gel. Air dry teeth and insert trays into the mouth.
2. Instruct patient to bite down lightly but firmly for one minute (80% effectiveness) or four minutes (100% effectiveness).
3. Remove trays. Instruct patient to expectorate any excess gel and not to eat or drink for at least 30 minutes.

Other Information:

- Store at controlled room temperature 59° to 86°F (15°-30°C). Protect from freezing.

Inactive Ingredients:

Citric Acid, FD&C Blue #1, FD&C Red #3, Flavor, Magnesium Aluminum Silicate, Phosphoric Acid, Polysorbate 20, Sodium Benzoate, Sodium Saccharin, Titanium Dioxide, Water, Xanthan Gum, Xylitol.

Manufactured for Burkhart Dental Supply
2502 S. 78th Street
Tacoma, Washington 98409
(800) 562-8176
www.burkhartdental.com

Net wt. 16 oz (454 g)

MADE IN THE USA

99092605 Rev 9/13

BURKHART

sodium fluoride gel

Product Information

Product Type

HUMAN PRESCRIPTION DRUG

Item Code (Source)

NDC:43498-102

Route of Administration	DENTAL
--------------------------------	--------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM FLUORIDE (UNII: 8ZYQ1474W7) (FLUORIDE ION - UNII:Q80VPU408O)	FLUORIDE ION	5.6 g in 454 g

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	
PHOSPHORIC ACID (UNII: E4GA8884NN)	
POLYSORBATE 20 (UNII: 7T1F30V5YH)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
WATER (UNII: 059QF0KO0R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
XYLITOL (UNII: VCQ006KQ1E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	purple	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43498-102-15	454 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/01/2013	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/14/2012	

Labeler - Burkhart Dental Supply Inc (027532357)