WHITENING- yayashi whitening gel, dentifrice Guangzhou Baishili Trading Co., Ltd

Emphasize that the toothpaste uses mild whitening ingredients that do not damage tooth enamel, is suitable for daily use, and helps users easily achieve teeth whitening effects

Warnings:

For external use only: Do not ingest the toothpaste. If swallowed, seek medical help immediately.

Avoid use on damaged or sensitive gums: If you have sensitive teeth, gums, or any oral health concerns, consult with a dentist before using this product.

Possible sensitivity: Some individuals may experience mild tooth or gum sensitivity with whitening products. Discontinue use if sensitivity persists or worsens, and consult a dentist if necessary.

Keep out of reach of children: This product is intended for adults and should be stored out of reach of children. Children under 12 years of age should only use this product with the guidance of a dentist.

Avoid contact with eyes: In case of accidental contact with eyes, rinse thoroughly with water. Seek medical attention if irritation continues.

Example: "For external use only. Do not swallow. Consult a dentist if you have sensitive teeth or gums. Discontinue use if sensitivity occurs. Keep out of reach of children and avoid contact with eyes."

Indications:

This product is intended for individuals looking to remove surface stains and achieve a natural whitening effect.

Suitable for adults experiencing mild tooth discoloration caused by coffee, tea, red wine, or smoking.

Not recommended for individuals with sensitive teeth or gums. Please consult a dentist before use if you have dental concerns.

Usage:

Use twice daily, morning and night. Apply a pea-sized amount of toothpaste to your toothbrush.

Gently brush teeth for at least two minutes, ensuring the toothpaste covers all tooth surfaces.

Rinse thoroughly after brushing; avoid swallowing.

For optimal whitening results, continue use for at least two weeks, and limit consumption of stain-causing foods and drinks.

Stop Use:

If irritation or discomfort occurs: Discontinue use immediately if you experience persistent irritation, discomfort, or unusual sensitivity in teeth or gums.

If symptoms worsen: Stop using the product if symptoms such as pain, inflammation,

or tooth sensitivity intensify. Consult a dentist before resuming use.

If allergic reaction develops: Discontinue use if you notice signs of an allergic reaction, such as redness, itching, or swelling in the mouth or on the gums.

Here's how the KEEP OUT OF REACH OF CHILDREN section could be presented for a whitening toothpaste:

Keep Out of Reach of Children:

This product is intended for adult use only. Do not allow children to use without supervision or the guidance of a dentist.

In case of accidental ingestion by a child, seek immediate medical assistance or contact a Poison Control Center.

Example: "Keep out of reach of children. For adult use only. If swallowed by a child, get medical help or contact a Poison Control Center immediately."

Here's how the DOSAGE & ADMINISTRATION section for a whitening toothpaste could be structured:

Dosage & Administration:

Recommended Dosage: Use a pea-sized amount of toothpaste per brushing session.

Frequency of Use: Brush teeth twice daily, in the morning and at night, or as directed by a dentist.

Administration Instructions:

Apply a pea-sized amount of toothpaste to a toothbrush. Brush teeth gently for at least two minutes, covering all tooth surfaces. Rinse thoroughly after brushing and avoid swallowing the product.

Duration for Optimal Results: For best whitening results, use consistently for a minimum of two weeks. Consult a dentist if additional whitening is desired or if you plan to use for an extended period.

Example: "Apply a pea-sized amount and brush twice daily for two minutes, ensuring full coverage. Rinse thoroughly after use. For optimal whitening, continue use for at least two weeks."

LYSOZYME SODIUM PHYTATE SORBITOL WATER SACCHARIN SODIUM SODIUM LAURYL SULFATE HYDRATED SILICA

XANTHAN GUM

GLYCERIN



WHITENING

yayashi whitening gel, dentifrice

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:84727-051

| Ac | ctive Ingredier | nt/Active | Moiety | | | | |
|--|---|---|--|-------------------|--------------------|---|--------------|
| | | Ingre | dient Name | | Basis of | Strength | Strength |
| ΗY | DRATED SILICA (U | NII: Y6O7T4G | 8P9) (HYDRATED SILICA - UNII:Y | 607T4G8P9) | HYDRATED | SILICA | 0.5 g in 0.5 |
| GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX) | | | | GLYCERIN 0.5 g ii | | 0.5 g in 0.5 | |
| XA | NTHAN GUM (UNII: | TTV12P4NEE |) (XANTHAN GUM - UNII:TTV12P4 | NEE) | XANTHAN G | SUM | 0.5 g in 0.5 |
| In | active Ingredi | ients | | | | | |
| | | I | ngredient Name | | | Str | ength |
| LY | SOZYME (UNII: 968 | JKA7T33) | | | 0.5 g in 0.5 g | | |
| SODIUM PHYTATE (UNII: 88496G1ERL) | | | | 0.5 g in 0.5 g | | | |
| SORBITOL (UNII: 506T60A25R) | | | | | 0.5 g in 0.5 g | | |
| so | RBITOL (UNII: 5067 | F60A25R) | | | | 0.5 g in 0.5 | i g |
| | ATER (UNII: 5067 | | | | | 0.5 g in 0.5 | i g |
| W/ SA | ATER (UNII: 059QF0 CCHARIN SODIUM | KOOR) (UNII: SB8ZI | | | | 0.5 g in 0.5 0.5 g in 0.5 | i g i g |
| W/ SA | ATER (UNII: 059QF0 | KOOR) (UNII: SB8ZI | | | | 0.5 g in 0.5 | i g i g |
| W/ SA SO | ATER (UNII: 059QF0 CCHARIN SODIUM | KOOR) (UNII: SB8ZI | | | | 0.5 g in 0.5 0.5 g in 0.5 | i g i g |
| w SA SO | ATER (UNII: 059QF0 CCHARIN SODIUM DIUM LAURYL SUI | KO0R) (UNII: SB8Z1 L FATE (UNII: | | | | 0.5 g in 0.5 0.5 g in 0.5 0.5 g in 0.5 | i g i g |
| w/ sa so Pa | ATER (UNII: 059QF0 CCHARIN SODIUM DOUM LAURYL SU ACKaging Item Code NDC:84727-051- 10 | KOOR) (UNII: SB8Z) LFATE (UNII: Pac | 368GB5141J) | | ting Start Date | 0.5 g in 0.5 0.5 g in 0.5 0.5 g in 0.5 | eting End |
| WASASO | ATER (UNII: 059QF0 CCHARIN SODIUM DIUM LAURYL SU ACKAGING Item Code NDC:84727-051- 01 | KOOR) (UNII: SB8ZI LFATE (UNII: Pac 00 g in 1 TUE roduct | 368GB5141J) kage Description E; Type 0: Not a Combination | C | ting Start Date | 0.5 g in 0.5 0.5 g in 0.5 0.5 g in 0.5 | eting End |
| WASASO | ATER (UNII: 059QF0 CCHARIN SODIUM DOUM LAURYL SU ACKaging Item Code NDC:84727-051- 10 | KOOR) (UNII: SB8ZI LFATE (UNII: Pac 00 g in 1 TUE roduct | 368GB5141J) kage Description E; Type 0: Not a Combination | L 11/01/2024 | ting Start Date | 0.5 g in 0.5 0.5 g in 0.5 0.5 g in 0.5 Mark | eting End |

Labeler - Guangzhou Baishili Trading Co., Ltd (709042161)

| Establishment | | | | | | | |
|-------------------------------------|---------|-----------|----------------------------|--|--|--|--|
| Name | Address | ID/FEI | Business Operations | | | | |
| Guangzhou Baishili Trading Co., Ltd | | 709042161 | manufacture(84727-051) | | | | |

Revised: 11/2024

Guangzhou Baishili Trading Co., Ltd