

**DR. SCHOLLS BLISTER PAIN RELIEF COMPLETE- lidocaine**  
**Scholl's Wellness Company LLC**

-----  
**Dr. Scholl's Blister Pain Relief Complete Kit**

***Drug Facts***

***Active ingredient***

Lidocaine 4%

***Purpose***

External Analgesic

***Use***

Temporarily relieves pain

***Warnings***

**For external use only**

- Avoid contact with eyes
- If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.

**Do not use**

- More than one patch on your body at a time
- If you have had an allergic reaction to lidocaine or other local anesthetics
- Do not use in large quantities, particularly over raw surfaces or blistered areas
- For more than one week without consulting a doctor.

**When using this product**

- Use only as directed. Read and follow all directions and warnings on this carton

**Stop use and ask a doctor if**

- condition worsens
- severe burning sensation, redness, rash or irritation develops
- symptoms persist for more than 7 days or clear up and occur again within a few days
- you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.

**If pregnant or breast-feeding,**

ask a health care professional before use.

## **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

### ***Directions***

#### **Adults and children over 12 years:**

**Step 1 (Lidocaine patch application)** • apply to the affected area not more than 3-4 times daily

- clean and dry affected area
- remove backing from lidocaine patch by firmly grasping both ends and gently pulling away backing
- once exposed portion of patch is positioned, carefully remove remaining backing completely to apply patch to affected area
- once applied, press firmly on patch to help adhesive adhere to blister
- do not remove patch and reapply
- once patch is applied, move to Step 2.

**Step 2 (Bandage application)** • remove backing film from hydrogel bandage marked "Step 1" and place center (adhesive side down) over skin to seal lidocaine patch and blister.

- remove backing film marked "Step 2" and firmly smooth down adhesive edge to skin to seal lidocaine patch and blister. Replace when bandage begins to loosen

**children 12 years or younger:** ask a doctor

### ***Other information***

Store between 20-25°C (68-77°F)

### ***Inactive ingredients***

Cellulose Gum, Dihydroxyaluminum Aminoacetate, Ethylhexylglycerin, Glycerol, Laurocapram, PEG-40 Hydrogenated Castor Oil, Phenoxyethanol, Polyacrylic Acid, Propylene Glycol, Silica Silylate, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Tocopherol Acetate, Vanillyl Butyl Ether, Urea, Water (Aqua)

### ***Questions?***

**1-866-360-3226**

### **Package Labeling:**

B1-CM1-20097-1-24.10.9

NEW

**Drug Facts**

**Active ingredient**  
Lidocaine 4% **Purpose**  
External Analgesic

**Use** Temporarily relieves pain

**Warnings**

**For external use only**

■ Avoid contact with eyes ■ If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.

**Do not use**

■ More than one patch on your body at a time ■ If you have had an allergic reaction to lidocaine or other local anesthetics ■ Do not use in large quantities, particularly over raw surfaces or blistered areas ■ For more than one week without consulting a doctor.

**When using this product**

■ Use only as directed. Read and follow all directions and warnings on this carton

**Stop use and ask a doctor if** ■ condition worsens ■ severe burning sensation, redness, rash or irritation develops ■ symptoms persist for more than 7 days or clear up and occur again within a few days ■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.

**If pregnant or breast feeding**, ask a health care professional before use.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** Adults and children over 12 years:

**Step 1 (Lidocaine patch application)** ■ apply to the affected area not more than 3-4 times daily ■ clean and dry affected area ■ remove backing from lidocaine patch by firmly grasping both ends and gently pulling away backing ■ once exposed portion of patch is positioned, carefully remove remaining backing completely to apply patch to affected area ■ once applied, press firmly on patch to help adhesive adhere to blister ■ do not remove patch and reapply ■ once patch is applied, move to Step 2.

**Step 2 (Bandage application)** ■ remove backing film from hydrogel bandage marked "Step 1" and place center (adhesive side down) over skin to seal lidocaine patch and blister. ■ remove backing film marked "Step 2" and firmly smooth down adhesive edge to skin to seal lidocaine patch and blister. Replace when bandage begins to loosen

**Children 12 years or younger:** ask a doctor

**Other information** Store between 20-25°C (68-77°F)

**Inactive ingredients** Cellulose Gum, Dihydroxyaluminum Aminoacetate, Ethylhexylglycerin, Glycerol, Laurecapram, PEG-40 Hydrogenated Castor Oil, Phenoxyethanol, Polyacrylic Acid, Propylene Glycol, Silica Silylate, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Tocopherol Acetate, Vanillyl Butyl Ether, Urea, Water (Aqua/Eau)

**Questions?** 1-866-360-3226

**FSA/HSA ELIGIBLE**

For additional information and instructions visit: [drscholls.com](http://drscholls.com)  
Pat. patents.drscholls.com

**INSTRUCTIONS FOR USE:**

**Lidocaine Patch Application**  
Clean and dry affected area. Remove backing from lidocaine patch and apply to blister. Press firmly on patch to help adhesive adhere to blister. Move to hydrogel bandage application.



**Hydrogel Bandage Application**  
**Step 1:** Remove backing film from hydrogel bandage marked "Step 1" and place center (adhesive side down) over blister.



**Step 2:** Remove backing film marked "Step 2" and firmly smooth down adhesive edge to skin to seal lidocaine patch and blister. Replace when strip begins to loosen.



**BLISTER PAIN RELIEF COMPLETE KIT**  
**SEAL & HEAL BANDAGES**

**LIDOCAINE 4%**

**CLINICALLY TESTED 12-HOUR PAIN RELIEF**

- ✓ Lidocaine patch numbs pain
- ✓ Hydrogel bandage seals & cushions blister



**BLISTER PAIN RELIEF COMPLETE KIT**  
**SEAL & HEAL BANDAGES**

- ✓ Maximum strength lidocaine without a prescription
- ✓ Only blister treatment with an active pain reliever


©/© Scholl's Wellness Company 2024  
Dist. by: Scholl's Wellness Company LLC, Parsippany, NJ 07054  
Made in China.



彩面



Lot: S269A



# DR. SCHOLL'S BLISTER PAIN RELIEF COMPLETE

lidocaine kit

| Product Information |                |                           |                |  |
|---------------------|----------------|---------------------------|----------------|--|
| <b>Product Type</b> | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:73469-0487 |  |

| Packaging |                  |                     |                      |                    |
|-----------|------------------|---------------------|----------------------|--------------------|
| #         | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
| 1         | NDC:73469-0487-8 | 1 in 1 KIT          | 02/15/2025           |                    |

| Quantity of Parts |                  |       |                        |  |
|-------------------|------------------|-------|------------------------|--|
| Part #            | Package Quantity |       | Total Product Quantity |  |
| Part 1            | 8                | PATCH | 2.96 g                 |  |

## Part 1 of 1

### DR SCHOLL S 4 LIDOCAINE EXTERNAL ANALGESIC

lidocaine patch

#### Product Information

Item Code (Source) NDC:75568-024

Route of Administration TOPICAL

#### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength     |
|--|-------------------|--------------|
| LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987) | LIDOCAINE         | 40 mg in 1 g |

#### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| CELLULOSE GUM (UNII: K679OBS311)                       |          |
| DIHYDROXYALUMINUM AMINOACETATE (UNII: DO250MG0W6)      |          |
| ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)                  |          |
| GLYCERIN (UNII: PDC6A3C0OX)                            |          |
| LAUROCAPRAM (UNII: 1F3X9DRV9X)                         |          |
| POLYOXYL 40 HYDROGENATED CASTOR OIL (UNII: 7YC686GQ8F) |          |
| PHENOXYETHANOL (UNII: HIE492ZZ3T)                      |          |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)                    |          |
| TARTARIC ACID (UNII: W4888I119H)                       |          |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)                    |          |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)          |          |
| VANILLYL BUTYL ETHER (UNII: S2ULN37C9R)                |          |
| UREA (UNII: 8W8T17847W)                                |          |
| WATER (UNII: 059QF0KO0R)                               |          |

#### Packaging

| # | Item Code        | Package Description                                  | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:75568-024-01 | 1 in 1 POUCH   |                      |                    |
| 1 |                  | 0.37 g in 1 PATCH; Type 0: Not a Combination Product |                      |                    |

#### Marketing Information

| Marketing | Application Number or Monograph | Marketing Start | Marketing End |
|-----------|---------------------------------|-----------------|---------------|
|-----------|---------------------------------|-----------------|---------------|

| Category           | Citation | Date       | Date |
|--------------------|----------|------------|------|
| OTC Monograph Drug | M017     | 02/15/2025 |      |

| Marketing Information |  |                      |                    |
|-----------------------|--|----------------------|--------------------|
| Marketing Category    | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug    | M017                                     | 02/15/2025           |                    |

**Labeler** - Scholl's Wellness Company LLC (117174744)

| Establishment                     |         |           |                             |
|-----------------------------------|---------|-----------|-----------------------------|
| Name                              | Address | ID/FEI    | Business Operations         |
| Shanghai Sibeï Cosmetic Co., Ltd. |         | 542969967 | pack(75568-024, 73469-0487) |

Revised: 12/2024

Scholl's Wellness Company LLC