

**CABINET PAIN RELIEVER AND FEVER REDUCER REFILL ACETAMINOPHEN-  
acetaminophen tablet  
Cabinet Health P.B.C.**

-----  
**Cabinet Pain Reliever & Fever Reducer Refill, Acetaminophen**

***Drug Facts***

***Active Ingredient (in each caplet)***

Acetaminophen 500 mg

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to: • the common cold • headache • backache • minor pain of arthritis • toothache • muscular aches • premenstrual and menstrual cramps
- temporarily reduces fever

***Warnings***

**Liver warning:**This product contains acetaminophen. Severe liver damage may occur if you take • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Allergy alert:**Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away

***Do not use***

- with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

***Ask a doctor before use if you have***

liver disease

***Ask a doctor or pharmacist before use if you are***

taking the blood thinning drug warfarin

***Stop use and ask a doctor if***

• pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions • do not take more than directed (see overdose warning)**

---

adults & children 12 years & over • take 2 caplets every 6 hours while symptoms last • do not take more than 6 caplets in 24 hours, unless directed by a doctor • do not use for more than 10 days unless directed by a doctor  
children under 12 years ask a doctor

---

**Other Information**

• store between 20–25°C (68–77°F)


**Inactive ingredients**

hydroxypropylmethylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, stearic acid, talc, titanium dioxide

**Questions or comments?**

**908-242-6108**

**Package Labeling:**



# CABINET:

## Pain Reliever & Fever Reducer Refill

COMPARES TO THE ACTIVE INGREDIENT IN  
**Extra Strength Tylenol®\***

**Acetaminophen 500 mg**

RELIEVES:  
HEADACHE • FEVER • MUSCLE ACHES &  
PAINS • MENSTRUAL CRAMPS

**120**  
CAPLETS

THIS PACKAGE FOR  
HOUSEHOLDS WITHOUT  
YOUNG CHILDREN



**TAMPER EVIDENT: DO NOT USE IF POUCH IS OPEN**

**Drug Facts**

**Drug Facts** (continued)

**Active Ingredient (in each caplet) Purpose**

Acetaminophen 500 mg..... Pain reliever/fever reducer

**Uses** • temporarily relieves minor aches and pains due to: • the common cold • headache • backache • minor pain of arthritis • toothache • muscular aches • premenstrual and menstrual cramps • temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away

**Do not use** • with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if** • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** • do not take more than directed (see overdose warning)

adults & children 12 years & over	• take 2 caplets every 6 hours while symptoms last • do not take more than 6 caplets in 24 hours, unless directed by a doctor • do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

**Other Information**

• store between 20-25°C (68-77°F)

**Inactive ingredients** hypermellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, stearic acid, talc, titanium dioxide

**Questions or comments?**  
908-242-6108

**PACKAGE NOT CHILD-RESISTANT**

**Plastic bottles were a headache of the past.**

\* This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Extra Strength Tylenol®

† Less than the limit of detection and consistent with gluten-free diet labeling per FDA



NDC: 82725-3027-2

WKD-4-10027  
Distributed by: Cabinet Health P.B.C.  
Question/Comments? 1908-242-6108  
www.cabinethealth.com

Lot / Expiration ▼



Independently Quality Tested



Gluten-Free Verified†



Packaging with this logo will turn into dirt in lomi.

\*\* Compost with an at-home composting device or at your nearest industrial composting facility. Appropriate facilities may not exist in your area.

**CABINET PAIN RELIEVER AND FEVER REDUCER REFILL ACETAMINOPHEN**

acetaminophen tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:82725-3027
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	S53
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:82725-3027-2	120 in 1 POUCH; Type 0: Not a Combination Product	05/12/2025	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	05/12/2025	

**Labeler** - Cabinet Health P.B.C. (117102391)

Revised: 10/2025

Cabinet Health P.B.C.