ASPERCREME- lidocaine patch Lead Chemical Co. Ltd.

Aspercreme Lidocaine Patch

Active ingredient - Lidocaine Patch US 1ct

Lidocaine 4%

Purpose - Lidocaine Patch US 1ct

Topical anesthetic

Use - Lidocaine Patch US 1ct

for the temporary relief of pain

Warnings - Lidocaine Patch US 1ct

For external use only

Do not use - Lidocaine Patch US 1ct

- on wounds or on irritated or damage skin
- more than 1 patch on your body at a time or with other topical analgesics at the same time
- · with a heating pad

When using this product - Lidocaine Patch US 1ct

- use only as directed
- do not bandage tightly
- avoid contact with the eyes and mucous membranes
- do not expose the area to local heat or to direct sunlight

Stop use and ask a doctor if - Lidocaine Patch US 1ct

- condition worsens or symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days
- redness is present or excessive skin irritation occurs
- you experience severe burning pain, swelling, or blistering where the product was applied

If pregnant or breast-feeding, - Lidocaine Patch US 1ct

ask a health professional before use.

Keep out of reach of children and pets. - Lidocaine Patch US 1ct

Used patches may still contain the drug product that can produce serious adverse effects if ingested by children or pets. If swallowed, get medical help or contact a Poison Control Center right away.

Directions - Lidocaine Patch US 1ct

- adults and children 12 years of age and older:
 - clean and dry affected area
 - remove backing from patch by firmly grasping both ends and gently pulling until backing separates in middle
 - carefully remove smaller portion of backing from patch and apply exposed portion of patch to affected area
 - once exposed portion of patch is positioned, carefully remove remaining backing to completely apply patch to affected area
 - apply to the affected area not more than 3 to 4 times daily
 - wash hands after applying or removing patch. Throw away the patch after folding sticky ends together.
- children under 12 years of age: ask a doctor

Inactive ingredients - Lidocaine Patch US 1ct

aluminum glycinate, aluminum hydroxide, cellulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, methylparaben, nonoxynol-30, polyacrylic acid, polysorbate 80, propylene glycol, silica, sodium polyacrylate, tartaric acid, titanium dioxide, urea, water

Active ingredient - Lidocaine Patch Singles US 1ct

Lidocaine 4%

Purpose - Lidocaine Patch Singles US 1ct

Topical anesthetic

Use - Lidocaine Patch Singles US 1ct

For the temporarily relief of pain

Warnings - Lidocaine Patch Singles US 1ct For external use only

Do not use - Lidocaine Patch Singles US 1ct

- on puncture wounds, cuts, irritated or swollen skin
- more than 1 patch on your body at a time or with other topical analgesics at the

- same time
- with a heating pad or apply local heat to the area of use

When using this product - Lidocaine Patch Singles US 1ct

- use only as directed
- do not bandage tightly
- avoid contact with the eyes
- dispose of used patch in manner that always keeps product away from children or pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and ask a doctor if - Lidocaine Patch Singles US 1ct

- condition worsens or symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days
- redness or irritation develops
- you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

If pregnant or breast feeding, - Lidocaine Patch Singles US 1ct ask a health professional before use.

Keep out of reach of children and pets. - Lidocaine Patch Singles US 1ctIf swallowed, get medical help or contact a Poison Control Center right away.

Directions - Lidocaine Patch Singles US 1ct adults and children over 12 years of age and older:

- clean and dry affected area
- remove backing from patch by firmly grasping both ends and gently pulling until backing separates in middle
- carefully remove smaller portion of backing from patch and apply exposed portion of patch to affected area
- once exposed portion of patch is positioned, carefully remove remaining backing to completely apply patch to affected area
- use 1 patch at a time and not more than 3 to 4 times daily

children under 12 years of age: consult a doctor

Inactive ingredients - Lidocaine Patch Singles US 1ct

aluminum glycinate, aluminum hydroxide, cellulose gum, glycerin, methyl acrylate/2-ethylhexyl acrylate copolymer, methylparaben, nonoxynol-30, polyacrylic acid, polysorbate 80, propylene glycol, silica, sodium polyacrylate, tartaric acid, titanium dioxide, urea, water

Package/Label Principal Display Panel - Lidocaine Patch US 1ct MAX STRENGTH

Aspercreme® with 4% LIDOCAINE PAIN RELIEF PATCH

NUMBS AWAY PAINFAST ACTING TARGETS NERVES FLEXIBLE FABRIC #1 LIDOCAINE BRAND*

PATCH

FRAGRANCE-FREE
Contains 1 Patch 3-15/16" x 5-1/2" (10 cm x 14 cm)
Pain Relieving Ointment on a Breathable Adhesive Pad

*Among OTC topical analgesics.

Dist. by Chattem, Inc., a Sanofi Company P.O. Box 2219, Chattanooga, TN 37409-0219 USA © 2022 www.aspercreme.com Made in Japan



Label

Package/Label Principal Display Panel - Lidocaine Patch Singles US 1ct MAX STRENGTH

Aspercreme® with 4% LIDOCAINE PAIN RELIEF PATCH

NUMBS AWAY PAINFAST ACTING TARGETS NERVES FLEXIBLE FABRIC #1 LIDOCAINE BRAND*

PATCH

FRAGRANCE-FREE Contains 1 Patch 3-15/16" x 5-1/2" (10 cm x 14 cm)

Pain Relieving Ointment on a Breathable Adhesive Pad

*Among OTC topical analgesics

CHATTEM®
A SANOFI COMPANY
Dist. by Chattem, Inc., a Sanofi Company
P.O. Box 2219, Chattanooga, TN
37409-0219 USA © 2020
www.aspercreme.com
Made in Japan



ASPERCREME

lidocaine patch

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:62168-0584

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	246 mg

Inactive Ingredients			
Ingredient Name	Strength		
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)			
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)			
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K6790BS311)			
GLYCERIN (UNII: PDC6A3C0OX)			
METHYLPARABEN (UNII: A2I8C7HI9T)			
NONOXYNOL-30 (UNII: JJX07DG188)			
POLYSORBATE 80 (UNII: 60ZP39ZG8H)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
TARTARIC ACID (UNII: W4888I119H)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
UREA (UNII: 8W8T17847W)			
WATER (UNII: 059QF0KO0R)			

Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:62168- 0584-4	400 in 1 CARTON	05/17/2021	03/31/2025	
1		1 in 1 POUCH; Type 0: Not a Combination Product			
2	NDC:62168- 0584-5	400 in 1 CARTON	02/02/2022		
2		1 in 1 POUCH; Type 0: Not a Combination Product			
3	NDC:62168- 0584-6	400 in 1 CARTON	07/15/2023		
3		1 in 1 POUCH; Type 0: Not a Combination Product			

Marketing II	nformation
--------------	------------

Marketing Application Number or Monograph Marketing Start Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	505G(a)(3)	02/16/2016	

Labeler - Lead Chemical Co. Ltd. (693727091)

Revised: 1/2024 Lead Chemical Co. Ltd.