SEFUDUN QUIT SMOKING NICOTINE PATCHES- nicotine patch Shenzhen Shandian Jingling Technology Co., Ltd.

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

SEFUDUN QUIT SMOKING NICOTINE PATCHES

nicotine

Stop smoking aid

Used patches have enough nicotine to poison children and petsif swallowed, get medical help or conlact a poison control centerfight away, Save pouch to use for patch disposal. Dispose af theused patches by folding sticky ends together and putting in pouch,

Use reduices withdrawal symploms, including nicotine craving associated with guitting smokind

if you are precnant or breast.feeding, only use this medicine onthe advice of your heaith care provider.Smoking can seribusly harm your child. Try to stop smoking withoutusing anvnicotine replacement medicine, This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known

if you are under 18 years of age, ask a doctor before use. Before usino this product, read the enclosed self-help guide forcomplete directions and other infolmation. Begin usio the patch on your guit day.

if you smoke more than 10 cigarettes per day, use the foilowingschedule below:

Weeks 1-4
Step 1
Use one 21 mgpatch / day
Weeks 5-6
Step 2
Use one 14 mgpatch / day
Weeks 7-8
Step 3
Use one 7 mgpatch day

methacrylic acid copolymer POLYESTER-5 (TG-38) ACRYLIC ACID SEFUDUN®



QUIT SMOKING AID

CONTROL CRAVINGS ALL-DAY!

- Helps You Quit Smoking.
- Controls Nicotine Cravings.
- **Olights Withdrawal Symptoms.**
- 0100% Natural Plant Based.

Step3



Drug Facts

Active ingredient (in each patch)

Purpose

MARKET SELL STREET, ST

Nicotine, 7 mg delivered over 24 hours

Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.

Smoking can seriously harm your child. Try to stop smoking without ng any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- Heart disease, recent heart attack, or irregular heartbeat.
 Nicotine can increase your heart rate.
- · High blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- An allergy to adhesive tape or have wkin problems, because you are more likely to get rashes.
- Stomach ulcer or diabetes.

Ask a doctor or pharmacist before use if you are:

When using this product

Stop use and ask a doctor if

- days, or if your skin swells, or you get a rash.
- irregular heartbeat or palpitations occur.
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat

Keep out of reach of children and pets

Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a poison control center

Drug Facts

Directions

- If you are under 18 years of age, ask a doctor before use
- Before using this product, read the enclosed self-help guide for complete directions and other information
- Begin using the patch on your quit day
- If you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1-4	Weeks 5-6	Weeks 7-8
Step 1	Step 2	Step 3
Use one 21 mg patch / day	Use one 14 mg patch / day	Use one 7 mg patch / day

- If you smoke 10 or less cigarettes per day, start with Step 2. for 6 weeks, then step 3 for 2 weeks.
- Apply one new patch every 24 hours on skin that is dry, clean
- Remove backing from patch and immediately press onto skill Hold for 10 seconds.
- Wash hands after applying or removing patch. Save pouch is use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- The used patch should be removed and a new one applied.
- and apply a new one in the morning.
- Do not wear more than one patch at a time.
- Do not cut patch in half or into smaller pieces.
- irritate your skin and loses strength after 24 hours.
- To avoid possible burns, remove patch before undergoing arts. MRI (magnetic resonance imaging) procedures.
- health care provider.

Other information • store at 20-25 °C (68-77°F)

Inactive ingredients acrylate adhesive, aluminized polyester, cellulose paper, methacrylic acid copolymer.

Comments or Questions? call + 1305 767 1517 Weekdays (8am-4pm ET)

nicotine patch

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:83804-107

Route of Administration TRANS DERMAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (LINII) 6M3C807Y6P) (NICOTINE LINII) 6M3C807Y6P)	NICOTINE	7 ma

VICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) NICOTINE 7 mg

Inactive Ingredients

Ingredient Name	Strength

ACRYLIC ACID (UNII: J94PBK7X8S)

METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)

POLYESTER-5 (TG-38) (UNII: 2L9351NW8W)

Packaging

# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:83804-107-	30 in 1 BOX; Type 0: Not a Combination Product	11/22/2024	

Marketing Information

Marketing information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		11/22/2024	

Labeler - Shenzhen Shandian Jingling Technology Co., Ltd. (418636058)

Establishment				
Name	Address	ID/FEI	Business Operations	
Shenzhen Shandian Jingling Technology Co., Ltd.		418636058	manufacture(83804-107)	

Revised: 11/2024 Shenzhen Shandian Jingling Technology Co., Ltd.