

**SAFETUSSIN HIGH BLOOD PRESSURE AND DIABETES COUGH, COLD AND FLU-  
acetaminophen, dextromethorphan hydrobromide, and guaifenesin liquid  
KRAMER LABORATORIES**

-----  
**SAFETUSSIN High Blood Pressure Nighttime COLD & FLU**

***Drug Facts***

<b><i>Active ingredients (in each 30 mL dose cup)</i></b>	<b><i>Purpose</i></b>
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant

**Uses**

temporarily relieves common cold/flu symptoms:

- cough
- sore throat
- headache
- minor aches and pains
- fever
- helps loosen phlegm (mucus) and thin bronchial secretions to get rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning**

This product contains acetaminophen.  
Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions.  
Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

## **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- If you have ever had an allergic reaction to this product or any of its ingredients
- In children under 12 years of age

## **Ask a doctor before use if you have**

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

## **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

## **When using this product**

Do not use more than as directed

## **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

## **Directions**

- do not take more than directed

- do not take more than 6 doses in any 24-hours period
- measure only with dosing cup provided
- dose as follows or as directed by a doctor

adults & children 12 years & older	30 mL per dose every 4 hours
children under 12 years of age	do not use

### Other information

- **each 30 mL contains:** sodium 19 mg
- store at room temperature. Do not refrigerate.

### Inactive ingredients

Anhydrous citric acid, edetate disodium, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose, xanthan gum.

### Distributed by

### PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label

SAFE FOR ADULTS WITH HIGH BLOOD PRESSURE & DIABETES

NDC 55505-217-36

**Safetussin®**

**MAX STRENGTH**

**MULTI-SYMPTOM**

**Cough, Cold & Flu**

**Acetaminophen / Pain Reliever 650 mg**

Dextromethorphan HBr / Cough Suppressant 20 mg

Guaifenesin / Expectorant 400 mg

Safe for adults with

High Blood Pressure

Diabetes

Relieves

- ✓ Headache
- ✓ Fever
- ✓ Sore Throat
- ✓ Body Aches & Pains
- ✓ Chest Congestion
- ✓ Cough

Safetussin® contains:

No decongestants

No Sugar

No Alcohol

No sorbitol

No fructose

No gluten

No dyes

Pharmacist Recommended Brand

Always read and follow label directions. Do not use longer than 7 consecutive days unless directed by a doctor. Ask a doctor if you have individual health questions.

8 FL OZ (237 mL)

Questions? Comments? Call 1-800-824-4894

©2023 Kramer Laboratories, Inc. Bridgewater, NJ 08807



# SAFETUSSIN HIGH BLOOD PRESSURE AND DIABETES COUGH, COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and guaifenesin liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55505-217
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55505-217-36	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/17/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/17/2023	

**Labeler** - KRAMER LABORATORIES (122720675)

Revised: 12/2025

KRAMER LABORATORIES