

CHILDRENS LORATADINE- loratadine solution
Strategic Sourcing Services LLC

Children's Loratadine

Drug Facts

Active ingredient (in each 5 mL teaspoonful)

Loratadine 5 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- use only with enclosed dosing cup

adults and children 6 years and over | 2 teaspoonfuls (tsp) daily; do not take more than 2

years and over	teaspoonfuls (tsp) in 24 hours
children 2 to under 6 years of age	1 teaspoonful (tsp) daily; do not take more than 1 teaspoonful (tsp) in 24 hours
children under 2 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- **do not use if bottle wrap imprinted with "SEALED FOR SAFETY" is broken or missing.**
- see bottom panel for lot number and expiration date
- store between 20° and 25°C (68° and 77°F)

Inactive ingredients

artificial peach flavor, citric acid monohydrate, glycerin, propylene glycol, purified water, sodium benzoate, sodium metabisulfite, sucrose

Questions?

Call **1-866-923-4914**

Distributed by McKesson
One Post Street
San Francisco, CA 94104

PRINCIPAL DISPLAY PANEL - 120 mL Bottle Carton

NDC 62011-0305-1

Health Mart ®

Compare to
Children's Claritin ®
active ingredient*

FRUIT FLAVOR

Children's
Loratadine
Oral Solution
USP, 5 mg/5 mL
(Antihistamine) ALLERGY

non-drowsy †

24 hour relief of:

- Sneezing • Runny Nose
- Itchy, Watery Eyes

- Itchy Throat or Nose

indoor & outdoor allergies

Ages 2 yrs & older
dosing cup included
alcohol free

WARNING: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

†when taken as directed.
see drug facts panel.

4 FL OZ
(120 mL)



CHILDRENS LORATADINE

loratadine solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:62011-0305

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII:7AJ03BO7QN)	LORATADINE	5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	yellow (colorless to slightly yellow)	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0305-1	1 in 1 CARTON	11/15/2016	
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076805	08/20/2004	

Labeler - Strategic Sourcing Services LLC (116956644)

Establishment

Name	Address	ID/FEI	Business Operations
Taro Pharmaceutical Industries Ltd.		600072078	manufacture(62011-0305)