

INFANTS PAIN AND FEVER- acetaminophen suspension WALGREENS

655L Walgreens Acetaminophen 160 mg suspension Infants

Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed.

See overdose warning

■ **this product does not contain directions or complete warnings for adult use.**

■ **shake well before using**

■ mL = milliliter

■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

■ push air out of syringe. Insert syringe tip into the bottle opening

■ flip bottle upside down. Pull yellow part of syringe to correct dose

■ dispense liquid slowly into child's mouth, toward inner cheek

■ repeat dose every 4 hours while symptoms last

■ do not give more than 5 times in 24 hours

■ replace cap tightly to maintain child resistance

DOSING CHART

Weight (lbs)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

*or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other information

■ **each 5 mL contains:** sodium 2 mg

■ store between 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, FD&C red no. 40, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments?

Call **1-877-290-4008**

Front of Package

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each 5 mL)

Purpose

Acetaminophen 160 mg Pain reliever/fever reducer

Uses

- temporarily:
 - reduces fever
 - relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Side of Package

Drug Facts (continued)

Directions

Do not take more than directed.

See overdose warning

- this product does not contain directions or complete warnings for adult use.
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into the bottle opening
- flip bottle upside down. Pull yellow part of syringe to correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

DOSING CHART

Weight (lbs)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

*or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, FD&C red no. 40, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments?

Call 1-877-290-4008

*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

**This product is not manufactured or distributed by Kennebec Inc., owner of the registered trademark Tylenol®.

655L 1224 REV-0325

NDC 0363-6555-04

Back of Package

Walgreens

Compare to the active ingredient in Infants' Tylenol®**



Infants' Pain & Fever

ACETAMINOPHEN 160 mg PER 5 mL ORAL SUSPENSION
PAIN RELIEVER / FEVER REDUCER

Alcohol Free

Aspirin Free

Ibuprofen Free

AGES
2-3
YEARS



Cherry flavored

USE ONLY WITH ENCLOSED SYRINGE.
See side panel for more information.

2 FL OZ (60 mL)

Side of Package

TAMPEN TYLENOL DO NOT USE IF TAMPON SEAL AROUND THE BOTTLE IS BROKEN OR MISSING

STEP 1.
See *Directions* for the correct dose. Shake well before using.




STEP 2.
Push air out of the syringe.



STEP 3.
Insert syringe tip into bottle opening then flip bottle upside down.



STEP 4.
Pull yellow part of syringe to the 1st dose line then push product back into bottle.



STEP 5.
Pull yellow part of syringe until it reaches and stays at the correct dose.



STEP 6.
Dispense liquid slowly into child's mouth.



IMPORTANT:

Please see dosing chart or ask a doctor for your child's correct dose.

Walgreens

Infants' Pain & Fever

ACETAMINOPHEN 160 mg PER 5 mL ORAL SUSPENSION
PAIN RELIEVER / FEVER REDUCER

• USE ONLY WITH ENCLOSED SYRINGE
AGES 2-3 YEARS



2 FL OZ (60 mL) Cherry flavored

TAMPER EVIDENT: DO NOT USE IF SHRINK BAND SEAL AROUND THE BOTTLE IS BROKEN OR MISSING

Drug Facts

Active Ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose Pain relief/fever reducer

Uses temporarily: ■ reduces fever

■ relieves minor aches and pains due to:

■ the common cold ■ flu ■ headache

■ sore throat ■ toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more

than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen

■ Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness

■ hives ■ rash ■ itchy skin ■ If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If a sore throat is severe, persists for more than 2 days, is accompanied or followed by fever,

headache, rash, malaise, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure

whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or

any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning).

When using and ask a doctor if ■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3

days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison

Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children

even if you do not notice any signs or symptoms.

Directions Do not take more than directed. See overdose warning.

■ This product does not contain directions for adult use. ■ Shake well before using

■ mL = milliliter ■ Find right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ Push air

out of syringe. Push syringe tip into the bottle opening. ■ Tip bottle upside down. Pull yellow part of syringe to

correct dose. ■ Dispense liquid slowly into child's mouth. ■ Repeat dose every 4 hours while

symptoms last. ■ Do not give more than 5 times in 24 hours. ■ Replace cap tightly to maintain child resistance

to tampering.

■ Assistance is available to help you use this product. Call 1-800-368-6969.

■ Do not use if you notice any signs or symptoms of a serious condition.

■ Do not use if you notice any signs or symptoms of a serious condition.

REV-0326

Varnish Omit Area

LOT:

EXP:

PEEL HERE

DISTRIBUTED BY WALGREEN CO. DISTRIBUTION FOR: WALGREEN CO. DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED walgreens.com © 2025 Walgreen Co. MADE IN INDIA

ITEM 776788 W00000-0000-0 W3ORG0125-F 655L 1224 NDC 0363-6555-04

REV-0326

Varnish Omit Area

LOT:

EXP:

PEEL HERE

DOSE CHART

Weight (lbs)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

*Or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other Information ■ Each 5 mL contains sodium 2 mg ■ Store between 20-25°C (68-77°F)


Inactive Ingredients anhydrous citric acid, FD&C red no. 40, thaxox, glycerin, hydroxyethylcellulose,

hydroxyethylcellulose and carbomer, hydroxyethylcellulose, propylparaben, purified water, sodium benzoate,


excipients, saccharin, sucrose, xanthan gum

Questions or comments? Call 1-877-200-4008


IMPORTANT: Please see dosing chart or ask a doctor for your child's correct dose.




STEP 1.
See **Directions**
for the
correct dose.
Shake well
before using.



STEP 2.
Push air
out of the
syringe.



STEP 3.
Insert syringe
tip into bottle
opening then
flip bottle
upside down.



STEP 4.
Pull yellow
part of syringe
to the 1st dose
line then push
product back
into bottle.



STEP 5.
Pull yellow
part of syringe
until it reaches
and stays at
the correct dose.



STEP 6.
Dispense liquid
slowly into
child's mouth.



Walgreens Pharmacist Recommended.
Our pharmacists recommend the Walgreens
brand. We invite you to compare to
national brands.

Adhesive Area

INFANTS PAIN AND FEVER

acetaminophen suspension

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6555	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	160 mg in 5 mL	
Inactive Ingredients				
Ingredient Name			Strength	
GLYCERIN (UNII: PDC6A3C00X)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
SUCROSE (UNII: C151H8M554)				
SORBITOL SOLUTION (UNII: 8KW3E207O2)				
WATER (UNII: 059QF0KO0R)				
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)				
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
XANTHAN GUM (UNII: TTV12P4NEE)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
Product Characteristics				
Color	red	Score		
Shape		Size		
Flavor	CHERRY	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6555-04	1 in 1 CARTON	05/03/2025	
1		60 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	05/03/2025	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(0363-6555)

Revised: 3/2025

WALGREENS